A Novel HIV-1 Nef Mutation in a Primary Pediatric Isolate Impairs MHC-Class I Downregulation and Cytopathicity

Ayub Ali,^{1,2} Robert L. Furler,³ Livia Pedroza-Martins,⁴ Arnaud D. Colantonio,^{1,5} Deborah Anisman-Posner,^{1,5} Yvonne Bryson,^{1,6} Otto O. Yang,^{1,2} and Christel H. Uittenbogaart^{1,5-7}

Abstract

HIV-1-induced cytopathicity of thymocytes is a major cause of reduced peripheral T cells and rapid disease progression observed in HIV-1-infected infants. Understanding the virulence factors responsible for thymocyte depletion has paramount importance in addressing the pathogenesis of disease progression in children. In this study, thymocyte depletion was analyzed following infection with two primary CXCR4-tropic HIV-1 pediatric isolates (PI), PI-2 and PI-2.1, which were serially derived from an *in utero*-infected infant. Although highly similar to each other, PI-2 showed markedly decreased thymocyte depletion *in vitro* compared with PI-2.1. Further analysis showed a novel deletion in the Nef protein (Nef Δ K7S) of PI-2, which was absent in PI-2.1. This deletion inhibited Nef-mediated major histocompatibility complex class I (MHC-I) downregulation in infected thymocytes *in vitro* and *in vivo*; in contrast, the mutated Nef continued to downregulate CD4 surface expression *in vitro*. These results suggest that HIV-1 Nef contributes to thymic damage in infants through selective functions.

Keywords: HIV-1, Nef, MHC-I downregulation, cytopathicity, thymocyte

Introduction

THE THYMUS IS PARTICULARLY ACTIVE during early life as the immune system develops in infants. Thymocyte infection with HIV-1 leads to severe thymic involution and dysfunction, resulting in declined thymopoiesis and subsequent depletion of peripheral T lymphocytes.¹ Although direct killing of thymocytes has been suggested as a major mechanism responsible for thymic dysfunction,^{2,3} HIV-1 also causes nonspecific activation and proliferation of intrathymic progenitor cells through secretion of proinflammatory cytokines in the thymic microenvironment that result in elevated viral infection, exhaustion of precursor cells, and destruction of thymic architecture.^{4,5} These conditions within the thymus have been observed in children infected with HIV-1 and are suggested to be a leading cause of rapid disease progression in perinatally infected children.^{5,6} Studies to understand the underlying pathological and molecular aspects of thymus dysfunction in HIV-1 infection suggest that the Nef protein of

HIV-1 plays a key role in impairing thymocyte development in addition to interfering with peripheral T cell function.⁷⁻¹⁰

Nef is a phosphorylated and myristoylated 27 kDa protein that is unique to HIV/SIV and is relatively conserved with a 75%–85% similarity within clade B of HIV-1.¹¹ Nef is a multifunctional protein that is suggested to play multiple key roles in HIV-1 and SIV pathogenesis.^{8,12} Deletion of Nef markedly attenuates SIV pathogenicity in the rhesus macaque model of AIDS,^{8,13} and dysfunctional Nef has been associated with delayed disease progression in humans.¹⁴⁻¹⁶ Nef has also been shown to impair T cell development^{10,17} and enhance viral replication and infectivity.^{18,19} Its expression begins early in the viral life cycle and localizes to the plasma membrane, cytoplasm, and within the budding virions. Nef promotes viral infectivity, immune evasion, and replication capacity by altering the activation state of infected cells. It does so by modulating cell signal transduction pathways and altering cell surface expression of receptors. Nef regulates various cell signaling molecules, such as Hck, Lck, Fyn, Lyn,

¹UCLA AIDS Institute, University of California, Los Angeles, Los Angeles, California.

²Division of Infectious Diseases, Department of Medicine, University of California, Los Angeles, Los Angeles, California.

³Division of Infectious Diseases, Department of Medicine, Weill Cornell Medicine, New York, New York.

⁴ANRS, French National Agency for Research on AIDS and Viral Hepatitis, Paris, France.

⁵Department of Microbiology, Immunology & Molecular Genetics, University of California, Los Angeles, Los Angeles, California.

⁶Department of Pediatrics, University of California, Los Angeles, Los Angeles, California.

⁷Jonsson Comprehensive Cancer Center, University of California, Los Angeles, Los Angeles, California.

Vav, Src, PAK1/2 (NAK), MAPK (Erk1), c-Raf1 kinase, PI3K, and Ask1.^{20,21} Nef downregulates cell surface receptors such as CD4,^{22–24} major histocompatibility complex class I (MHC-I),^{23–25} MHC-II, and MHCII-associated invariant chain (Ii, CD74),²⁶ CD28,²⁷ and host restriction factor SERINC5.²⁸

Nef-mediated downregulation of surface expression of MHC-I and CD4 has been extensively studied *in vitro*. The downregulation of MHC-I A and B molecules protects infected cells from CTL-mediated killing.^{29–31} Previously, we described two closely related CXCR4-tropic HIV-1 primary isolates [pediatric isolates (PI)-2.1 and PI-2] serially derived from an *in utero*-infected infant, and their dissimilar abilities to cause thymocyte depletion.³² PI-2.1 showed aggressive thymocyte depletion *in vitro*, while isolate PI-2 showed diminished cytopathicity. Given the central role of Nef in HIV-1 disease progression, we performed a detailed analysis of Nef sequences of these two pediatric isolates and their corresponding abilities to downregulate MHC-I and CD4.

Materials and Methods

HIV-1 pediatric isolates and virus stocks

HIV-1 primary isolates and plasma samples for virus isolation were obtained as close as possible to birth from 10 of 34 children born to 204 HIV-1-seropositive mothers monitored prospectively in the Los Angeles Pediatric AIDS Consortium between June 1988 and May 1995.^{32–34} The isolates were characterized as syncytium inducing (SI) or nonsyncytium inducing (NSI) by using the MT-2 assay.³⁵ The characteristics of some of the children providing the viral isolates were previously described: PI-2 is an SI HIV-1 isolate obtained 1 week after birth from an infant infected *in utero* who developed AIDS and died within 6 months.³⁴ PI-2.1 was isolated from plasma 1 month after birth from the same infant. PI-2 and PI-2.1 isolation and propagation were done by directly adding serum from an infected infant to a pool of purified activated allogeneic CD4⁺ T cells, prepared as described previously.^{36,37} Briefly, allogeneic CD4⁺ T cells from three normal donors were individually purified by capture in CD4 monoclonal antibody-coated tissue culture flasks (Applied Immune Sciences, Santa Clara, CA) and activated by stimulation with 200 ng/mL anti-human CD3 antibody (OKT3; Ortho Biotech, Inc., Raritan, NJ) plus 5,000 U/mL recombinant human interleukin-2 (IL-2) for 5 days. Cells from three donors were combined, cryopreserved in liquid nitrogen, and then thawed and cultured in medium with IL-2 for 2–3 days before infection. The makeup by phenotype of the CD4⁺ T cell pool at the time of infection by the pediatric isolates was as follows: 95% CD4⁺ CD8[−], 5% CD4⁺ CD8⁺, <1% CD8⁺ CD4[−], 60% CD27⁺, and 50% HLA-DR⁺. In addition, 94% of the cells from the stimulated CD4⁺ T cell pool expressed CXCR4 and 30%–40% expressed CCR5 (data not shown).³² The same CD4⁺ T cell pool was used for isolating PI-2 and PI-2.1 virus from plasma (passage 1) and for preparing virus stocks (passage 2) and for preparing the control viruses including CCR5-tropic virus HIV-1 NF-NSX (NF-NSX) and the CXCR4-tropic virus HIV-1 NL4-3 (NL4-3).^{38,39} The infectivity of the pediatric isolates and of the molecular-clone-derived laboratory viruses was determined by limiting dilution assays in PBMC. The CCR5-tropic

pediatric isolate PI-8 was prepared similarly to PI-2 and PI-2.1 and used as control for the Western blot studies.³²

Cloning and sequencing of the nef gene in pediatric HIV-1 isolates

RNA was isolated from plasma using the TRIzol reagent (Invitrogen, Carlsbad, CA) and was reverse transcribed into complementary DNA (cDNA) using the High-Capacity cDNA Reverse Transcription Kits (Applied Biosystems, Foster City, CA). Reverse transcription was primed with a random primer (Applied Biosystems) and following nested polymerase chain reaction (PCR), the cDNA of full-length *nef* was amplified. First and second rounds of PCR were carried out using Nef 8687F (5-GTAGCTGAAGGGACAGATAGGGTTAT-3) and Nef 9589R (5-TAGTTAGCCAGAGAGCTCCCA-3) as the outer primer set and Nef 8787 XbaIF (5-GCTCTAGAATG GGTGGCAAGTGCTCAA-3) and Nef 9495R (5-TTATATG CAGCATCTGAGGGC-3) as the inner primer set, respectively. Numbering of primers is based on the NL4-3 sequence. High Fidelity Advantage[®] HD Polymerase Mix (Clontech, Mountain View, CA) was used for PCR, according to the manufacturer's instructions. After cloning of PCR products into the Topo TA cloning vector (Invitrogen), plasmid DNAs were isolated from at least six colonies for sequencing with the ABI 3130 Genetic Analyzer (Applied Biosystems).

Production of single-round infectious, pseudotyped HIV-1 reporter virus

The reporter HIV-1 construct AA1305#18 plasmid containing an NL4-3 backbone with reporter antigens heat shock antigen (HSA) or mouse CD24 (mCD24) and deletions in Vpu, Env, and Vpr was used to produce VSV-G pseudotyped virus as described previously.⁴⁰ Insertion of isolated *nef* genes was done using PCR products that were gel purified with Quick Spin Column (Qiagen, Valencia, CA) and digested with *Xba*I and *Bsp*EI restriction enzymes (New England Biolabs, Ipswich, MA). Single-round infectious pseudovirus was produced by cotransfecting 293T cells with the whole-genome plasmid AA1305#18 containing *nef* alleles and a plasmid encoding the envelope glycoprotein from VSV-G using BioT transfection reagent (Bioland, Paramount, CA).

Western blot

HIV-1 *nef* genes from the pediatric isolates (PI) were cloned into a plasmid containing the NL4-3 genome (pNL4-3) to make infectious virus as described previously.⁴¹ To construct NL4-3/PI-*nef*, the *nef* gene of NL4-3 was replaced with either individual *nef* obtained from Topo cloning or quasispecies or *nef* obtained from the nested PCR as described above. T1 cells were infected with NL4-3 virus containing *nef* alleles from either wild-type NL4-3 or alleles from the pediatric isolates (multiplicity of infection [MOI] = 1). On day 5, lysates harvested from the infected cells were run onto 10% polyacrylamide gel electrophoresis (PAGE) for protein separation. Following transfer to a PVDF membrane (Millipore, Billerica, MA), HIV-1 proteins were detected by an anti-Nef antibody (Nef#2949; NIH AIDS Reagent Program) or an anti-Gag antibody (Gag#6457; NIH AIDS Reagent Program) followed by detection using the SuperSignal

West Pico Chemiluminescent Substrate (Thermo Scientific, Rockford, IL).

Immunophenotyping

Surface expression of MHC-I and CD4 in HIV-1-infected and uninfected cells was done using flow cytometry. Study of MHC-I downregulation by Nef on infected thymocytes^{40,42} with FITC- and PE-conjugated monoclonal antibodies was done as previously described.^{42–45} PE-conjugated or FITC-conjugated KC57 mAb (Beckman Coulter, Brea, CA) was used to determine intracellular HIV-1_{Gag} protein expression in distinct thymocyte subsets^{43,46} and was performed following surface immunophenotyping.^{44,45} Study of CD4 downregulation by Nef on infected cells was described previously.⁴⁰ Briefly, 2×10^6 T1 cells were infected with 100 μ L (10 ng p24) of viral stock. Three days postinfection, surface expression of CD4 and mCD24 was detected with PE-conjugated human anti-CD4 (BioLegend, San Diego, CA) and FITC-conjugated mouse anti-CD24 (BioLegend). Multiparameter data acquisition was done on a dual-laser FACS Calibur flow cytometer (BDIS, San Jose, CA) or an MACSQuant VYB, and data analysis was performed with FCS Express 6.0.

In vitro infection of postnatal thymocytes

Normal human postnatal thymus specimens were obtained from infants and children undergoing corrective cardiac surgery. The tissue was cut into small pieces, red blood cells were removed by NH_4Cl /Tris lysis buffer, and the mechanically disrupted tissue was passed over a cell strainer to generate a single-cell suspension of thymocytes as previously described.^{39,44,46} Thymocytes were infected *in vitro* with CXCR4-tropic pediatric isolates (PI-2 and PI-2.1), the CXCR4-tropic molecular clone NL4-3, the CCR5-tropic molecular clone NF-NSX, or mock infected with the supernatant from the same CD4^+ T cell pool used for virus stock preparation. The thymocytes were cultured in serum-free medium for 14 days with IL-2 and IL-4 as previously described.³² All thymocyte infections with pediatric isolates were standardized by using equal numbers of virus infectious units/ 10^7 cells. HIV-1 virion production was measured by HIV-1 Gag p24 enzyme-linked immunosorbent assay as previously described.⁴⁴

In vivo infection of SCID-hu mice

C.B-17 SCID mice were bred at UCLA and implanted with human fetal thymus and liver graft (thy/liv) under the murine kidney capsule.⁴⁷ Four to six months postimplantation, the thy/liv grafts of SCID-hu mice implanted with thy/liv grafts from the same donor were infected with different HIV-1 isolates (PI-2, PI-2.1, NL4-3, or NFN-SX) by injecting 10–30 ng of p24 in a 50- μ L volume directly into the graft as previously described.⁴⁴ Mock-infected implants, used as controls in the experiments, were prepared by injecting implants with 50 μ L of appropriate control supernatants. Two different experiments consisting of 10–12 SCID-hu mice per 3-week time point were conducted. MHC-I and CD4 phenotyping in HIV-1-infected cells was done at 3 weeks postinfection as previously described.⁴⁵

Results

PI-2 is less cytopathic to thymocytes in vitro than PI-2.1 or NL4-3

HIV-1 strains isolated from *in utero* and perinatally infected children vary in rates of disease progression and exhibit differential effects on thymopoiesis during *in vitro* culture.^{6,32} Beyond the observation that CXCR4-tropic HIV isolates cause more thymocyte depletion than CCR5-tropic HIV isolates,⁴⁸ it is unclear what molecular mechanisms determine differential effects on thymocyte depletion and thymic development by HIV-1. Two CXCR4-tropic HIV-1 isolates (PI-2 and PI-2.1) were isolated from an *in utero*-infected infant and were found to infect CD4^+ thymocytes differently.³² Primary human thymocytes were infected *in vitro* with PI-2, PI-2.1, or NL4-3 viruses in parallel with a mock-infection control and cultured in serum-free medium supplemented with IL-2 and IL-4 for 14 days. Flow cytometric analysis of surface CD4 and CD8 expression was done on the thymocytes to measure subset depletion. Apoptotic and dead cells were excluded using 7-AAD, and quadrants were set using isotype controls. Following *in vitro* infection of human thymocytes, PI-2 was less efficient in depleting CD4^+ thymocytes (9% CD4^+ depletion) compared with PI-2.1 (40% CD4^+ depletion) or the CXCR4-tropic molecular clone NL4-3 (90% CD4^+ depletion) as related to the percentage of CD4^+ cells in mock-infected thymocytes (Fig. 1A). Despite initial infection with equal infectious units, PI-2 *in vitro* replication was also lower than PI-2.1 as measured by p24 levels in the culture supernatants (Fig. 1B).

PI-2 contains amino acid mutations in the Nef protein

The Nef protein was previously shown to affect the pathogenicity of HIV-1 in cultured thymocytes¹⁷ and the thy/liv implant in the SCID-hu mouse model.⁴⁹ Sequencing of the *nef* alleles found in PI-2 and PI-2.1 was done to examine a possible cause of their discrepancy in thymic cytopathicity. Analysis of the two *nef* sequences revealed that the amino acids (aa) of these two isolates were highly similar, and the majority of reported functional motifs were conserved in both isolates (Fig. 2A). The myristoylation residue (G2) required for membrane localization, the oligomerization domain needed for all functions, and motifs such as WL₅₈, LL₁₆₅, DD₁₇₅, and EE₁₅₅, which are required for CD4 binding and lysosomal degradation function, were found to be preserved in both isolates. However, a deletion of seven amino acids (Nef Δ K7S) from amino acid positions 82 to 88 (₈₂KGALDLS₈₈) was present in PI-2 but not in PI-2.1. This sequence falls within the core domain of Nef (aa 72–203), which is a relatively conserved and structured type II polyproline helix (aa 72–77) followed by two α -helices (aa 81–120).⁵⁰ This novel deletion is also 4 amino acids downstream of a polyproline helix (₇₂PXXPXR/K₇₇) binding site for Src-homology domain 3 (SH3) that mediates cellular activation and MHC-I downregulation.^{15,29} In addition to Nef Δ K7S, a single amino acid mutation at position 71 from K to R (K71R), adjacent to the SH3 binding site, was observed in PI-2.

Nef mutations in PI-2 do not inhibit expression of structural proteins

To assess if the identified mutations in the *nef* gene of PI-2 inhibited Nef protein expression, Western blot analysis was

FIG. 1. Thymocytes infected with the CXCR4-tropic HIV isolate PI-2 *in vitro* show less CD4 depletion and lower HIV-Gag p24 levels than thymocytes infected with PI-2.1 or NL4-3. Thymocytes were infected *in vitro* with CXCR4-tropic pediatric isolates (PI-2 and PI-2.1), the CXCR4-tropic NL4-3 molecular clone NL4-3, or mock infected with the supernatant from the same cells (CD4⁺ T cell pool). (A) Fourteen days postinfection, thymocytes were stained for CD4 and CD8. Apoptotic and dead cells were excluded using 7-AAD, and quadrants were set using isotype controls. (B) Viral replication was followed by measuring HIV Gag p24 antigen in the supernatants. PI, pediatric isolates.

done on cells infected with CXCR4-tropic molecular clone NL4-3 (lane 1), CCR5-tropic pediatric isolate PI-8³² (lane 2), and CXCR4-tropic isolate PI-2 (lane 3). The results show that PI-2-infected cells expressed similar levels of HIV-1 Gag (Fig. 2B) and Nef (Fig. 2C) as cells infected with PI-8 or NL4-3 and used as controls. Accordingly, PI-2-infected cells express HIV-1 Nef despite mutations.

PI-2 is impaired in MHC-I downregulation *in vitro* and *in vivo*

While a myriad of cellular effects have been shown for Nef, two of its most clearly defined functions are to downregulate cell surface expression of MHC-I and CD4 mole-

cules. These functions were assessed in thymocyte cultures infected with PI-2 and PI-2.1 and compared with cultures infected with the HIV-1 molecular clones NL4-3 or NFN-SX (NL4-3 with a substituted CCR5-tropic Env from strain JR-FL). Mock-infected cultures were prepared by adding supernatants of uninfected activated allogeneic CD4⁺ T cell pools. Thymocytes infected *in vitro* (Fig. 3) were analyzed for surface expression of MHC-I. Following infection, MHC-I downregulation was assessed by the mean fluorescence intensity (MFI) of MHC-I in HIV-1 Gag⁺ cells (KC57⁺). The cells were surface stained with antibodies to MHC-I (HLA-ABC-FITC) and CD3-APC followed with intracellular staining for HIV-1 Gag proteins using the KC57-PE antibody. Isotype controls were used to set the quadrants. MHC-I expression on mock-infected thymocytes is shown in Figure 3A. Cells productively infected (KC57⁺) with CXCR4-tropic NL4-3 containing a functional Nef (Fig. 3B) had decreased expression of MHC-I (MFI=263) compared with uninfected (KC57⁻) cells (MFI=745). Cells infected with the CCR5-tropic virus NF-NSX containing a functional Nef (Fig. 3C) had a decreased expression of MHC-I (MFI=217) compared with uninfected cells (MFI=551). Cells infected with the pediatric isolate PI-2.1 containing a functional Nef (Fig. 3D) also had decreased MHC-I expression (MFI=220) compared with uninfected cells (MFI=639). However, cells infected with the pediatric isolate PI-2 containing the Nef mutation (Fig. 3E) showed a smaller decrease in MHC-I surface expression (MFI=417) compared with uninfected cells (MFI=573).

The ability of the pediatric isolate Nef proteins to downregulate MHC-I was also tested *in vivo* using an SCID-hu mouse model implanted with human fetal thy/liv tissue that forms a thymus-like organoid as previously described.^{44,45} SCID-hu mice were infected with NL4-3, PI-2, PI-2.1, or mock-negative control supernatant. MHC-I downregulation in productively infected (KC57⁺) thymocytes was measured *ex vivo* at 3 weeks postinfection by measuring the HLA-ABC MFI. As shown in Figure 4A, cells infected with NL4-3 (*n*=5 mice) and PI-2.1 (*n*=5 mice) showed decreased mean surface expression of HLA-ABC compared with cells infected with PI-2 (*n*=6 mice). PI-2 expressing thymocytes had similar levels of MHC-I expression as mock infection (*n*=5 mice). Statistical analysis was done using two-tailed unpaired *t* tests. Similar to the *in vitro* experiments, infections of thy/liv implants with PI-2, PI-2.1, or NL4-3 again demonstrated diminished capabilities of PI-2 to downregulate MHC-I following infection, whereas PI-2.1 exhibited comparable MHC-I downregulation with NL4-3 (Fig. 4A). The percentages of CD4⁺ expression on thymocytes in thy/liv implants following infection with PI-2 were the same as in mock-infected implants, contrary to our earlier *in vitro* experiments. However, PI-2.1-infected implants showed a similar decrease in the percentage of CD4⁺ thymocytes as NL4-3-infected implants (Fig. 4B).

HIV-1 PI-2 Nef downregulates CD4 but not MHC-I expression in the absence of Vpu and Env *in vitro*

To investigate the small discrepancy in CD4 downregulation between our *in vitro* and *in vivo* experiments, we analyzed the Nef proteins separately from other HIV-1 proteins, Env and Vpu, which have previously been reported

126

FIG. 4. Pediatric HIV isolate 2 (PI-2) is impaired in MHC-I downregulation *in vivo*, but is less cytopathic than PI-2.1. SCID mice implanted with human fetal thy/liv (SCID-hu mice) were infected with NL4-3 or pediatric isolates PI-2.1 and PI-2. (A) MHC-I downregulation was measured *ex vivo* at 3 weeks postinfection. The MFI values show the levels of MHC-I on productively infected (KC57⁺) thymocytes. Cells infected with NL4-3 ($n=5$ mice), PI-2.1 ($n=5$ mice), and PI-2 ($n=6$ mice). Error bars depict standard error of the mean. Statistical analysis was done using two-tailed unpaired *t* tests. PI-2 expressing thymocytes had similar levels of MHC-I expression as mock infection ($n=5$ mice, data not shown). (B) Percentages of thymocytes expressing CD4 were measured *ex vivo* at 3 weeks postinfection. The percentages of CD4⁺ thymocytes in PI-2 infected are similar to the mock-infected thy/liv implants. thy/liv, thymus/liver.

to contribute to CD4 surface downregulation.^{51,52} To test the effect of Nef separately from Env and Vpu while maintaining their physiologic expression levels, PI-2 and PI-2.1 *nef* genes were inserted into an NL4-3-based genome containing deletions of *env* and *vpu*.⁴⁰ Single-cycle infection was done using VSV-G pseudotyped viruses expressing a CD24 tag and the various *nef* genes. In agree-

ment with our earlier *in vitro* data obtained in thymocytes infected with the original pediatric isolates, PI-2 Nef was unable to downregulate MHC-I in infected cells as much as Nef from PI-2.1 and NL4-3 (Fig. 5A–E). Confirming our previous experiment, PI-2 Nef was still able to downregulate CD4 surface expression *in vitro*, even in the absence of Vpu and Env (Fig. 5F–J).

FIG. 5. HIV PI-2 Nef downregulates CD4 but not MHC-I expression in the absence of Vpu, Env, and Vpr. To assess the MHC-class I downregulation of the *nef* genes in pediatric isolates PI-2.1 and PI-2 in the absence of Vpu, Env, and Vpr proteins, the *nef* genes were cloned out of the PI-2.1, PI-2, and NL4-3 isolates and placed in an NL4-3 vector lacking Vpu, Env, and Vpr. To determine the effect of PI-2.1 and PI-2 Nef proteins on MHC-I and CD4 downregulation, VSV-G pseudotyped virus, including a CD24 tag and the various *nef* genes, was used to infect T1 cells. HLA-A2 and CD4 surface expression in T1 cells infected with (A, F) mock control; (B, G) VSV-G Nef pseudotyped defective NL4-3 lacking Vpu, Env, and Vpr; (C, H) NL4-3 ΔNef; (D, I) VSV-G pseudotyped PI-2.1 *nef* lacking *vpu*, *env*, and *vpr*; or (E, J) VSV-G pseudotyped PI-2 *nef* lacking *vpu*, *env*, and *vpr*.

Discussion

Following comparison of two primary HIV-1 pediatric isolates from the same *in utero*-infected infant, PI-2 and PI-2.1, we identified a unique in-frame deletion Nef Δ K7S in PI-2, which attenuates the depletion of thymocytes *in vitro*. Nef proteins bearing this mutation are defective for MHC-I downregulation, but this effect is not due to reduced Nef expression or global Nef dysfunction. Nef proteins bearing this mutation were still able to downregulate CD4 surface expression *in vitro*. This novel Δ K7S in-frame deletion is in a location that is not known to contain any specific motifs required for previously defined Nef functions. However, it is adjacent to the SH3 binding site that is critical for multiple Nef functions, including cellular activation and MHC-I downregulation.^{53,54} It is possible that Δ K7S alters the tertiary structure of Nef and thus interferes with interactions with host factors. While there is a K versus R polymorphism that falls within the PXXP motif between residues 72–75 when comparing PI-2.1 with PI-2, this is unlikely to contribute to impaired MHC-I downregulation by PI-2. Although the NL4-3 Nef sequence has T71 (Fig. 2A), the subtype B consensus sequence is actually 71R, and thus, PI-2 actually matches “wild-type” Nef more closely than PI-2.1 or NL4-3 at position 71. Moreover, K71R is a conservative substitution. Although highly unlikely, however, we cannot definitively rule out that 71R somehow contributed to the impaired MHC-I downregulation by PI-2.

Previous data show that various mutations in Nef affect oligomerization as well as ternary complex formation with cell molecules such as AP-1 and MHC-I,^{50,55–57} and a survey of *in vivo* Nef sequences demonstrated frequent examples of Nef with impaired MHC-I downregulation despite no identifiable mutations in the defined motifs involved in this function.⁵⁸ Similarly, the significance of the K71R substitution mutation in PI-2 Nef is unclear. This mutation is also located in proximity to the canonical sequence required for SH3 binding. Mutational studies have demonstrated that residues flanking the motif play critical role in determining binding specificity to SH3 domain.⁵⁹

Others have reported that Nef-mediated CD4 downregulation correlates with increased viral pathogenesis in thymocytes, and MHC-I downregulation has diminished influence in cytopathicity.⁶⁰ In contrast, our findings indicate that PI-2-mutated *nef* containing virus was less cytotoxic in thymocytes than PI-2.1, despite having a similar ability to downregulate surface expression of CD4 *in vitro*. PI-2-infected cells did not downregulate MHC-I but did decrease CD4 surface expression *in vitro*, indicating that the novel identified mutations in PI-2 Nef may not have affected binding to several host factors reported to be involved in CD4 downregulation.^{61,62} Point mutations in the *nef* gene (WL57AA, LL164AA, and DD174AA) of the NL4-3 molecular clone abrogate CD4 downregulation and decrease Nef-mediated pathogenesis.⁶⁰ These mutations were not present in PI-2. Van Nuffel *et al.* have previously reported that Nef:PAK2 and an intact VGF motif within Nef are required for reduced thymic output.⁶³ Despite having intact VGF and PAK2 binding motifs and representative CD4 downregulation *in vitro*, PI-2 still lacked the virulence and thymic depletion seen in thymocytes infected with PI-2.1 or molecular clones, indicating a correlation between MHC-I downregulation and Nef's ability to deplete thymocytes.

Given the importance of MHC-I molecules in T cell development, it is unclear whether the defect in MHC-I downregulation by HIV isolate PI-2 is directly related to the attenuated thymic cytopathicity, or if the novel identified mutations disrupt another pathway that potentiates the negative effect of Nef in the thymopoiesis. Further unraveling this phenomenon may provide greater insight into the particularly rapid disease progression seen in HIV-1-infected children, and may offer therapeutic targets for this particularly vulnerable population.

Sequence Data

The nucleotide sequences for PI-2 and PI-2.1 *nef* genes have been deposited in GenBank under accession numbers KC702883 and KC702882, respectively.

Acknowledgments

The following reagents were obtained from the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: anti-Nef antibody (Nef#2949), anti-Gag antibody (Gag#6457), and the HIV-1 NL4-3 infectious molecular clone (pNL4-3) from Dr. Malcolm Martin (cat. no. 114).³⁸ The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Authors' Contributions

A.A., L.P.-M., R.L.F., O.O.Y., and C.H.U. conceived and designed the experiments; A.A., L.P.-M., A.D.C., and D.A.-P. performed the experiments; A.A., L.P.-M., R.L.F., A.D.C., D.A.-P., and C.H.U. analyzed the data; Y.B. and O.O.Y. contributed reagents/materials/analysis tools and advice; A.A., R.L.F., O.O.Y., and C.H.U. wrote the article.

Author Disclosure Statement

The authors do not have any financial or nonfinancial competing interests.

Funding Information

This work was supported by the National Institutes of Health, grant numbers: HD29341, AI054286, AI32440, and CA016042. This work was also supported by the UCLA AIDS Institute, UCLA Center for AIDS Research (CFAR) NIH AI28697, James B. Pendleton Charitable Trust, and the McCarthy Family Foundation.

References

1. Douek DC, Betts MR, Hill BJ, *et al.*: Evidence for increased T cell turnover and decreased thymic output in HIV infection. *J Immunol* 2001;167:6663–6668.
2. Bonyhadi ML, Rabin L, Salimi S, *et al.*: HIV induces thymus depletion *in vivo*. *Nature* 1993;363:728–732.
3. Hanna Z, Kay DG, Rebai N, Guimond A, Jothy S, Jolicoeur P: Nef harbors a major determinant of pathogenicity for an AIDS-like disease induced by HIV-1 in transgenic mice. *Cell* 1998;95:163–175.
4. Bandera A, Ferrario G, Saresella M, *et al.*: CD4⁺ T cell depletion, immune activation and increased production of regulatory T cells in the thymus of HIV-infected individuals. *PLoS One* 2010;5:e10788.

5. Su L, Kaneshima H, Bonyhadi M, *et al.*: HIV-1-induced thymocyte depletion is associated with indirect cytopathogenicity and infection of progenitor cells in vivo. *Immunity* 1995;2:25–36.
6. Kourtis AP, Ibegbu C, Nahmias AJ, *et al.*: Early progression of disease in HIV-infected infants with thymus dysfunction. *N Engl J Med* 1996;335:1431–1436.
7. Pennington DJ, Jenkins SA, Brady HJ, Miles CG, Dzierzak EA, Abraham DJ: HIV-1 Nef severely impairs thymocyte development and peripheral T-cell function by a CD4-independent mechanism. *Genes Funct* 1997;1:321–335.
8. Kestler HW, 3rd, Ringler DJ, Mori K, *et al.*: Importance of the *nef* gene for maintenance of high virus loads and for development of AIDS. *Cell* 1991;65:651–662.
9. Skowronski J, Parks D, Mariani R: Altered T cell activation and development in transgenic mice expressing the HIV-1 *nef* gene. *EMBO J* 1993;12:703–713.
10. Stove V, Naessens E, Stove C, Swigut T, Plum J, Verhasselt B: Signaling but not trafficking function of HIV-1 protein Nef is essential for Nef-induced defects in human intrathymic T-cell development. *Blood* 2003;102:2925–2932.
11. Jubier-Maurin V, Saragosti S, Perret JL, *et al.*: Genetic characterization of the *nef* gene from human immunodeficiency virus type 1 group M strains representing genetic subtypes A, B, C, E, F, G, and H. *AIDS Res Hum Retroviruses* 1999;15:23–32.
12. Arien KK, Verhasselt B: HIV Nef: Role in pathogenesis and viral fitness. *Curr HIV Res* 2008;6:200–208.
13. Daniel MD, Kirchhoff F, Czajak SC, Sehgal PK, Desrosiers RC: Protective effects of a live attenuated SIV vaccine with a deletion in the *nef* gene. *Science* 1992;258:1938–1941.
14. Kirchhoff F, Greenough TC, Brettler DB, Sullivan JL, Desrosiers RC: Brief report: Absence of intact *nef* sequences in a long-term survivor with nonprogressive HIV-1 infection. *N Engl J Med* 1999;332:228–232.
15. Deacon NJ, Tsykin A, Solomon A, *et al.*: Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and recipients. *Science* 1995;270:988–991.
16. Learmont JC, Geczy AF, Mills J, *et al.*: Immunologic and virologic status after 14 to 18 years of infection with an attenuated strain of HIV-1. A report from the Sydney Blood Bank Cohort. *N Engl J Med* 1999;340:1715–1722.
17. Verhasselt B, Naessens E, Verhofstede C, *et al.*: Human immunodeficiency virus *nef* gene expression affects generation and function of human T cells, but not dendritic cells. *Blood* 1999;94:2809–2818.
18. Harris M: HIV: A new role for Nef in the spread of HIV. *Curr Biol* 1999;9:R459–R461.
19. Zuo J, Suen J, Wong A, *et al.*: Functional analysis of HIV type 1 *Nef* gene variants from adolescent and adult survivors of perinatal infection. *AIDS Res Hum Retroviruses* 2012;28:486–492.
20. Landi A, Iannucci V, Nuffel AV, Meuwissen P, Verhasselt B: One protein to rule them all: Modulation of cell surface receptors and molecules by HIV Nef. *Curr HIV Res* 2011;9:496–504.
21. Geyer M, Fackler OT, Peterlin BM: Structure—Function relationships in HIV-1 Nef. *EMBO Rep* 2001;2:580–585.
22. Garcia JV, Miller AD: Serine phosphorylation-independent downregulation of cell-surface CD4 by *nef*. *Nature* 1991;350:508–511.
23. Mann JK, Byakwaga H, Kuang XT, *et al.*: Ability of HIV-1 Nef to downregulate CD4 and HLA class I differs among viral subtypes. *Retrovirology* 2013;10:100.
24. Mann JK, Chopera D, Omarjee S, *et al.*: Nef-mediated down-regulation of CD4 and HLA class I in HIV-1 subtype C infection: Association with disease progression and influence of immune pressure. *Virology* 2014;468–470:214–225.
25. Schwartz O, Marechal V, Le Gall S, Lemonnier F, Heard JM: Endocytosis of major histocompatibility complex class I molecules is induced by the HIV-1 Nef protein. *Nat Med* 1996;2:338–342.
26. Stumptner-Cuvelette P, Morchoisne S, Dugast M, *et al.*: HIV-1 Nef impairs MHC class II antigen presentation and surface expression. *Proc Natl Acad Sci U S A* 2001;98:12144–12149.
27. Swigut T, Shohdy N, Skowronski J: Mechanism for down-regulation of CD28 by Nef. *EMBO J* 2001;20:1593–1604.
28. Rosa A, Chande A, Ziglio S, *et al.*: HIV-1 Nef promotes infection by excluding SERINC5 from virion incorporation. *Nature* 2015;526:212–217.
29. Collins KL, Chen BK, Kalams SA, Walker BD, Baltimore D: HIV-1 Nef protein protects infected primary cells against killing by cytotoxic T lymphocytes. *Nature* 1998;391:397–401.
30. Ali A, Ng HL, Dagarag MD, Yang OO: Evasion of cytotoxic T lymphocytes is a functional constraint maintaining HIV-1 Nef expression. *Eur J Immunol* 2005;35:3221–3228.
31. Mwimanzu F, Toyoda M, Mahiti M, *et al.*: Resistance of major histocompatibility complex class B (MHC-B) to Nef-mediated downregulation relative to that of MHC-A is conserved among primate lentiviruses and influences antiviral T cell responses in HIV-1-infected individuals. *J Virol* 2018;92:1–19.
32. Pedroza-Martins L, Boscardin WJ, Anisman-Posner DJ, Schols D, Bryson YJ, Uittenbogaart CH: Impact of cytokines on replication in the thymus of primary human immunodeficiency virus type 1 isolates from infants. *J Virol* 2002;76:6929–6943.
33. Ganeshan S, Dickover RE, Korber BT, Bryson YJ, Wolinsky SM: Human immunodeficiency virus type 1 genetic evolution in children with different rates of development of disease. *J Virol* 1997;71:663–677.
34. von Seidlein L, Plaeger S, Dickover R, Krogstad P, Stiehm ER, Bryson Y: Primary human immunodeficiency virus type 1 infection during pregnancy associated with transmission of SI/MT-2 cell tropic virus and precipitous loss of CD4 cells in mother and infant. *Pediatr Infect Dis J* 1998;17:528–530.
35. Koot M, Vos AH, Keet RP, *et al.*: HIV-1 biological phenotype in long-term infected individuals evaluated with an MT-2 cocultivation assay. *AIDS* 1992;6:49–54.
36. Ferbas J, Kaplan AH, Hausner MA, *et al.*: Virus burden in long-term survivors of human immunodeficiency virus (HIV) infection is a determinant of anti-HIV CD8⁺ lymphocyte activity. *J Infect Dis* 1995;172:329–339.
37. Dickover RE, Dillon M, Leung KM, *et al.*: Early prognostic indicators in primary perinatal human immunodeficiency virus type 1 infection: Importance of viral RNA and the timing of transmission on long-term outcome. *J Infect Dis* 1998;178:375–387.
38. Adachi A, Gendelman HE, Koenig S, *et al.*: Production of acquired immunodeficiency syndrome-associated retrovirus

- in human and nonhuman cells transfected with an infectious molecular clone. *J Virol* 1986;59:284–291.
39. Pedroza-Martins L, Gurney KB, Torbett BE, Uittenbogaart CH: Differential tropism and replication kinetics of human immunodeficiency virus type 1 isolates in thymocytes: Coreceptor expression allows viral entry, but productive infection of distinct subsets is determined at the postentry level. *J Virol* 1998;72:9441–9452.
 40. Ali A, Realegeno S, Yang OO, Lewis MJ: Simultaneous assessment of CD4 and MHC-I downregulation by Nef primary isolates in the context of infection. *J Virol Methods* 2009;161:297–304.
 41. Ali A, Jamieson BD, Yang OO: Half-genome human immunodeficiency virus type 1 constructs for rapid production of reporter viruses. *J Virol Methods* 2003;110:137–142.
 42. Schmid I, Uittenbogaart CH, Keld B, Giorgi JV: A rapid method for measuring apoptosis and dual-color immunofluorescence by single laser flow cytometry. *J Immunol Methods* 1994;170:145–157.
 43. Schmid I, Uittenbogaart CH, Giorgi JV: A gentle fixation and permeabilization method for combined cell surface and intracellular staining with improved precision in DNA quantification. *Cytometry* 1991;12:279–285.
 44. Gurney KB, Colantonio AD, Blom B, Spits H, Uittenbogaart CH: Endogenous IFN- α production by plasmacytoid dendritic cells exerts an antiviral effect on thymic HIV-1 infection. *J Immunol* 2004;173:7269–7276.
 45. Gurney KB, Uittenbogaart CH: Human immunodeficiency virus persistence and production in T-cell development. *Clin Vaccine Immunol* 2006;13:1237–1245.
 46. Uittenbogaart CH, Anisman DJ, Jamieson BD, *et al.*: Differential tropism of HIV-1 isolates for distinct thymocyte subsets in vitro. *AIDS* 1996;10:F9–F16.
 47. Aldrovandi GM, Feuer G, Gao L, *et al.*: The SCID-hu mouse as a model for HIV-1 infection. *Nature* 1993;363:732–736.
 48. Correa R, Munoz-Fernandez MA: Viral phenotype affects the thymic production of new T cells in HIV-1-infected children. *AIDS* 2001;15:1959–1963.
 49. Jamieson BD, Aldrovandi GM, Planelles V, *et al.*: Requirement of human immunodeficiency virus type 1 nef for in vivo replication and pathogenicity. *J Virol* 1994;68:3478–3485.
 50. Arold S, Hoh F, Domergue S, *et al.*: Characterization and molecular basis of the oligomeric structure of HIV-1 nef protein. *Protein Sci* 2000;9:1137–1148.
 51. Kerkau T, Bacik I, Bennink JR, *et al.*: The human immunodeficiency virus type 1 (HIV-1) Vpu protein interferes with an early step in the biosynthesis of major histocompatibility complex (MHC) class I molecules. *J Exp Med* 1997;185:1295–1305.
 52. Yewdell JW, Hill AB: Viral interference with antigen presentation. *Nat Immunol* 2002;3:1019–1025.
 53. Madrid R, Janvier K, Hitchin D, *et al.*: Nef-induced alteration of the early/recycling endosomal compartment correlates with enhancement of HIV-1 infectivity. *J Biol Chem* 2005;280:5032–5044.
 54. Coleman SH, Madrid R, Van Damme N, *et al.*: Modulation of cellular protein trafficking by human immunodeficiency virus type 1 Nef: Role of the acidic residue in the ExxxLL motif. *J Virol* 2006;80:1837–1849.
 55. Noviello CM, Benichou S, Guatelli JC: Cooperative binding of the class I major histocompatibility complex cytoplasmic domain and human immunodeficiency virus type 1 Nef to the endosomal AP-1 complex via its mu subunit. *J Virol* 2008;82:1249–1258.
 56. Wonderlich ER, Williams M, Collins KL: The tyrosine binding pocket in the adaptor protein 1 (AP-1) μ 1 subunit is necessary for Nef to recruit AP-1 to the major histocompatibility complex class I cytoplasmic tail. *J Biol Chem* 2008;283:3011–3022.
 57. Singh RK, Lau D, Noviello CM, Ghosh P, Guatelli JC: An MHC-I cytoplasmic domain/HIV-1 Nef fusion protein binds directly to the μ subunit of the AP-1 endosomal coat complex. *PLoS One* 2009;4:e8364.
 58. Lewis MJ, Balamurugan A, Ohno A, Kilpatrick S, Ng HL, Yang OO: Functional adaptation of Nef to the immune milieu of HIV-1 infection in vivo. *J Immunol* 2008;180:4075–4081.
 59. Kuo LS, Baugh LL, Denial SJ, *et al.*: Overlapping effector interfaces define the multiple functions of the HIV-1 Nef polyproline helix. *Retrovirology* 2012;9:47.
 60. Stoddart CA, Geleziunas R, Ferrell S, *et al.*: Human immunodeficiency virus type 1 Nef-mediated downregulation of CD4 correlates with Nef enhancement of viral pathogenesis. *J Virol* 2003;77:2124–2133.
 61. Landi A, Timermans CG, Naessens E, Vanderstraeten H, Stove V, Verhasselt B: The human immunodeficiency virus (HIV) Rev-binding protein (HRB) is a co-factor for HIV-1 Nef-mediated CD4 downregulation. *J Gen Virol* 2016;97:778–785.
 62. Landi A, Vermeire J, Iannucci V, *et al.*: Genome-wide shRNA screening identifies host factors involved in early endocytic events for HIV-1-induced CD4 down-regulation. *Retrovirology* 2014;11:118.
 63. Van Nuffel A, Arien KK, Stove V, *et al.*: Primate lentiviral Nef proteins deregulate T-cell development by multiple mechanisms. *Retrovirology* 2013;10:137.

Address correspondence to:

Christel H. Uittenbogaart
 Department of Microbiology, Immunology
 & Molecular Genetics
 University of California, Los Angeles
 615 Charles E. Young Drive South
 Los Angeles, CA 90095

E-mail: uittenbo@ucla.edu