FORUM ORIGINAL RESEARCH COMMUNICATION

Decreased Soluble Guanylate Cyclase Contributes to Cardiac Dysfunction Induced by Chronic Doxorubicin Treatment in Mice

Sara Vandenwijngaert¹, Melissa Swinnen², Ann-Sophie Walravens², Manu Beerens², Hilde Gillijns², Ellen Caluwé², Robert E. Tainsh¹, Daniel I. Nathan¹, Kaitlin Allen¹, Peter Brouckaert³, Jozef Bartunek^{2,4}, Marielle Scherrer-Crosbie⁵, Kenneth D. Bloch^{1,5}, Donald B. Bloch^{1,6}, Stefan P. Janssens^{2,*} and Emmanuel S. Buys^{1,*}

Abstract

Aims: The use of doxorubicin, a potent chemotherapeutic agent, is limited by cardiotoxicity. We tested the hypothesis that decreased soluble guanylate cyclase (sGC) enzyme activity contributes to the development of doxorubicin-induced cardiotoxicity.

Results: Doxorubicin administration (20 mg/kg, intraperitoneally [IP]) reduced cardiac sGC activity in wild-type (WT) mice. To investigate whether decreased sGC activity contributes to doxorubicin-induced cardiotoxicity, we studied mice with cardiomyocyte-specific deficiency of the sGC $\alpha 1$ -subunit (mice with cardiomyocyte-specific deletion of exon 6 of the sGC $\alpha 1$ allele [sGC $\alpha 1^{-/-CM}$]). After 12 weeks of doxorubicin administration (2 mg/kg/week IP), left ventricular (LV) systolic dysfunction was greater in sGC $\alpha 1^{-/-CM}$ than WT mice. To further assess whether reduced sGC activity plays a pathogenic role in doxorubicin-induced cardiotoxicity, we studied a mouse model in which decreased cardiac sGC activity was induced by cardiomyocyte-specific expression of a dominant negative sGC $\alpha 1$ mutant (DNsGC $\alpha 1$) upon doxycycline removal (Tet-off). After 8 weeks of doxorubicin administration, DNsGC $\alpha 1^{tg/+}$, but not WT, mice displayed LV systolic dysfunction and dilatation. The difference in cardiac function and remodeling between DNsGC $\alpha 1^{tg/+}$ and WT mice was even more pronounced after 12 weeks of treatment. Further impairment of cardiac function was attenuated when DNsGC $\alpha 1$ gene expression was inhibited (beginning at 8 weeks of doxorubicin treatment) by administering doxycycline. Furthermore, doxorubicin-associated reactive oxygen species generation was higher in sGC $\alpha 1$ -deficient than WT hearts.

Innovation and Conclusion: These data demonstrate that a reduction in cardiac sGC activity worsens doxorubicin-induced cardiotoxicity in mice and identify sGC as a potential therapeutic target. Various pharmacological sGC agonists are in clinical development or use and may represent a promising approach to limit doxorubicin-associated cardiotoxicity. *Antioxid. Redox Signal.* 26, 153–164.

Keywords: cardiovascular, cardiomyopathy, chemotherapy, doxorubicin, cyclic nucleotides, soluble guanylate cyclase

¹Department of Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital Research Institute and Harvard Medical School, Anesthesia Center for Critical Care Research, Boston, Massachusetts.

²Department of Cardiovascular Sciences, KU Leuven, Leuven, Belgium.

³Department of Biomedical Molecular Biology, Ghent University and Flanders Institute for Biotechnology, Ghent, Belgium.

⁴Cardiovascular Center, OLV Hospital, Aalst, Belgium.

⁵Cardiovascular Research Center, Massachusetts General Hospital Research Institute and Harvard Medical School, Boston, Massachusetts.

⁶Department of Medicine, Massachusetts General Hospital Research Institute and Harvard Medical School, Boston, Massachusetts.

*These two authors contributed equally.

Innovation

Adverse effects observed in chemotherapy patients receiving dexrazoxane, the only drug approved to attenuate doxorubicin-induced cardiotoxicity in patients, have restricted its use. This study illustrates that doxorubicin reduces cardiac soluble guanylate cyclase (sGC) activity and that reduced sGC activity exacerbates doxorubicin-induced cardiotoxicity. Increasing interest in sGC as a therapeutic target is driving clinical development of pharmacological sGC agonists, and the first drug in this class was recently approved to treat pulmonary hypertension. Because our data suggest that sGC protects against doxorubicin-induced cardiotoxicity, pharmacological sGC agonists could represent a novel therapeutic strategy to treat cardiotoxicity in patients undergoing chemotherapy.

Introduction

SINCE ITS DISCOVERY more than five decades ago, the anthracycline, doxorubicin, has become a central component of therapeutic regimens to treat various malignancies, including lymphoma, sarcoma, some leukemias, and breast cancer. Unfortunately, the clinical use of this potent chemotherapeutic agent is hampered by cardiotoxicity. The early adverse cardiac effects of doxorubicin are often transient and resolve spontaneously (18, 33, 37, 43). Of far greater concern is the development of chronic cardiomyopathy, leading to symptomatic congestive heart failure in up to 5% of doxorubicin-treated patients (8, 23, 46). In childhood cancer survivors, the long-term risk of late-onset doxorubicin-associated heart failure can be as high as 12% at the age of 40 (9). The most important risk factor for the development of doxorubicin-induced cardiomyopathy is the cumulative dose.

Although various pathogenic mechanisms have been suggested to underlie doxorubicin-induced cardiotoxicity, many studies support a pivotal role of oxidative stress (11, 22, 30, 32, 50, 56). Cardiomyocytes are particularly prone to damage by free radicals because of the high oxidative metabolism, low concentrations of antioxidant enzymes, and abundance of mitochondria (14, 15).

With a univalent redox potential of approximately -320 mV, doxorubicin is a good substrate for intracellular oxidoreductases such as xanthine oxidase, NADPH cytochrome P450 reductase, and NADH dehydrogenase (12–14, 49). Reduction of the quinone moiety of doxorubicin results in an unstable semiquinone radical that donates an electron to molecular oxygen, thereby forming superoxide radicals. If free iron is present, the semiquinone radical can also form a complex with iron, resulting in an anthracycline–iron free radical complex that in turn reduces molecular oxygen (31).

Dexrazoxane (ICRF-187), the only drug currently in clinical use to protect against doxorubicin-induced cardiotoxicity, is thought to confer cardioprotection by chelating iron and subsequently reducing anthracycline–iron free radical complex formation and superoxide radicals (21). Because of reports of lower tumor responsiveness or enhanced myelosuppression in chemotherapy-receiving patients treated with dexrazoxane, the US Food and Drug Administration and the European Medicines Agency restricted the use of dexrazoxane (38, 47). Although a growing body of literature suggests

that dexrazoxane does not adversely affect therapeutic response (28, 40, 41, 53), additional mechanism-based therapies for doxorubicin-induced cardiotoxicity are needed.

The second messenger cyclic guanosine 3',5'-monophosphate (cGMP) regulates a vast array of physiological processes vital to endothelial, vascular smooth muscle, and cardiomyocyte function. cGMP is generated by particulate guanylate cyclase in response to natriuretic peptides and by soluble guanylate cyclase (sGC) upon activation by nitric oxide (NO) and, less potently, carbon monoxide (CO). Disruption of the cGMP signaling pathway has been implicated in adverse cardiac remodeling and heart failure. Interestingly, doxorubicin administration reduced cardiac cGMP levels, which could be prevented by inhalation of CO (45).

In this study, we investigated the role of sGC in doxorubicin-induced cardiotoxicity. We observed that doxorubicin administration decreased cardiac sGC activity and that reduced sGC activity exacerbated reactive oxygen species (ROS) formation in the heart and development of doxorubicin-induced cardiac dysfunction.

Results

Doxorubicin administration reduces cardiac sGC activity

To examine the effect of doxorubicin treatment on sGC enzymatic activity in the heart, the increase in cGMP production induced by the NO donor diethylenetriamine NONOate (DETA-NO) was measured in cardiac extracts from wild-type (WT) mice 24 h after doxorubicin (20 mg/kg, intraperitoneally [IP]) or saline administration. Doxorubicin treatment reduced cardiac NO-stimulated sGC activity by $\sim 20\%$ (Fig. 1A).

sGC is a heterodimeric enzyme comprising a $\beta 1$ subunit and either an $\alpha 1$ or $\alpha 2$ subunit, with the sGC $\alpha 1\beta 1$ isoform being the predominant isoform in the heart (6). Immunoblot analysis did not reveal changes in sGC $\alpha 1$ or sGC $\beta 1$ protein expression in hearts of doxorubicin-treated mice (Supplementary Fig. S1; Supplementary Data are available online at www.liebertpub.com/ars), indicating that the doxorubicin-induced decrease in cardiac sGC activity is not due to altered sGC expression.

It is conceivable that the oxidative stress associated with doxorubicin administration results in direct oxidative modification of sGC, potentially leading to loss of its prosthetic heme moiety (2, 4, 26, 42, 48) and decreased enzyme activity. To explore this possibility, we administered the radical scavenger tempol to WT mice before treatment with doxorubicin (20 mg/kg, IP) or saline and evaluated cardiac sGC activity in the presence of DETA-NO. Tempol administration prevented the doxorubicin-induced decrease in cardiac NO-stimulated sGC activity (Fig. 1B), suggesting that this decrease in activity is due to oxidative modification of sGC.

Our hypothesis that doxorubicin treatment results in oxidation of sGC was further corroborated by the observation that BAY 58-2667 (cinaciguat), an sGC agonist specifically targeting oxidized and heme-free sGC (42), increased sGC activity in cardiac extracts obtained from doxorubicin- and saline-treated WT mice to the same extent (Fig. 1C). Thus, whereas in hearts of doxorubicin-treated mice, activation of sGC by NO—requiring the heme group of sGC to be in a reduced state—was impaired, sGC activation by BAY 58-2667 was not.

FIG. 1. Cardiac NO-stimulated sGC activity is decreased in mice treated with DOX. (A) NO-stimulated sGC activity was lower in cardiac extracts from mice treated with DOX (20 mg/kg IP, 24 h) than from mice administered saline ($n=10$ each). (B) Tempol administration before DOX or saline treatment prevented this decrease in cardiac NO-stimulated sGC activity ($n=9$ and 8 , respectively). (C) In addition, incubation with the sGC activator, BAY 58-2667, increased sGC activity in the cardiac extracts obtained from DOX- and saline-treated mice to the same extent ($n=10$ each). Data are expressed relative to sGC activity in the contemporaneous saline-treated control group. $*p<0.05$ versus saline. DOX, doxorubicin; IP, intraperitoneal; NO, nitric oxide; sGC, soluble guanylate cyclase.

Constitutive myocardial sGC α 1 deficiency exacerbates doxorubicin-induced cardiac dysfunction

To evaluate whether the doxorubicin-induced decrease in sGC activity contributes to the development of cardiotoxicity, we studied mice with cardiomyocyte-specific reduction of sGC activity (sGC α 1 $^{-/-CM}$). Cre-mediated deletion of exon 6 of sGC α 1 in cardiomyocytes attenuated the ability of DETA-NO to activate sGC in the heart: the relative fold increase in cGMP synthesis was 0.64 ± 0.09 versus 1.00 ± 0.18 in sGC α 1 $^{-/-CM}$ and WT mice, respectively ($n=10$ and 8 , $p<0.05$). Restriction of sGC α 1 deficiency to the heart in sGC α 1 $^{-/-CM}$ mice was confirmed by polymerase chain reaction (PCR) analysis (Supplementary Fig. S2). Baseline echocardiographic parameters, including left ventricular (LV) end-systolic and end-diastolic internal diameters and fractional shortening (FS), were similar in sGC α 1 $^{-/-CM}$ mice and WT littermates (Supplementary Table S1).

Cardiotoxicity was induced in sGC α 1 $^{-/-CM}$ and WT mice ($n=14$ for both) by administering doxorubicin (2 mg/kg IP, once weekly) for 12 weeks, as previously reported (51). Following 8 weeks of doxorubicin administration, echocardiography revealed that end-systolic internal diameter (LVID_{ES}) was increased (Fig. 2A) and FS decreased (Fig. 2B) in both sGC α 1 $^{-/-CM}$ and WT mice compared with baseline measurements obtained before the first doxorubicin injection.

However, after 12 weeks, doxorubicin-induced systolic dysfunction was significantly greater in sGC α 1 $^{-/-CM}$ mice than in WT littermates: LVID_{ES} was greater (Fig. 2A) and FS lower (Fig. 2B) in sGC α 1 $^{-/-CM}$ than in WT mice. In addition, compared with baseline, LV end-diastolic internal diameter (LVID_{ED}) was modestly increased after 12 weeks of doxorubicin treatment in sGC α 1 $^{-/-CM}$ mice (3.3 ± 0.04 vs. 3.1 ± 0.03 mm, respectively, $p<0.05$), but not in WT mice (3.2 ± 0.05 vs.

3.1 ± 0.03 mm, $p>0.05$). Invasive LV pressure-volume measurements confirmed more pronounced systolic dysfunction, as evidenced by the reduced ejection fraction (EF), in doxorubicin-treated sGC α 1 $^{-/-CM}$ versus WT mice (Table 1).

Similar to previous reports (10, 35, 36), doxorubicin administration in mice was associated with a decrease in heart rate (HR), assessed *via* invasive hemodynamic measurements (Table 1) and echocardiography (sGC α 1 $^{-/-CM}$ + doxorubicin vs. sGC α 1 $^{-/-CM}$ + saline; 426 ± 18 vs. 505 ± 18 beats per minute [bpm] and WT + doxorubicin vs. WT + saline; 428 ± 17 vs. 531 ± 13 bpm, $p<0.05$ for doxorubicin vs. saline in both genotypes). HR did not differ between doxorubicin-treated sGC α 1 $^{-/-CM}$ and WT mice ($p>0.05$).

Furthermore, doxorubicin administration reduced survival rates to a similar extent in sGC α 1 $^{-/-CM}$ and WT mice (71% vs. 86%, respectively, $n=14$ for both, $p>0.05$), as well as body weights ($-23 \pm 3\%$ vs. $-19 \pm 2\%$, respectively, $n=10$ and 12 , $p>0.05$).

Doxorubicin-associated cardiotoxicity is increased in mice with inducible cardiomyocyte-specific expression of a dominant negative sGC α 1 mutant

To further assess whether reduced sGC activity plays a pathogenic role in doxorubicin-induced cardiotoxicity, a second mouse model with inducible cardiomyocyte-specific sGC α 1 deficiency was generated and studied. Upon withdrawal of doxycycline from the diet, a dominant negative sGC α 1 mutant (DNsGC α 1, Tet-Off system) is expressed that competes with sGC α 1 and sGC α 2 for binding to sGC β 1, thereby inhibiting the formation of the two catalytically active sGC heterodimers in the heart.

Four weeks after doxycycline removal from the diet, the ability of DETA-NO to activate sGC was attenuated in the

FIG. 2. Greater systolic dysfunction in sGCα1^{-/-CM} than WT mice after 12 weeks of DOX treatment. Echocardiographic analysis revealed decreased LV systolic function in sGCα1^{-/-CM} and WT mice ($n=14$ each) after 8 weeks of DOX administration, illustrated by an increased LVID_{ES} (A) and a reduced FS (B). After 12 weeks of DOX treatment, systolic dysfunction was significantly greater in sGCα1^{-/-CM} than in WT mice. [†] $p < 0.05$ versus baseline and ^{*} $p < 0.05$ versus WT (same time point). FS, fractional shortening; LV, left ventricular; LVID_{ES}, left ventricular end-systolic internal diameter; sGCα1^{-/-CM}, mice with cardiomyocyte-specific deletion of exon 6 of the sGCα1 allele; WT, wild-type.

hearts of mice with cardiomyocyte-specific expression of a dominant negative mutation of sGCα1 (DNsGCα1^{tg/+}) mice: the relative fold increase in cGMP synthesis was 0.59 ± 0.07 versus 1.00 ± 0.11 in DNsGCα1^{tg/+} and WT mice, respectively ($n=8$ each, $p < 0.05$). Baseline echocardiographic parameters were similar in DNsGCα1^{tg/+} mice and WT littermates (Supplementary Table S2).

After 8 weeks of doxorubicin treatment (2 mg/kg IP, once weekly), systolic dysfunction and dilatation, characterized by increased LVID_{ES} and LVID_{ED} (Fig. 3A, B) and decreased FS (Fig. 3C), were apparent in DNsGCα1^{tg/+} mice, but not WT mice. After 12 weeks of doxorubicin treatment, LV functional impairment and dilatation had further progressed in DNsGCα1^{tg/+} mice. Overall, these data are consistent with the greater doxorubicin-induced systolic dysfunction observed in sGCα1^{-/-CM} mice and suggest that sGC may protect against cardiac dysfunction associated with chronic doxorubicin treatment.

Of note, treating mice with doxorubicin for 12 weeks decreased survival rates to a similar extent in DNsGCα1^{tg/+} and WT mice (67% vs. 56%, respectively, $n=30$ and 34, $p > 0.05$), as well as body weights ($-12 \pm 0\%$ vs. $-15 \pm 1\%$, respectively, $n=18$ and 17, $p > 0.05$).

Doxorubicin-induced cardiotoxicity in mice with induced expression of mutated sGCα1 is attenuated by reversal of sGCα1 mutant expression

The observation that systolic dysfunction occurs within 8 weeks of doxorubicin treatment in DNsGCα1^{tg/+}, but not WT, mice provided a model to test the therapeutic potential of removing sGC activity inhibition by readministering doxycycline. DNsGCα1^{tg/+} and WT mice were exposed to doxorubicin for 12 weeks (2 mg/kg IP, once weekly), and after 8 weeks of doxorubicin treatment, doxycycline was added to the diet of both DNsGCα1^{tg/+} and WT mice for an additional 8 weeks, restoring cardiac sGC activity in DNsGCα1^{tg/+} mice (Fig. 4A).

Cardiac function was assessed by echocardiography after 12 weeks of doxorubicin administration (T_{12 weeks}) and 4 weeks later (T_{16 weeks}). After 12 weeks of doxorubicin treatment, including 4 weeks of doxycycline administration, cardiac dysfunction and dilatation were still more pronounced in DNsGCα1^{tg/+} than in WT mice (Fig. 4B). However, 4 weeks later, functional deterioration and adverse remodeling had progressed further in WT mice, whereas cardiac dysfunction and dilatation were reduced in DNsGCα1^{tg/+} mice (Fig. 4B). These findings suggest that restoring sGC activity can attenuate doxorubicin-induced LV dysfunction in mice.

Morphological changes associated with doxorubicin administration were similar in DNsGCα1^{tg/+} and WT hearts

To assess potential differences in the degree of doxorubicin-induced cardiac atrophy between sGCα1-deficient mice and WT mice, we measured heart weights in both genotypes following 12 weeks of doxorubicin or saline administration. Doxorubicin treatment decreased normalized heart weight by 18% and 24% in sGCα1^{-/-CM} and WT mice, respectively, compared with saline-treated controls (Supplementary Table S3, $p > 0.05$). Similarly, in doxorubicin-treated DNsGCα1^{tg/+} and WT mice, heart weight was decreased by 17% and 15%, respectively, compared with saline-treated mice (Supplementary Table S3, $p > 0.05$).

To further explore the mechanisms underlying the difference in doxorubicin-induced cardiac dysfunction between sGCα1-deficient mice and WT mice, we examined fibrosis, apoptosis, inflammation, and vascular density in hearts of DNsGCα1^{tg/+} and WT mice following 12 weeks of doxorubicin administration.

To investigate cardiac fibrosis, we measured collagen deposition and expression levels of two key mediators of fibrosis: transforming growth factor (TGF)-β1 and connective tissue growth factor (CTGF). CTGF is the profibrotic mediator through which TGF-β1 triggers connective tissue cell proliferation, adhesion, migration, and the synthesis of extracellular matrix. Analysis of Sirius red-stained tissue sections using circularly polarized light revealed comparable deposition of thick, tightly packed, red birefringent collagen fibers as well as thin, loosely assembled, green birefringent collagen fibers in the two genotypes (Supplementary Table S4). Consistent with these results, TGF-β1 and CTGF transcript levels were similar in doxorubicin-treated DNsGCα1^{tg/+} and WT mice (Supplementary Table S4).

In addition, the number of apoptotic cardiomyocytes was similar in DNsGCα1^{tg/+} and WT mice after 12 weeks of doxorubicin treatment. In accordance with these results, cardiac

TABLE 1. GREATER SYSTOLIC DYSFUNCTION IN sGC α 1^{-/-}CM THAN WILD-TYPE MICE AFTER 12 WEEKS OF DOXORUBICIN ADMINISTRATION

	WT + saline (n = 6)	sGC α 1 ^{-/-} CM + saline (n = 5)	WT + DOX (n = 12)	sGC α 1 ^{-/-} CM + DOX (n = 9)	Adjusted p-value
ESV (μ l)	19 $\pm$ 1	19 $\pm$ 1	26 $\pm$ 3	37 $\pm$ 3 ^{†,*}	0.01
EDV (μ l)	43 $\pm$ 2	41 $\pm$ 2	51 $\pm$ 2	58 $\pm$ 2 [†]	0.07
EF (%)	60 $\pm$ 2	58 $\pm$ 3	55 $\pm$ 4	42 $\pm$ 4*	0.03
MAP (mmHg)	104 $\pm$ 5	96 $\pm$ 6	105 $\pm$ 5	102 $\pm$ 4	0.94
E _a (mmHg/ μ l)	3.6 $\pm$ 0.3	3.6 $\pm$ 0.4	4.0 $\pm$ 0.4	3.9 $\pm$ 0.3	1.00
P _{ES} (mmHg)	91 $\pm$ 8	83 $\pm$ 4	97 $\pm$ 4	91 $\pm$ 5	0.67
P _{ED} (mmHg)	2.5 $\pm$ 0.5	2.6 $\pm$ 0.3	3.3 $\pm$ 0.6	4.5 $\pm$ 0.7 [†]	0.11
dP/dt _{max} (mmHg/s)	13,499 $\pm$ 1433	11,995 $\pm$ 601	13,503 $\pm$ 678	10,387 $\pm$ 922*	0.03
dP/dt _{min} (mmHg/s)	-12,054 $\pm$ 1809	-11,420 $\pm$ 950	-13,531 $\pm$ 846	-10,943 $\pm$ 991 [†]	0.21
Tau (ms)	5.1 $\pm$ 0.1	4.9 $\pm$ 0.2	5.6 $\pm$ 0.2	6.3 $\pm$ 0.4 ^{†,*}	0.048
PRSW	83 $\pm$ 9	84 $\pm$ 6	70 $\pm$ 9	49 $\pm$ 7 [†]	0.22
E _{ES} (mmHg/ μ l)	7.4 $\pm$ 0.6	7.3 $\pm$ 1.4	4.0 $\pm$ 0.8	2.3 $\pm$ 0.3 [†]	0.73
EDPVR (mmHg/ μ l)	0.22 $\pm$ 0.03	0.18 $\pm$ 0.02	0.16 $\pm$ 0.02	0.19 $\pm$ 0.02	0.56
HR (bpm)	576 $\pm$ 8	614 $\pm$ 12	528 $\pm$ 9 [†]	530 $\pm$ 14 [†]	1.00

Invasive hemodynamic measurements revealed increased systolic dysfunction in sGC α 1^{-/-}CM compared with WT mice after 12 weeks of DOX administration.

Multiplicity-adjusted *p*-values for comparison between WT + DOX and sGC α 1^{-/-}CM + DOX are reported. [†]*p* < 0.05 versus saline and **p* < 0.05 versus WT + doxorubicin.

bpm, beats per minute; DOX, doxorubicin; dP/dt_{max}, maximum first derivative of developed LV pressure; dP/dt_{min}, minimum first derivative of developed LV pressure; E_a, arterial elastance; EDPVR, end-diastolic pressure volume relationship; EDV, end-diastolic volume; E_{ES}, end-systolic elastance; EF, ejection fraction; ESV, end-systolic volume; HR, heart rate; LV, left ventricular; MAP, mean arterial pressure; P_{ED}, end-diastolic pressure; P_{ES}, end-systolic pressure; PRSW, preload recruitable stroke work; sGC, soluble guanylate cyclase; sGC α 1^{-/-}CM, mice with cardiomyocyte-specific deletion of exon 6 of the sGC α 1 allele; Tau, time constant for isovolumic relaxation; WT, wild-type.

mRNA expression levels of antiapoptotic B cell lymphoma 2 (Bcl-2) and proapoptotic Bcl-2-associated X protein (Bax) did not differ between the two genotypes (Supplementary Table S4).

In addition, the number of infiltrated CD45-positive leukocytes, a measure of cardiac inflammation, was similar in hearts of DNsgc α 1^{tg/+} and WT mice after 12 weeks of doxorubicin administration (relative to the number of cardiomyocyte nuclei; Supplementary Table S4).

Furthermore, vascular density was comparable in DNsgc α 1^{tg/+} and WT hearts after 12 weeks doxorubicin treatment (Supplementary Table S4), suggesting that a lower density of capillaries, supplying cardiomyocytes with oxygen and nutrients, did not contribute to the greater cardiac dysfunction in doxorubicin-treated DNsgc α 1^{tg/+} mice.

Doxorubicin-induced ROS generation is increased in sGC α 1-deficient mice

To consider the possibility that doxorubicin induced higher levels of oxidative stress in sGC α 1-deficient than in WT mice, we investigated the degree of doxorubicin-induced ROS generation in cardiomyocytes isolated from sGC α 1-deficient and WT mice. sGC α 1^{-/-} and WT cardiomyocytes were exposed to doxorubicin (50 μ M) for 3 h and then incubated with chloromethyl derivative of 2',7'-dichlorodihydrofluorescein diacetate (CM-H₂DCFDA). This cell-permeable compound is deacetylated by intracellular esterases and then oxidized by hydrogen peroxide (5) and other oxidants (20) to produce fluorescent 2',7'-dichlorofluorescein (DCF). Three hours after exposure to doxorubicin, greater DCF fluorescence, and thus ROS production, was observed in sGC α 1^{-/-} than in WT cardiomyocytes (Fig. 5A).

Similarly, reactive nitrogen species-mediated tyrosine nitration was more pronounced in sGC α 1^{-/-} than WT hearts 24 h

after doxorubicin administration (Fig. 5B). In addition, higher expression of malondialdehyde (MDA) was observed in DNsgc α 1^{tg/+} than WT hearts after 12 weeks of doxorubicin treatment (Fig. 5C). MDA is a product of lipid peroxidation, frequently initiated by hydroxyl and hydroperoxyl radicals (3).

Taken together, these results suggest that doxorubicin-induced oxidative stress is more pronounced when cardiac sGC activity is reduced. This in turn could lead to enhanced sGC oxidation and decrease in activity and subsequent exacerbation of ROS levels *via* a yet to be identified negative feedback system.

Discussion

The long-term adverse cardiac effects of doxorubicin, which limit the use of this effective chemotherapeutic agent, emphasize the need for novel therapeutic approaches to prevent and treat cardiotoxicity associated with doxorubicin treatment. Our data identify sGC as a potential therapeutic target for doxorubicin-induced cardiotoxicity.

Twenty-four hours after doxorubicin administration, cardiac sGC activity was reduced in mice. Since cardiac sGC expression was not altered by doxorubicin treatment, the observed decrease in activity likely results from post-translational modification of sGC. Oxidative modification of sGC, leading to loss of its prosthetic heme group and generation of NO-insensitive sGC, was previously reported in cardiovascular diseases associated with increased oxidative stress (2, 4, 26, 42, 48). *In vivo* administration of the radical scavenger, tempol, or targeting oxidized and heme-free sGC by adding the sGC activator, BAY 58-2667, *ex vivo* rescued the doxorubicin-associated decrease in cardiac sGC activity, suggesting that doxorubicin-induced oxidative modification of sGC represents an underlying mechanism for the reduction in activity.

FIG. 3. Greater DOX-induced LV dysfunction and dilatation in DNsgCα1^{tg/+} than in WT mice. After 8 weeks of DOX treatment, LVID_{ES} and LVID_{ED} were increased (**A**, **B**) and FS decreased (**C**) in DNsgCα1^{tg/+}, but not in WT mice ($n=20$ each). After 12 weeks of DOX administration, systolic dysfunction and dilatation were more pronounced in DNsgCα1^{tg/+} than in WT mice. * $p<0.05$ versus baseline and † $p<0.05$ versus WT (same time point). DNsgCα1^{tg/+}, mice with cardiomyocyte-specific expression of a dominant negative mutation of sGCα1; LVID_{ES}, left ventricular end-systolic internal diameter; LVID_{ED}, left ventricular end-diastolic internal diameter; FS, fractional shortening.

To determine whether reduced sGC activity contributes to the development of doxorubicin-induced cardiotoxicity, doxorubicin was administered to mice with a cardiomyocyte-specific deficiency in sGCα1. After 12 weeks of doxorubicin treatment, LV systolic dysfunction was more pronounced in sGCα1^{-/-CM} than in WT mice. Similarly, mice with induced expression of a dominant negative mutated sGCα1 (DNsgCα1^{tg/+}) displayed greater LV systolic dysfunction and remodeling than WT mice after 12 weeks of doxorubicin treatment. Together, these data reveal a protective role for sGC against doxorubicin-induced cardiotoxicity.

FIG. 4. Attenuated LV dysfunction in DOX-treated DNsgCα1^{tg/+} mice after reversal of sGCα1 mutant expression. (**A**) Doxycycline was added to the diet of DNsgCα1^{tg/+} and WT mice after 8 weeks of DOX treatment to initiate reversal of sGCα1 mutant expression in DNsgCα1^{tg/+} mice. LV function and dimensions were assessed via TTE after 12 weeks of DOX treatment (T_{12 weeks}) and 4 weeks after halting treatment (T_{16 weeks}). (**B**) After 12 weeks of DOX treatment and 4 weeks of doxycycline administration (T_{12 weeks}), DOX-induced systolic dysfunction and remodeling were still greater in DNsgCα1^{tg/+} than in WT mice ($n=28$ and 21, respectively). However, 4 weeks after cessation of DOX treatment (and 8 weeks of doxycycline administration), systolic function and dilatation had further deteriorated in WT mice, but improved in DNsgCα1^{tg/+} mice. † $p<0.05$ versus 12 weeks and * $p<0.05$ versus WT (same time point). TTE, transthoracic echocardiography.

FIG. 5. Increased DOX-induced reactive oxygen species production in sGC α 1-deficient hearts. (A) DCF fluorescence, and thus reactive oxygen species production, was greater in cardiomyocytes isolated from sGC α 1^{-/-} than from WT mice after *in vitro* exposure to DOX (50 μ M for 3 h, average of three independent experiments). Fluorescence intensity was normalized to protein concentration and expressed relative to levels in untreated cardiomyocytes (*i.e.*, without DOX). (B) In addition, immunoblotting and densitometric analysis revealed higher levels of tyrosine nitration in sGC α 1^{-/-} than WT hearts 24 h after DOX administration (20 mg/kg IP). (C) Similarly, more MDA-protein adducts were observed in DNsgc α 1^{tg/+} than WT hearts after 12 weeks of DOX administration. Immunoblotting for Gapdh allowed to control for sample variability. * p < 0.05 versus WT. DCF, 2',7'-Dichlorofluorescein; MDA, malondialdehyde; 3-NT, 3-nitrotyrosine.

Furthermore, we wanted to explore whether restoring sGC activity could attenuate established cardiac dysfunction. Since DNsgc α 1^{tg/+} mice displayed LV dysfunction and remodeling after 8 weeks of doxorubicin administration, whereas WT mice did not, the impact of restoring cardiac sGC activity on doxorubicin-induced cardiotoxicity could be evaluated by initiating reversal of sGC α 1 mutant expression in DNsgc α 1^{tg/+} mice at this time point. Halting expression of mutated sGC α 1 in DNsgc α 1^{tg/+} mice after 8 weeks of doxorubicin administration, resulting in restored sGC activity, eventually reduced LV dysfunction. Four weeks after halting doxorubicin treatment, cardiac function was even less impaired in DNsgc α 1^{tg/+} mice than in WT animals.

It is possible that mechanisms to compensate for reduced cardiac sGC activity are triggered in the DNsgc α 1^{tg/+} mouse model. When expression of mutated sGC α 1 is halted in these DNsgc α 1^{tg/+} mice, the resulting increase in sGC activity in combination with potential compensatory mechanisms could lead to enhanced cGMP signaling in comparison with WT mice and may contribute to improvement of cardiac function beyond the level observed in doxorubicin-treated WT mice.

Despite the observation that doxorubicin-induced cardiac dysfunction was greater in sGC α 1^{-/-CM} and DNsgc α 1^{tg/+}

mice than in WT littermates, heart weights were similar in doxorubicin-treated sGC α 1-deficient and WT mice. In addition, cardiac Bcl-2 and Bax mRNA expression and cardiomyocyte apoptosis did not differ between DNsgc α 1^{tg/+} and WT mice following 12 weeks of doxorubicin treatment. These results indicate that the increased cardiac dysfunction in doxorubicin-treated sGC α 1-deficient mice did not result from a greater loss of cardiomyocytes. In addition, cardiac fibrosis, inflammation, and vascular density did not differ between sGC α 1 mutant and WT mice following 12 weeks of doxorubicin administration.

We cannot exclude that the degree of cardiac fibrosis, apoptosis, inflammation, or vascular density might differ at earlier or later time points. In addition, the difference in cardiac function between doxorubicin-treated sGC α 1-deficient and WT mice did not reflect an effect of sGC α 1 deficiency on blood pressure, which was similar in doxorubicin-treated sGC α 1^{-/-CM} and WT mice.

Oxidative stress has been suggested to play a prominent role in doxorubicin-associated cardiotoxicity. In mice overexpressing endothelial NO synthase (NOS3), doxorubicin-induced cardiotoxicity was greater than in WT mice due to abundant ROS production resulting from NOS3-mediated

reduction of doxorubicin (34). We also observed increased doxorubicin-induced ROS production in sGC α 1-deficient cardiomyocytes and greater oxidative stress levels in sGC α 1-deficient hearts after *in vivo* exposure to doxorubicin.

It is possible that the more pronounced doxorubicin-induced oxidative stress in sGC α 1-deficient cardiomyocytes leads to greater contractile impairment. Oxidative stress can alter expression, phosphorylation, or function of calcium regulatory proteins such as sarcoplasmic reticulum calcium ATPase 2A and ryanodine receptor 2 (1, 25, 54). Oxidative stress also induces structural modifications of sarcomeric proteins and disrupts mitochondrial function in cardiomyocytes (44). Additional studies are required to evaluate whether increased oxidative stress levels in sGC α 1-deficient cardiomyocytes result in greater contractile impairment and, if so, which mechanisms are involved.

Further support for a protective role of NO-sGC-cGMP signaling in doxorubicin-induced cardiotoxicity was previously provided by the observation that supplementing dietary nitrate reduced doxorubicin-induced cardiotoxicity (57). This cardioprotection was associated with a decrease in ROS generation, lipid peroxidation, and mitochondrial respiratory chain damage (57).

In addition, pharmacological inhibition of the cGMP-catabolizing enzyme, phosphodiesterase type 5 (PDE5), with sildenafil or tadalafil attenuated doxorubicin-associated cardiotoxicity in mice (17, 24). These PDE5 inhibitors reduced doxorubicin-induced oxidative stress, disruption of the mitochondrial membrane potential, apoptosis, and depletion of prosurvival proteins in the heart (17, 24). Because the ability of PDE5 inhibitors to augment cGMP levels is inherently limited by cGMP synthesis, strategies designed to enhance cGMP synthesis may be more effective in preventing doxorubicin-induced cardiotoxicity. Nitroglycerin and related NO donors are currently used to treat various cardiovascular diseases. However, nitrates produce off-target (cGMP-independent) effects and induce tolerance.

As many beneficial effects of NO are mediated through sGC, pharmacological agents directly targeting sGC could represent a novel mechanism-based approach to limit cardiotoxicity in patients undergoing doxorubicin-based chemotherapy. The therapeutic potential of targeting sGC to prevent or even reverse doxorubicin-induced cardiotoxicity is supported by the observation that doxorubicin-induced LV dysfunction in DNsgc α 1^{tg/+} mice is attenuated by removing inhibition of sGC activity. Future studies exploring the therapeutic potential of sGC agonists for the treatment of cardiotoxicity associated with doxorubicin treatment are warranted.

In conclusion, doxorubicin administration reduces cardiac sGC activity in mice, potentially *via* oxidative modification of the enzyme. Our studies in two genetically modified mouse models of cardiac sGC α 1 deficiency demonstrate that reduced cardiac sGC activity contributes to the development of doxorubicin-induced cardiotoxicity, identifying sGC as a plausible therapeutic target to treat this devastating side effect of anthracycline-based chemotherapy. Carefully designed preclinical studies with pharmacological sGC agonists are required to test whether these drugs may provide an additional tool to cardiologists and oncologists for preventing chronic cardiomyopathy and heart failure in doxorubicin-treated patients.

Materials and Methods

Experimental animals

This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Housing and all procedures involving experimental animals were approved by the Institutional Animal Care and Use Committees of Massachusetts General Hospital (Subcommittee on Research Animal Care).

Cardiomyocyte-specific sGC α 1^{-/-CM} mice were generated by crossing mice in which exon 6 of the sGC α 1 fl/fl, generated by our group (7) with mice expressing Cre recombinase under the control of an α -myosin heavy chain promoter (α MHC-Cre^{+/+}). In cardiomyocytes of sGC α 1 fl/fl ^{α MHC-Cre^{+/+}} offspring (referred to as sGC α 1^{-/-CM} mice), Cre recombinase induces deletion of exon 6, encoding a portion of the catalytic domain of sGC α 1 required for enzyme activity. These mice were maintained on a C57BL/6J background (The Jackson Laboratory, Farmington, CT) and sGC α 1 fl/fl ^{α MHC-Cre^{-/-}} littermates served as WT controls.

To establish the DNsgc α 1^{tg/+} mouse model, a first mouse line was generated in which expression of a mutated sGC α 1-encoding gene is driven by a doxycycline-responsive promoter element (tetracycline operator [TetO]). The dominant negative point mutation introduced into the sGC α 1-encoding gene results in an amino acid change in the catalytic region of sGC α 1 (D529A, Asp to Ala) (55). Following implantation of microinjected FVB zygotes into a pseudopregnant Swiss foster mother, offspring were genotyped, and transgenic founders were backcrossed for >6 generations onto a C57BL/6N background (Taconic, Hudson, NY).

This mouse line was crossed with a second mouse line expressing the gene encoding a tetracycline transactivator (tTA) protein under the control of an α MHC promoter, kindly provided by Dr. D.A. Dichek (University of Washington Medical Center, Seattle, WA), and maintained on a C57BL/6N background (Taconic, Hudson, NY).

When doxycycline is removed from the diet (doxycycline-containing chow, 200 mg/kg; Harlan Laboratories, Huntingdon, United Kingdom) of dual heterozygous TetO-DNsgc α 1^{+/+ α MHC-tTA^{+/+}} offspring (referred to as DNsgc α 1^{tg/+} mice), doxycycline-mediated repression of transcriptional activation is abolished (Tet-Off system), and tTA binds to and activates transcription from the TetO promoter element, inducing expression of the sGC α 1 mutant specifically in cardiomyocytes. DNsgc α 1^{tg/+} mice and their WT controls (TetO-DNsgc α 1^{-/- α MHC-tTA^{-/-}} littermates) were conceived and raised in the presence of doxycycline. When the mice were 4–6 weeks old, doxycycline was withdrawn, and 4 weeks later, mice were used for experiments.

Doxorubicin treatment regimen

For acute studies, 8-week-old C57BL/6J mice (The Jackson Laboratory) were injected with a single dose of doxorubicin (20 mg/kg IP, using a 2 mg/ml stock of doxorubicin hydrochloride obtained from Pfizer, New York, NY) or saline and sacrificed 24 h later. A subset of these mice were provided with the radical scavenger, tempol (1-oxy-2,2,6,6-tetramethyl-4-hydroxypiperidine; Thermo Fisher Scientific, Waltham, MA), *via* drinking water (1 mM) 4 days before

doxorubicin or saline treatment and *via* gavage (30 mg/kg) 1 h before treatment.

For chronic studies, 8–12-week-old male and female DN α GC α 1^{tg/+}, sGC α 1^{-/-CM}, and WT mice were administered a low dose of doxorubicin for 12 weeks (2 mg/kg IP, once weekly).

Transthoracic echocardiography

To measure LV systolic function and dimensions, sGC α 1^{-/-CM} mice were lightly sedated (ketamine, 100 mg/kg, IP) and echocardiography was performed using a 13 MHz ultrasound probe (Vivid 7; GE Healthcare, Wilmington, MA). DN α GC α 1^{tg/+} mice were sedated with 1.5% isoflurane and echocardiograms were obtained using an MS400 transducer on a Vevo 2100 scanner (VisualSonics, Inc., Toronto, Canada). LVID_{ED} and LVID_{ES}, interventricular septal thickness at end-diastole, LV posterior wall thickness at end-diastole, and HR were measured and FS calculated, as previously described (39).

Invasive hemodynamic measurements

Mice were anesthetized by IP injection with ketamine (120 mg/kg), fentanyl (90 μ g/kg), and rocuronium (10 mg/kg), intubated, and mechanically ventilated (FiO₂ = 1, 10 μ l/g, 120 breaths per minute). A fluid-filled catheter was introduced into the carotid artery to record arterial blood pressure and HR. After thoracotomy, a pressure–volume conductance catheter (PVR-1030; Millar Instruments, Houston, TX) was introduced through the apex into the LV, as described previously (19).

LV end-systolic and end-diastolic volumes and LV pressures were measured, and EF, arterial elastance, the maximum and minimum first derivative of developed LV pressure (dP/dt_{max} and dP/dt_{min}), and the time constant for isovolumic relaxation (Tau) were calculated. The preload recruitable stroke work, end-systolic elastance, and end-diastolic pressure–volume relationship were obtained by transiently occluding the inferior vena cava. All parameters were measured using LabChart (ADInstruments, Colorado Springs, CO).

sGC activity measurements

sGC enzyme activity was measured as described previously (7). Cardiac tissues were homogenized and supernatants (containing 50 μ g of protein) were incubated for 10 min at 37°C in a reaction mixture with DETA-NO (1 mmol/l) or BAY 58-2667 (100 μ mol/l). cGMP in the reaction mixture was measured using a commercial radioimmunoassay (Cayman Chemical, Ann Arbor, MI). sGC enzyme activity was expressed as picomoles of cGMP produced per minute per milligram of protein in cardiac extract supernatant, normalized to the contemporaneous control group (saline-treated mice when sGC activity was measured in doxorubicin-treated mice and WT mice when sGC activity was measured in sGC α 1-deficient mice).

Histological and immunoblot analysis

Immunohistochemistry was performed as described previously (52). Images were obtained using an Axiovert 200M imaging microscope (Zeiss, Oberkochen, Germany). To assess the degree of cardiac fibrosis, the area of collagen deposition was traced on Sirius red-stained tissue sections using circularly polarized light, allowing evaluation of tightly packed, red birefringent collagen and thin, loosely assem-

bled, green birefringent collagen. The degree of fibrosis was expressed as the area of red or green birefringent collagen relative to the area of the examined LV tissue area.

Apoptotic cardiomyocytes were detected by labeling DNA strand breaks using TUNEL (ApopTag; Millipore, Billerica, MA), according to the manufacturer's instructions. The index of apoptosis was determined by dividing the number of immunolabeled cardiomyocyte nuclei by the area of the examined LV tissue area.

Vascular density was analyzed on BS1-lectin-stained tissue sections (Sigma-Aldrich, St.-Louis, MO) by dividing the number of positively stained small blood vessels by the number of cardiomyocyte nuclei.

Mouse cardiac tissue was homogenized in RIPA buffer containing Halt protease and phosphatase inhibitors (Thermo Fisher Scientific). Homogenates were transferred to Eppendorf tubes, incubated on ice for 20 min, centrifuged at 14,000 g for 10 min, and supernatants collected. Proteins were separated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis, transferred to Immobilon-FL polyvinylidene difluoride membranes (Millipore) by semidry electroblotting, blocked for 1 h in Odyssey blocking buffer (LI-COR Biosciences, Lincoln, NE), and probed overnight with antibodies specific for sGC α 1 (Abcam, Cambridge, MA), sGC β 1 (Sigma-Aldrich), 3-nitrotyrosine (3-NT; Millipore), or glyceraldehyde-3-phosphate dehydrogenase (Cell Signaling Technology, Danvers, MA). Bound fluorophore-coupled secondary antibodies were visualized using the Odyssey system (LI-COR Biosciences).

For MDA immunoblotting, proteins were transferred to a nitrocellulose membrane, which was blocked for 1 h in 5% nonfat milk. Bound primary antibody (Millipore) was detected using a horseradish peroxidase-conjugated secondary antibody (Dako, Heverlee, Belgium) and visualized using ECL Plus Western blotting detection reagents (GE Healthcare, Buckinghamshire, United Kingdom).

For all immunoblots, densitometric analysis was performed using ImageJ software (NIH, whole lane densitometry for 3-NT and MDA immunoblots).

Quantitative real-time polymerase chain reaction

Cardiac expression of genes involved in fibrosis and apoptosis was measured using quantitative real-time polymerase chain reaction (TaqMan or SYBR green PCR master mix; Life Technologies, Carlsbad, CA). Relative mRNA levels were analyzed using the Livak method (29).

Measurements of DCF fluorescence

Cardiomyocytes were isolated from global sGC α 1^{-/-} mice and WT littermates by Langendorff perfusion, as previously described (27). After plating cardiomyocytes onto laminin-coated plates, the cells were incubated with 50 μ mol/l doxorubicin (Sigma-Aldrich) for 3 h, washed with phosphate-buffered saline, and lysed by adding 100 μ l T-PER (Thermo Fisher Scientific). Next, 140 μ l of 50 μ M CM-H₂DCFDA (Life Technologies) was added to 10 μ l cell lysate (96-well plate). Following incubation at 25°C for 10 min in the dark, DCF fluorescence was measured using a spectrofluorometer (BMG Labtech, Ortenberg, Germany), as previously described (16). Fluorescence intensity was normalized to protein concentration and expressed relative to levels in untreated cardiomyocytes (*i.e.*, without doxorubicin).

Statistical analysis

Values are presented as mean $\pm$ standard error of the mean. When comparing two groups, normal distribution of the data was assessed using the Shapiro–Wilk test, and an unpaired two-tailed *t*-test (parametric) or a Mann–Whitney test (nonparametric) was used. Serial echocardiographic data were analyzed by two-way analysis of variance (ANOVA) and Sidak's multiple comparisons test. For these experiments, the required samples sizes were calculated to yield statistical power of ≥ 0.80 ($\alpha = 0.05$). Analysis of pressure–volume measurements was performed via one-way ANOVA and Sidak's multiple comparisons test (when passed the Shapiro–Wilk normality test) or the Kruskal–Wallis test and Dunn's multiple comparisons test (when failed the Shapiro–Wilk normality test). Survival was analyzed via the Kaplan–Meier method.

Acknowledgments

The authors would like to thank Dr. Cornelius Busch and Dr. Pieter Vermeersch for their help in generating the DN_sGC $\alpha 1^{tg/+}$ mice and Dr. David Dichek for providing the α MHC-tTA mice. This work was supported by a Postdoctoral Fellowship from the Belgian American Education Foundation (to S.V.), Scientist Development Grant 10SDG2610313 from the American Heart Association (to E.S.B.), Research Grant from KU Leuven (PF10/014, to S.P.J.), NIH grant R01EY022746 (to E.S.B.), grants from the FWO-Vlaanderen and the UGent-GOA programs (to P.B.), NIH Grant R01DK082971 (to K.D.B., D.B.B.), and the Leducq Foundation (to K.D.B., D.B.B.).

Author Disclosure Statement

No competing financial interests exist.

References

- Adachi T, Weisbrod RM, Pimentel DR, Ying J, Sharov VS, Schoneich C, and Cohen RA. S-glutathiolation by peroxynitrite activates SERCA during arterial relaxation by nitric oxide. *Nat Med* 10: 1200–1207, 2004.
- Ahrens I, Habersberger J, Baumlin N, Qian H, Smith BK, Stasch JP, Bode C, Schmidt HH, and Peter K. Measuring oxidative burden and predicting pharmacological response in coronary artery disease patients with a novel direct activator of haem-free/oxidised sGC. *Atherosclerosis* 218: 431–434, 2011.
- Ayala A, Munoz MF, and Arguelles S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev* 2014: 360438, 2014.
- Boerrigter G, Costello-Boerrigter LC, Cataliotti A, Lapp H, Stasch JP, and Burnett JC, Jr. Targeting heme-oxidized soluble guanylate cyclase in experimental heart failure. *Hypertension* 49: 1128–1133, 2007.
- Brandt R and Keston AS. Synthesis of diacetyldichlorofluorescein: a stable reagent for fluorometric analysis. *Anal Biochem* 11: 6–9, 1965.
- Budworth J, Meillerais S, Charles I, and Powell K. Tissue distribution of the human soluble guanylate cyclases. *Biochem Biophys Res Commun* 263: 696–701, 1999.
- Buyes ES, Sips P, Vermeersch P, Raher MJ, Rogge E, Ichinose F, Dewerchin M, Bloch KD, Janssens S, and Brouckaert P. Gender-specific hypertension and responsiveness to nitric oxide in sGC $\alpha 1$ knockout mice. *Cardiovasc Res* 79: 179–186, 2008.
- Buzdar AU, Marcus C, Smith TL, and Blumenschein GR. Early and delayed clinical cardiotoxicity of doxorubicin. *Cancer* 55: 2761–2765, 1985.
- Chow EJ, Chen Y, Kremer LC, Breslow NE, Hudson MM, Armstrong GT, Border WL, Feijen EA, Green DM, Meacham LR, Meeske KA, Mulrooney DA, Ness KK, Oeffinger KC, Sklar CA, Stovall M, van der Pal HJ, Weathers RE, Robison LL, and Yasui Y. Individual prediction of heart failure among childhood cancer survivors. *J Clin Oncol* 33: 394–402, 2015.
- Desai VG, Herman EH, Moland CL, Branham WS, Lewis SM, Davis KJ, George NI, Lee T, Kerr S, and Fuscoe JC. Development of doxorubicin-induced chronic cardiotoxicity in the B6C3F1 mouse model. *Toxicol Appl Pharmacol* 266: 109–121, 2013.
- Dickey JS, Gonzalez Y, Aryal B, Mog S, Nakamura AJ, Redon CE, Baxa U, Rosen E, Cheng G, Zielonka J, Parekh P, Mason KP, Joseph J, Kalyanaraman B, Bonner W, Herman E, Shacter E, and Rao VA. Mito-tempol and dextrazoxane exhibit cardioprotective and chemotherapeutic effects through specific protein oxidation and autophagy in a syngeneic breast tumor preclinical model. *PLoS One* 8: e70575, 2013.
- Doroshov JH. Anthracycline antibiotic-stimulated superoxide, hydrogen peroxide, and hydroxyl radical production by NADH dehydrogenase. *Cancer Res* 43: 4543–4551, 1983.
- Doroshov JH. Effect of anthracycline antibiotics on oxygen radical formation in rat heart. *Cancer Res* 43: 460–472, 1983.
- Doroshov JH and Davies KJ. Redox cycling of anthracyclines by cardiac mitochondria. II. Formation of superoxide anion, hydrogen peroxide, and hydroxyl radical. *J Biol Chem* 261: 3068–3074, 1986.
- Doroshov JH, Locker GY, and Myers CE. Enzymatic defenses of the mouse heart against reactive oxygen metabolites: alterations produced by doxorubicin. *J Clin Invest* 65: 128–135, 1980.
- Esposito G, Vos M, Vilain S, Swerts J, De Sousa Valadas J, Van Meensel S, Schaap O, and Verstreken P. Aconitase causes iron toxicity in *Drosophila* pink1 mutants. *PLoS Genet* 9: e1003478, 2013.
- Fisher PW, Salloum F, Das A, Hyder H, and Kukreja RC. Phosphodiesterase-5 inhibition with sildenafil attenuates cardiomyocyte apoptosis and left ventricular dysfunction in a chronic model of doxorubicin cardiotoxicity. *Circulation* 111: 1601–1610, 2005.
- Guglin M, Aljayeh M, Saiyad S, Ali R, and Curtis AB. Introducing a new entity: chemotherapy-induced arrhythmia. *Europace* 11: 1579–1586, 2009.
- Ichinose F, Buys ES, Neilan TG, Furutani EM, Morgan JG, Jassal DS, Graveline AR, Searles RJ, Lim CC, Kaneki M, Picard MH, Scherrer-Crosbie M, Janssens S, Liao R, and Bloch KD. Cardiomyocyte-specific overexpression of nitric oxide synthase 3 prevents myocardial dysfunction in murine models of septic shock. *Circ Res* 100: 130–139, 2007.
- Jakubowski W and Bartosz G. 2,7-Dichlorofluorescein oxidation and reactive oxygen species: what does it measure? *Cell Biol Int* 24: 757–760, 2000.
- Jones RL. Utility of dextrazoxane for the reduction of anthracycline-induced cardiotoxicity. *Expert Rev Cardiovasc Ther* 6: 1311–1317, 2008.
- Kalivendi SV, Konorev EA, Cunningham S, Vanamala SK, Kaji EH, Joseph J, and Kalyanaraman B. Doxorubicin

- activates nuclear factor of activated T-lymphocytes and Fas ligand transcription: role of mitochondrial reactive oxygen species and calcium. *Biochem J* 389: 527–539, 2005.
23. Kilickap S, Akgul E, Aksoy S, Aytemir K, and Barista I. Doxorubicin-induced second degree and complete atrio-ventricular block. *Europace* 7: 227–230, 2005.
 24. Koka S, Das A, Zhu SG, Durrant D, Xi L, and Kukreja RC. Long-acting phosphodiesterase-5 inhibitor tadalafil attenuates doxorubicin-induced cardiomyopathy without interfering with chemotherapeutic effect. *J Pharmacol Exp Ther* 334: 1023–1030, 2010.
 25. Lancel S, Zhang J, Evangelista A, Trucillo MP, Tong X, Siwik DA, Cohen RA, and Colucci WS. Nitroxyl activates SERCA in cardiac myocytes via glutathiolation of cysteine 674. *Circ Res* 104: 720–723, 2009.
 26. Lapp H, Mitrovic V, Franz N, Heuer H, Buerke M, Wolfertz J, Mueck W, Unger S, Wensing G, and Frey R. Cinaciguat (BAY 58-2667) improves cardiopulmonary hemodynamics in patients with acute decompensated heart failure. *Circulation* 119: 2781–2788, 2009.
 27. Lim CC, Apstein CS, Colucci WS, and Liao R. Impaired cell shortening and relengthening with increased pacing frequency are intrinsic to the senescent mouse cardiomyocyte. *J Mol Cell Cardiol* 32: 2075–2082, 2000.
 28. Lipshultz SE, Scully RE, Lipsitz SR, Sallan SE, Silverman LB, Miller TL, Barry EV, Asselin BL, Athale U, Clavell LA, Larsen E, Moghrabi A, Samson Y, Michon B, Schorin MA, Cohen HJ, Neuberg DS, Orav EJ, and Colan SD. Assessment of dexrazoxane as a cardioprotectant in doxorubicin-treated children with high-risk acute lymphoblastic leukaemia: long-term follow-up of a prospective, randomised, multicentre trial. *Lancet Oncol* 11: 950–961, 2010.
 29. Livak KJ and Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{−(Delta Delta C(T))} method. *Methods* 25: 402–408, 2001.
 30. Milei J, Boveris A, Llesuy S, Molina HA, Storino R, Ortega D, and Milei SE. Amelioration of adriamycin-induced cardiotoxicity in rabbits by prenylamine and vitamins A and E. *Am Heart J* 111: 95–102, 1986.
 31. Muindi J, Sinha BK, Gianni L, and Myers C. Thiol-dependent DNA damage produced by anthracycline-iron complexes. The structure–activity relationships and molecular mechanisms. *Mol Pharmacol* 27: 356–365, 1985.
 32. Myers CE, McGuire WP, and Liss RH, Ifrim I, Grotzinger K, and Young RC. Adriamycin: the role of lipid peroxidation in cardiac toxicity and tumor response. *Science* 197: 165–167, 1977.
 33. Nakamae H, Tsumura K, Terada Y, Nakane T, Nakamae M, Ohta K, Yamane T, and Hino M. Notable effects of angiotensin II receptor blocker, valsartan, on acute cardiotoxic changes after standard chemotherapy with cyclophosphamide, doxorubicin, vincristine, and prednisolone. *Cancer* 104: 2492–2498, 2005.
 34. Neilan TG, Blake SL, Ichinose F, Raher MJ, Buys ES, Jassal DS, Furutani E, Perez-Sanz TM, Graveline A, Janssens SP, Picard MH, Scherrer-Crosbie M, and Bloch KD. Disruption of nitric oxide synthase 3 protects against the cardiac injury, dysfunction, and mortality induced by doxorubicin. *Circulation* 116: 506–514, 2007.
 35. Olson LE, Bedja D, Alvey SJ, Cardounel AJ, Gabrielson KL, and Reeves RH. Protection from doxorubicin-induced cardiac toxicity in mice with a null allele of carbonyl reductase 1. *Cancer Res* 63: 6602–6606, 2003.
 36. Riad A, Bien S, Gratz M, Escher F, Westermann D, Heimesaat MM, Bereswill S, Krieg T, Felix SB, Schultheiss HP, Kroemer HK, and Tschope C. Toll-like receptor-4 deficiency attenuates doxorubicin-induced cardiomyopathy in mice. *Eur J Heart Fail* 10: 233–243, 2008.
 37. Rudzinski T, Ciesielczyk M, Religa W, Bednarkiewicz Z, and Krzeminska-Pakula M. Doxorubicin-induced ventricular arrhythmia treated by implantation of an automatic cardioverter-defibrillator. *Europace* 9: 278–280, 2007.
 38. Salzer WL, Devidas M, Carroll WL, Winick N, Pullen J, Hunger SP, and Camitta BA. Long-term results of the pediatric oncology group studies for childhood acute lymphoblastic leukemia 1984–2001: a report from the children's oncology group. *Leukemia* 24: 355–370, 2010.
 39. Scherrer-Crosbie M, Ullrich R, Bloch KD, Nakajima H, Nasser B, Aretz HT, Lindsey ML, Vancon AC, Huang PL, Lee RT, Zapol WM, and Picard MH. Endothelial nitric oxide synthase limits left ventricular remodeling after myocardial infarction in mice. *Circulation* 104: 1286–1291, 2001.
 40. Schwartz CL, Wexler LH, Krailo MD, Teot LA, Devidas M, Steinherz LJ, Goorin AM, Gebhardt MC, Healey JH, Sato JK, Meyers PA, Grier HE, Bernstein ML, and Lipshultz SE. Intensified chemotherapy with dexrazoxane cardioprotection in newly diagnosed nonmetastatic osteosarcoma: a report from the Children's Oncology Group. *Pediatr Blood Cancer* 63: 54–61, 2015.
 41. Seif AE, Walker DM, Li Y, Huang YS, Kavcic M, Torp K, Bagatell R, Fisher BT, and Aplenc R. Dexrazoxane exposure and risk of secondary acute myeloid leukemia in pediatric oncology patients. *Pediatr Blood Cancer* 62: 704–709, 2015.
 42. Stasch JP, Schmidt PM, Nedvetsky PI, Nedvetskaya TY, Ak HS, Meurer S, Deile M, Taye A, Knorr A, Lapp H, Muller H, Turgay Y, Rothkegel C, Tersteegen A, Kemp-Harper B, Muller-Esterl W, and Schmidt HH. Targeting the heme-oxidized nitric oxide receptor for selective vasodilatation of diseased blood vessels. *J Clin Invest* 116: 2552–2561, 2006.
 43. Steinberg JS, Cohen AJ, Wasserman AG, Cohen P, and Ross AM. Acute arrhythmogenicity of doxorubicin administration. *Cancer* 60: 1213–1218, 1987.
 44. Steinberg SF. Oxidative stress and sarcomeric proteins. *Circ Res* 112: 393–405, 2013.
 45. Suliman HB, Carraway MS, Ali AS, Reynolds CM, Welty-Wolf KE, and Piantadosi CA. The CO/HO system reverses inhibition of mitochondrial biogenesis and prevents murine doxorubicin cardiomyopathy. *J Clin Invest* 117: 3730–3741, 2007.
 46. Swain SM, Whaley FS, and Ewer MS. Congestive heart failure in patients treated with doxorubicin: a retrospective analysis of three trials. *Cancer* 97: 2869–2879, 2003.
 47. Tebbi CK, London WB, Friedman D, Villaluna D, De Alarcon PA, Constine LS, Mendenhall NP, Spoto R, Chauvenet A, and Schwartz CL. Dexrazoxane-associated risk for acute myeloid leukemia/myelodysplastic syndrome and other secondary malignancies in pediatric Hodgkin's disease. *J Clin Oncol* 25: 493–500, 2007.
 48. Thoonen R, Cauwels A, Decaluwe K, Geschka S, Tainsh RE, Delanghe J, Hochepeid T, De Cauwer L, Rogge E, Voet S, Sips P, Karas RH, Bloch KD, Vuylsteke M, Stasch JP, Van de Voorde J, Buys ES, and Brouckaert P. Cardiovascular and pharmacological implications of haem-deficient

- NO-unresponsive soluble guanylate cyclase knock-in mice. *Nat Commun* 6: 8482, 2015.
49. Thornalley PJ and Dodd NJ. Free radical production from normal and adriamycin-treated rat cardiac sarcosomes. *Biochem Pharmacol* 34: 669–674, 1985.
 50. Timolati F, Ott D, Pentassuglia L, Giraud MN, Perriard JC, Suter TM, and Zuppinger C. Neuregulin-1 beta attenuates doxorubicin-induced alterations of excitation-contraction coupling and reduces oxidative stress in adult rat cardiomyocytes. *J Mol Cell Cardiol* 41: 845–854, 2006.
 51. van Almen GC, Swinnen M, Carai P, Verhesen W, Cleutjens JP, D'Hooge J, Verheyen FK, Pinto YM, Schroen B, Carmeliet P, and Heymans S. Absence of thrombospondin-2 increases cardiomyocyte damage and matrix disruption in doxorubicin-induced cardiomyopathy. *J Mol Cell Cardiol* 51: 318–328, 2011.
 52. Vandenwijngaert S, Pokreisz P, Hermans H, Gillijns H, Pellens M, Bax NA, Coppiello G, Oosterlinck W, Balogh A, Papp Z, Bouten CV, Bartunek J, D'Hooge J, Luttun A, Verbeke E, Herregods MC, Herijgers P, Bloch KD, and Janssens S. Increased cardiac myocyte PDE5 levels in human and murine pressure overload hypertrophy contribute to adverse LV remodeling. *PLoS One* 8: e58841, 2013.
 53. Vrooman LM, Neuberg DS, Stevenson KE, Asselin BL, Athale UH, Clavell L, Cole PD, Kelly KM, Larsen EC, Laverdiere C, Michon B, Schorin M, Schwartz CL, Cohen HJ, Lipshultz SE, Silverman LB, and Sallan SE. The low incidence of secondary acute myelogenous leukaemia in children and adolescents treated with dexrazoxane for acute lymphoblastic leukaemia: a report from the Dana-Farber Cancer Institute ALL Consortium. *Eur J Cancer* 47: 1373–1379, 2011.
 54. Xu L, Eu JP, Meissner G, and Stamler JS. Activation of the cardiac calcium release channel (ryanodine receptor) by poly-S-nitrosylation. *Science* 279: 234–237, 1998.
 55. Yuen PS, Doolittle LK, and Garbers DL. Dominant negative mutants of nitric oxide-sensitive guanylyl cyclase. *J Biol Chem* 269: 791–793, 1994.
 56. Zhou S, Palmeira CM, and Wallace KB. Doxorubicin-induced persistent oxidative stress to cardiac myocytes. *Toxicol Lett* 121: 151–157, 2001.
 57. Zhu SG, Kukreja RC, Das A, Chen Q, Lesnefsky EJ, and Xi L. Dietary nitrate supplementation protects against doxorubicin-induced cardiomyopathy by improving mitochondrial function. *J Am Coll Cardiol* 57: 2181–2189, 2011.

Address correspondence to:

Dr. Emmanuel S. Buys

Department of Anesthesia, Critical Care,
and Pain Medicine

Massachusetts General Hospital Research Institute
and Harvard Medical School
Anesthesia Center for Critical Care Research
55 Fruit Street, Thier 511B
Boston, MA 02114

E-mail: ebuys@mgh.harvard.edu

Date of first submission to ARS Central, October 19, 2015;
date of final revised submission, July 28, 2016; date of acceptance, August 3, 2016.

Abbreviations Used

3-NT	= 3-nitrotyrosine
ANOVA	= analysis of variance
Bax	= B cell lymphoma 2-associated X protein
Bcl-2	= B cell lymphoma 2
bpm	= beats per minute
cGMP	= cyclic guanosine 3',5'-monophosphate
CM-H ₂ DCFDA	= chloromethyl derivative of 2',7'-dichlorodihydrofluorescein diacetate
CO	= carbon monoxide
CTGF	= connective tissue growth factor
DCF	= 2',7'-dichlorofluorescein
DETA-NO	= diethylenetriamine NONOate
DNsGC α 1 ^{tg/+}	= mice with cardiomyocyte-specific expression of a dominant negative mutation of sGC α 1
DOX	= doxorubicin
dP/dt _{max}	= maximum first derivative of developed LV pressure
dP/dt _{min}	= minimum first derivative of developed LV pressure
E _a	= arterial elastance
EDPVR	= end-diastolic pressure volume relationship
EDV	= end-diastolic volume
E _{ES}	= end-systolic elastance
EF	= ejection fraction
ESV	= end-systolic volume
FS	= fractional shortening
Gapdh	= glyceraldehyde-3'-phosphate dehydrogenase
HR	= heart rate
IP	= intraperitoneal
IVS _{ED}	= interventricular septal thickness at end-diastole
LV	= left ventricular
LVID _{ED}	= left ventricular end-diastolic internal diameter
LVID _{ES}	= left ventricular end-systolic internal diameter
LVPW _{ED}	= left ventricular posterior wall thickness at end-diastole
MAP	= mean arterial pressure
MDA	= malondialdehyde
MHC	= myosin heavy chain
NO	= nitric oxide
NOS3	= endothelial nitric oxide synthase
PCR	= polymerase chain reaction
PDE5	= phosphodiesterase type 5
P _{ED}	= end-diastolic pressure
P _{ES}	= end-systolic pressure
PRSW	= preload recruitable stroke work
ROS	= reactive oxygen species
sGC	= soluble guanylate cyclase
sGC α 1 ^{-/-CM}	= mice with cardiomyocyte-specific deletion of exon 6 of the sGC α 1 allele
Tau	= time constant for isovolumic relaxation
tetO	= tetracycline operator
TGF- β 1	= transforming growth factor- β 1
tTA	= tetracycline transactivator
TTE	= transthoracic echocardiography
WT	= wild-type