COMPREHENSIVE INVITED REVIEW

Protein Promiscuity in H₂O₂ Signaling

David Young,¹⁻³ Brandan Pedre,¹⁻³ Daria Ezerina,¹⁻³ Barbara De Smet,¹⁻⁵ Aleksandra Lewandowska,¹⁻⁵ Maria-Armineh Tossounian,¹⁻³ Nandita Bodra,¹⁻⁵ Jingjing Huang,¹⁻⁵ Leonardo Astolfi Rosado,¹⁻³ Frank Van Breusegem,^{2,4,5} and Joris Messens¹⁻³

Abstract

Significance: Decrypting the cellular response to oxidative stress relies on a comprehensive understanding of the redox signaling pathways stimulated under oxidizing conditions. Redox signaling events can be divided into upstream sensing of oxidants, midstream redox signaling of protein function, and downstream transcriptional redox regulation. **Recent Advances:** A more and more accepted theory of hydrogen peroxide (H₂O₂) signaling is that of a thiol peroxidase redox relay, whereby protein thiols with low reactivity toward H₂O₂ are instead oxidized through an oxidative relay with thiol peroxidases.

Critical Issues: These ultrareactive thiol peroxidases are the upstream redox sensors, which form the first cellular port of call for H₂O₂. Not all redox-regulated interactions between thiol peroxidases and cellular proteins involve a transfer of oxidative equivalents, and the nature of redox signaling is further complicated through promiscuous functions of redox-regulated “moonlighting” proteins, of which the precise cellular role under oxidative stress can frequently be obscured by “polygamous” interactions. An ultimate goal of redox signaling is to initiate a rapid response, and in contrast to prokaryotic oxidant-responsive transcription factors, mammalian systems have developed redox signaling pathways, which intersect both with kinase-dependent activation of transcription factors, as well as direct oxidative regulation of transcription factors through peroxiredoxin (Prx) redox relays.

Future Directions: We highlight that both transcriptional regulation and cell fate can be modulated either through oxidative regulation of kinase pathways, or through distinct redox-dependent associations involving either Prxs or redox-responsive moonlighting proteins with functional promiscuity. These protein associations form systems of crossregulatory networks with multiple nodes of potential oxidative regulation for H₂O₂-mediated signaling. *Antioxid. Redox Signal.* 30, 1285–1324.

Keywords: thiol redox, transcription factors, peroxidase signaling, moonlighting functionality

Table of Contents

I. Introduction	1286
II. Applied Techniques for Redox Biology	1287
A. Redox proteomics	1287
B. Structural techniques for studying redox-regulated proteins	1289
C. Genetically encoded redox biosensors	1290
III. Downstream Redox Regulation: Coordinating an Oxidant Response Through Redox-Sensing Transcription Factors	1291
A. Signal transducer and activator of transcription 3	1292
B. Oxidative stress regulator OxyR	1292
C. MarR family	1292

Reviewing Editors: Myra Conway, Alessandro Matte, Helmut Sies, Mark Wilson, Merridee Wouters, and Jing Yang

¹Center for Structural Biology, VIB, Brussels, Belgium.

²Brussels Center for Redox Biology, Brussels, Belgium.

³Structural Biology Brussels, Vrije Universiteit Brussel, Brussels, Belgium.

⁴Department of Plant Biotechnology and Bioinformatics, Ghent University, Ghent, Belgium.

⁵Center for Plant Systems Biology, VIB, Ghent, Belgium.

IV. Midstream Redox Signaling by Moonlighting Proteins	1295
A. Human protein deglycase DJ-1	1295
B. Glyceraldehyde 3-phosphate dehydrogenase	1297
C. Peptidyl-prolyl <i>cis-trans</i> isomerase	1298
D. Transglutaminase 2	1298
V. Upstream Redox Sensing: Prxs as Specific Sensors and Mediators of the Oxidative Stress Response	1299
A. Relationship between redox state and Prx conformation	1300
B. Thiol peroxidase redox relays	1300
C. Nonperoxidatic roles for Prx	1302
D. Complementary roles of Prx and Trx	1305
VI. Conclusion	1306

I. Introduction

CELL SIGNALING AFFORDS ORGANISMS a method of delicate control over cellular balance, and enables the rapid coordination of responses to external stimuli and stressors. Intracellular signaling is formed from a variety of mechanisms, from enzymatic posttranslational modifications of substrate proteins, such as acetylation or phosphorylation, to the sensing of mediatory small molecules, such as Ca^{2+} or cyclic nucleotides. The reversible posttranslational modification of protein thiols presents another axis of cellular signaling, yet despite decades of research, our understanding of oxidative signaling networks remains woefully incomplete. Reactive oxygen species (ROS), such as superoxide ($\text{O}_2^{\bullet-}$), hydroxyl radical ($\bullet\text{OH}$), singlet oxygen (O^1), and peroxide (ROOH), are intracellular initiators of oxidative modifications, and of these, hydrogen peroxide (H_2O_2) is considered most likely to

play a role in oxidative signaling due to its relatively longer half-life and specificity toward protein thiols.

For many years, H_2O_2 within the cell has been regarded as playing a damaging and detrimental role for cell survival, but an appropriate and fine-tuned dose of H_2O_2 is required for normal cellular functioning (303, 338, 339). H_2O_2 is highly oxidizing because of the presence of a peroxide bond (O–O), of which the chemical reduction is rather limited by its high activation energy (419), making H_2O_2 very selective in its reactions with metal centers and specialized protein thiols (404). With the exception of plants, in which the main H_2O_2 source is thought to be glycolate oxidase, H_2O_2 is formed predominantly from $\text{O}_2^{\bullet-}$ dismutation that either happens in a spontaneous ($10^5 \text{ M}^{-1} \cdot \text{s}^{-1}$) or in an activation energy-reduced fashion by the enzyme superoxide dismutase (SOD; $10^9 \text{ M}^{-1} \cdot \text{s}^{-1}$) (98, 99) (Fig. 1). Whereas the cellular electron transport chains are a potential $\text{O}_2^{\bullet-}$ source, the contribution

FIG. 1. Central role of H_2O_2 in sensing, signaling, and regulation. Endogenous sources of H_2O_2 are shown. To prevent damage to lipids, DNA, and proteins, superoxide is rapidly converted into H_2O_2 by compartment-specific SODs. H_2O_2 is produced in several cellular organelles and diffuses across membranes. Cysteine sulfurs react with the redox metabolite H_2O_2 , initiating protein conformational changes (upstream sensing). This information is signaled *via* multiple, often complex, pathways with involvement of several functionally divergent proteins (midstream signaling), and ultimately mechanisms important for the integrity and fitness of the cell are activated (downstream regulation). This results in the upregulation of Prxs, GPxs, and other antioxidant enzymes, which play an active role in the upstream sensing of H_2O_2 . In prokaryotes, H_2O_2 can react directly with oxidant-sensing TF. GPxs, glutathione peroxidases; H_2O_2 , hydrogen peroxide; Prxs, peroxiredoxins; SODs, superoxide dismutases; TF, transcription factors.

of the mitochondrial respiratory chain is generally minor (246, 348). Instead, nicotinamide adenine dinucleotide phosphate [NAD(P)H] oxidases (Fig. 1), also known as rubidium oxidase (RBOH), NADPH oxidase (NOX), and dual oxidase (DUOX), are the main H_2O_2 sources that transfer electrons from NADPH to molecular oxygen (119). These membrane-spanning enzyme complexes generate H_2O_2 through two-electron reduction of oxygen or through one-electron reduction to $\text{O}_2^{\bullet-}$ and subsequent reduction to H_2O_2 . Their activities are controlled by growth factors and cytokines and have an array of physiological and pathophysiological functions (13, 119). In this study, the oxidative burst used to eliminate invading microorganisms highly dependent on NOX activities and mutations that affect the NOX2 activities lead to recurrent infections and impaired pathogen clearance (140, 376). Besides pathogen removal, tissue wounding induces the generation of NOX-dependent H_2O_2 that is required for leukocyte recruitment and wound healing (223, 261). Because of the spatiotemporal differentiation of locally high H_2O_2 concentrations, the H_2O_2 -induced redox signal has to take place in the vicinity of its production, as also recently mentioned in The Redox Code (163). In general, H_2O_2 is a suitable second messenger, because (i) few protein targets are kinetically relevant for H_2O_2 reduction, such as thiol peroxidases and metal centers (404); (ii) it is stable enough to diffuse and generate a gradient from the source (100, 404); (iii) its production is fine-tuned and can be controlled by external stimuli, such as growth factors, insulin, and environmental or mechanical stresses (83, 145, 231); and (iv) its scavenging enzyme activity can be regulated, as observed in peroxiredoxin (Prx) overoxidation (65, 350, 375).

In addition to its role in cellular functioning, intracellular H_2O_2 production is also induced by biotic and abiotic stresses from the cellular environment, thus, bringing the cells under oxidative stress. The term “oxidative stress” appeared in the literature in the mid-eighties (337). Since then, research on oxidative stress responses and the linked redox switches is flourishing. Oxidative stress is now regarded as an imbalance between oxidants and antioxidants in favor of the oxidants, first triggering a redox signaling response, but in the long run resulting in disruption or blocking of signaling pathways, with molecular and cellular damages as a consequence (340).

Despite the highly oxidative nature of H_2O_2 , the rate constants of its reaction with protein thiols vary over several orders of magnitude ($\sim 0.1\text{--}10^8 \text{ M}^{-1} \cdot \text{s}^{-1}$), with some cysteines positioned in pockets catalytically geared toward H_2O_2 reduction. The proteins most catalytically reactive toward H_2O_2 are the primary oxidant sensors of the cell and are therefore likely to play a key role in either the initiation of intracellular signaling or rapid induction of a transcriptional response (Fig. 2). Induction of a transcriptional response can be mediated either by transcriptional regulators capable of directly sensing H_2O_2 (and so contain a “peroxidatic” catalytic cysteine), or through an oxidant signal transduction *via* interaction with a facilitating enzyme catalytically reactive toward H_2O_2 . Both methods of H_2O_2 -mediated transcriptional regulation are discussed within this review in the context of “downstream redox regulation,” and here it is compelling to compare the more simplistic, direct model of oxidant-responsive transcriptional regulation developed by prokaryotes, to the indirect Prx-mediated mode of oxidative regulation adopted by mammalian transcription factors.

As the enzymes most catalytically reactive toward H_2O_2 , Prxs constitute the entry point of oxidative signaling, and in

their recently consolidated role in oxidant transmission, Prxs can be considered as frontline messengers of oxidative signaling, a role discussed within this review in terms of “upstream redox sensing.” Aside from the modulation of transcriptional regulation, signaling through thiol oxidation—Prx facilitated or otherwise—requires specific protein targets to elicit specific functions conducive to the goal of a cellular oxidative response. Considering that H_2O_2 flux can be both rapid, and random, it appears beneficial for cells to possess constitutively expressed proteins, which can alter function in a redox-dependent manner in response to oxidative signaling. Unlike Prxs, which are obligate antioxidant enzymes with a conserved function, such proteins would effectively be redox-responsive moonlighting proteins, and a selection of such moonlighting proteins are the focus of the section Midstream Redox Signaling by Moonlighting Proteins of this review.

II. Applied Techniques for Redox Biology

A. Redox proteomics

Advances in the sensitivity of mass spectrometry (MS) have facilitated an expanding characterization of the redoxome of reversibly oxidized cysteine residues. Recent reviews have highlighted the importance of this technology in the field of redox proteomics (30, 412). Current techniques for the selective enrichment and/or labeling of redox-sensitive cysteines for MS include the on-resin capture of oxidizable cysteines from soluble cell extracts by means of Thiopropyl–Sephadex beads and the adaption of thiol-reactive alkylating agents, such as iodoacetamide (IAM) or *N*-ethylmaleimide (NEM), for quantitative techniques, such as isotope-coded affinity tags (ICATs), or iodoacetyl-based tandem mass tags (iodo-TMTs) (107, 108, 291). These techniques rely on the blocking of free thiols followed by the reduction of oxidized cysteines and subsequent labeling/affinity capture. Whereas ICAT involves the differential labeling of reduced and reversibly oxidized cysteines with “light” and “heavy” ICAT reagents, respectively, iodo-TMT uses a variety of mass reporters, thereby allowing the simultaneous assessment of different oxidant conditions.

A potential drawback to the use of IAM or NEM in the alkylation of free thiols is their crossreactivity with sulfenylated cysteines, although the resulting thioether derivative can be cleaved by reducing agents and, hence, does not necessarily interfere with selective identification of sulfonates (301). A further limitation of IAM-based probes is their potential for toxicity when used at high concentrations (1, 213). In past proteomic approaches, the use of reductants of relative specificity, such as arsenite for sulfenylated cysteines (321) and ascorbate for S-nitrosylated cysteines (153), allowed a partial distinction between oxidative modification types, and iodo-TMT has been resourcefully demonstrated to differentiate sulfenylation, sulfinylation, S-nitrosylation, and S-glutathionylation (355, 405).

In addition to these approaches that employ electrophiles to target thiols, an array of cyclic carbon nucleophiles with specific reactivity to sulfenic acid have arisen based on 5,5-dimethyl-1,3-cyclohexanedione (dimedone) (124). Azide- and alkyne-functionalized “chemical reporter” analogs of dimedone (DAz-1, DAz-2, DYn-1, and DYn-2) for biotinylation successfully identified the proteomics of sulfenylated proteins *in vivo* (207, 269, 276, 300, 331), but the biotin tag had to be photocleaved postenrichment to avoid compromising peptide

FIG. 2. Redox regulation through H_2O_2 -modified protein sulfurs. (A) Cysteine thiols reacting with H_2O_2 to form a sulfenic acid. Identification of sulfenylation in proteins ($-\text{SOH}$) is important, because it presents a crossroad between regulation signaling through thiol switches ($-\text{S-S}-$) and damage signaling through overoxidation to sulfinic ($-\text{SO}_2\text{H}$) and sulfonic ($-\text{SO}_3\text{H}$) acids. (B) Oxidation of some methionine sulfurs and formation of R- and S- sulfoxides ($-\text{SO}-$). These sulfoxides in proteins are also at the crossroad between regulation signaling by methionine sulfoxide reductases and damage signaling by overoxidation to sulfone ($-\text{SO}_2-$). LMW, low-molecular-weight; Msr, methionine sulfoxide reductase; Srx, sulfiredoxin.

ionization in the MS identification (Fig. 3) (413, 414). However, the specificity of dimedone toward sulfenic acids has recently been put to question (97, 136, 137, 353). There has been experimental evidence of dimedone labeling cyclic sulfinyl amides, which can be formed in a sulfenic acid-independent manner *via* the reaction of a glutathionylated thiol or disulfide bond with an amide of the protein backbone (97). A recent study has also demonstrated that a large portion of dimedone-tagged proteins are susceptible to cleavage by dithiothreitol (DTT), indicating that the dimedone-bound species was not

Cys-SOH, but rather cysteine perthiosulfenic acid (Cys-SSOH), the result of the oxidation of a persulfidated thiol (136). Moreover, there have been some indications of dimedone treatment *per se* leading to increased intracellular “ROS” levels, as detected by dichlorofluorescein (DCF) fluorescence (288). Hence, it is strongly advised to confirm proteins detected as sulfenylated by dimedone-based probes using other methods.

As an alternative to chemical approaches to sulfenylation trapping, a genetically encoded probe has been developed. This probe, termed YAP1C, is based on the C-terminal

FIG. 3. In vitro or in vivo sulfenome trapping with dimedone-based chemical probes. Specific trapping of the sulfenylated proteins *in vitro* or *in vivo* by dimedone-based chemical probes (DAZ-1, DAZ-2, DYn-1, and DYn-2). A biotin can be added to the tagged protein by a click reaction between alkyne and azide. The biotinylated targets can be visualized by Western blot with an anti-streptavidin antibody. Alternatively, the targets can be enriched on streptavidin beads, after identification by MS. DA, azide-functionalized dimedone; DY, alkyne-functionalized dimedone; MS, mass spectrometry.

region (residues 565 to 650) of the *Saccharomyces cerevisiae* (baker's yeast) transcription factor yeast API-like protein (YAP1) (264, 364, 365, 400), with only Cys598 retained and other cysteines mutated to alanine or threonine (364). Cys598 of YAP1C has proven to specifically form stable mixed disulfides with sulfenylated cysteines, resulting in a protein complex that can be isolated through inclusion of an affinity tag to YAP1C (264, 364, 365, 400). The major advantages of the YAP1C probes when compared with the more common chemical-based approaches to trap sulfenylated proteins are that they are expressed in the cell, thus directly circumventing any permeability issues, and that they can specifically be targeted to tissues or organelles. Potential disadvantages of the YAP1C probes include the necessity for genetic modification of the target organism and the sensitivity of the mixed disulfide to cleavage by endogenous cellular reductants.

For proteomic characterization of reversible disulfides with physiological significance, thioredoxin (Trx) has recently been utilized as a tool for specific capture of cellular redox targets (8, 206, 254, 278, 283, 421). Both the tandem mass tag (TMT) and ICAT proteomic methods have been adapted to identify Trx targets, merely by using Trx as the postalkylation reductant instead of a chemical reducing agent, such as DTT or tris(2-carboxyethyl)phosphine (TCEP) (278, 421). Other *in vivo* approaches involve replacement of endogenous Trx with a resolving cysteine mutant of Trx, which forms a stable mixed disulfide with oxidized cysteines of the target proteins. The resulting Trx–target complexes have been enriched by immobilized metal affinity chromatography, in which a poly-His tag has been engineered (283), Trx-Sepharose resin (8, 206, 278), or by immunoaffinity (254). Proteins captured in a mixed disulfide with Trx can elute with a reducing agent and can be separated by two-dimensional gel electrophoresis for MS identification (8, 206, 254, 278, 283).

A similar approach has been used to identify intracellular targets of *Arabidopsis thaliana* 2-Cys peroxiredoxin A (PrxA), with Prx–target complexes isolated by coimmunoprecipitation, followed by separation on sodium dodecyl sulfate–polyacrylamide gel electrophoresis and identification by nano liquid chromatography–tandem MS (42). In addition to the detection of covalently bound partners of Trx or Prx, the determination of transient interaction partners has been attempted through techniques that do not depend on the mixed disulfide stability. A classic and well-established technique is the yeast two-hybrid system that fuses “bait” and “prey” sequences to mutually required domains of a transcriptional activator of a reporter gene (91). Interaction of bait and prey results in nuclear translocation and transcription of the reporter gene, thereby providing a readable output. A yeast two-hybrid system has been used to identify an interaction between the *Salmonella* virulence-related effector, SlrP (small leucine-rich protein [E3 ubiquitin-protein ligase]), and human Trx (16). Two-hybrid approaches have also been adapted for use in mammalian cell lines, such as the KInase Substrate Sensor (KISS) assay that exploits a kinase pathway, involving glycoprotein 130 (gp130), nonreceptor tyrosine-protein kinase 2 (TYK2), and signal transducer and activator of transcription 3 (STAT3), to control STAT3-dependent reporter gene expression in response to the bait–prey interaction (211).

More recently developed techniques for assessing protein–protein interactions include proximity-based labeling meth-

ods, such as proximity-dependent biotinylation. Site-directed mutation of the biotin/bioadenosine monophosphate (AMP)-binding domain of the biotin ligase, BirA, enabled the development of techniques based on promiscuous biotinylating fusion proteins, such as BioID and BioID2 (55, 173, 312, 313). The intracellular expression of a BirA fused with the bait protein allows highly sensitive labeling of any transiently interacting partner proteins in a proximity-dependent manner. Biotinylated proteins can then be efficiently enriched by avidin or streptavidin for MS-based identification. Although this technique provides a sensitive tagging method of noncovalent redox-mediated interactions without omission of any transient mixed disulfide complexes, its evident drawback is the non-specific nature of the labeling of the proteins in the immediate proximity of the bait, instead of the discerning labeling of direct interaction partners only. In an attempt to overcome this, a derivative of BioID that incorporates protein fragment complementation was developed. In this approach, BirA is split into two fragments that are fused to two interaction partners; hence, BirA-mediated biotinylation of vicinal proteins occurs in a more defined manner, only when and where the two proteins interact (67, 326). Recently, BioID has been applied in the identification of redox-related interactions in the study of interacting partners of Trx-interacting protein (Txnip) in a mammalian cell line (101). Of the 31 interacting partners identified, 17 were found to be independent of mixed disulfides through the additional use of a Txnip-BioID control, in which the redox-active cysteine of Txnip (Cys247) had been mutated to serine.

B. Structural techniques for studying redox-regulated proteins

X-ray crystallography is the preferred technique for the determination of three-dimensional macromolecular protein structures, which can provide specific insight into the molecular features that govern redox sensitivity of thiols/disulfides. Atomic resolution crystallography has enabled the visualization of the intermediate oxidation states of peroxidic cysteines (280) and facilitated the analysis of disulfide strain energy and other structural determinants of disulfide redox potentials (309, 325). However, practical issues can arise when attempting to crystallize redox-sensitive proteins, as partial oxidation of proteins may result in mixed populations of conformational or oligomeric states with variable stabilities, which obstructs crystallization. The utilization of either reducing or oxidizing agents can successfully drive the protein toward a single redox state, although the choice of the reducing agents can be an important factor, with TCEP generally favored over DTT, because of its enhanced aqueous stability and efficacy over a wide pH range (112). Once crystals of a redox-sensitive protein are obtained, further complications can occur from the use of synchrotron radiation to collect diffraction data. Typical synchrotron diffraction experiments with photon flux in the range of 10^{11} – 10^{12} photons per second subject crystals to a radiation dosage in the order of 10^7 Gy ($\text{J} \cdot \text{kg}^{-1}$), with the absorbed dose depending on the X-ray wavelength and the atomic composition of the protein (249).

One of the first evidence of specific radiation damages to a protein is the electron capture-induced cleavage of disulfide bonds (5), which is particularly undesirable in the case of functional redox-active disulfides. The sensitivity of disulfide bonds toward ionizing radiation varies greatly and depends

generally on the local microenvironment and, with the exception of Trx1 from *Litopenaeus vannamei* (white-leg shrimp), increases with solvent accessibility (36, 297, 401). In a case study of lysozyme crystals, saturated radicalization of disulfide bonds was observed at a cumulative dose of 0.5–0.7 MGy, followed by bond breakage at 1.05 MGy (357). In addition to the radiolytic cleavage of disulfides, oxidation of protein thiols by radiation has also been proposed, although its occurrence in protein crystals is poorly characterized. Oxidation of thiol sulfurs by ionizing radiation has been suggested to proceed *via* a thiyl radical ($\text{RS}^\bullet$) intermediate, which can be formed either by direct electron capture at the thiol or through reaction with a hydroxyl radical ($^\bullet\text{OH}$), which is formed upon radiolysis of water. The thiyl radical can then react with molecular oxygen to form a persulfenic acid, which then either rearranges to a sulfinic acid (RSO_2H), or condenses to a sulfenic acid (RSOH) (409). Radiation-induced oxidation of cysteine to its sulfenylated form have been proposed to occur in the crystal structures of rat trypsin (402) and isocyanide hydratase (188), and the radiation-induced thiol oxidation of yeast alcohol dehydrogenase in solution has been characterized (307). MS analysis of radiation-exposed cysteinyl peptides suggests a predominant oxidation species of overoxidized $\text{R-SO}_2\text{H/R-SO}_3\text{H}$ forms over an R-SOH (409). Structural biologists wishing to identify physiologically relevant cysteine oxidation in crystal structures should always consider the possibility of artefactual cysteine oxidation during X-ray diffraction data collection.

A possible alternative to X-ray crystallography for structural studies is nuclear magnetic resonance (NMR) spectroscopy. NMR allows for solution study of small proteins or protein fragments, and has the benefit of bypassing the need for crystallization, although the high concentrations of protein required for this technique can sometimes lead to aggregation. NMR can be applied effectively to study redox-dependent structural dynamics of proteins that adopt different conformations according to the redox status and are prone to partial oxidant-induced unfolding, as seen for heat shock protein 33 (Hsp33) (117, 205). Small-angle X-ray scattering (SAXS), as a complementary biophysical technique, has also provided useful insights into redox-driven structural changes, as for instance, into the large-scale oxidation-induced structural rearrangement of the mycothiol-dependent oxidoreductase, Rv2466c (2), later identified as mycoredoxin-2 (310).

Early applications of NMR spectroscopy enabled the first structural characterization of cysteine sulfenylation (59), disulfide bond formation (332), and thiol pK_a determination by monitoring chemical shift changes (96). In view of the increasing number of redox-regulated protein–protein interactions revealed by proteomic studies (42), together with the apparent role of scaffold proteins in facilitating peroxidase redox relays (17), there is a clear need for the structural characterization of transient protein complexes specific to redox processes. Crystallographic studies of such complexes are limited by both the required stability (relatively high affinity) and reasonable conformational homogeneity of the complex. However, with the development in diamagnetic and paramagnetic NMR spectroscopy techniques, characterization of transient protein interactions through NMR may increasingly become an applied field (221, 320).

As cryo-electron microscopy (cryo-EM) has not the same restrictions of conformational homogeneity due to the computational sorting and filtering of sample images, it lends

itself well to the structural study of multiprotein complexes (80, 164, 165). In addition to nonnative complexes, cryo-EM is also especially well suited to analyze large native macromolecular assemblies of subunits. Examples of redox-regulated functions characterized by cryo-EM include the observation of the self-assembling filamentous high-molecular-weight (HMW) chaperone form of PrxIII (293, 416) and the involvement of CXXC motifs of the viral chaperone protein UL32 in disulfide bond regulation (3).

Methodological improvements in sample preparation and image processing software and technological advancements in phase plates (for in-focus phase contrast) and detector camera qualities have evolved single-particle cryo-EM into a viable tool for structural study at the molecular level (114). The resolution of electron microscopy density maps has increased remarkably over the years, with a recent determination of the structure of glutamate dehydrogenase at 1.8 Å resolution (236), and will progress even further (390). As an alternative to X-ray crystallography, single-particle cryo-EM also bears the advantage of being more amenable to the study of membrane proteins, as detergent micelles are often obstructive to crystal formation (although this issue is partially overcome by advances in the usage of lipidic cubic phase crystallization) (134). Approximately 30% of proteins are membrane bound (392), yet they comprise only 3% of the crystal structures and 10% of the cryo-EM structures (299). There is an increasing recognition of redox-regulated mechanisms that drive ion channel conductance in response to oxidative stress (26, 398). A distinct lack of structural information regarding redox-regulated ion channels could be addressed by the expanding prevalence of cryo-EM application in the structural biology field, with a shortage of trained users and affordable access to high-quality electron cryomicroscopes as a current bottleneck (390).

C. Genetically encoded redox biosensors

To assess the biological implications of redox-related processes, it is critical to be able to monitor specific redox species and changes in their levels, preferably with a subcellular resolution. One set of methods involves the use of cell-permeable dyes that react with cellular ROS, leading to fluorescence, of which the signal intensity is proportional to the ROS level (116). Traditional dyes such as 2',7'-dichlorofluorescein diacetate (DCFDA) and its derivatives are plagued by several problems: (i) they lack specificity toward the different forms of ROS, (ii) their subcellular targeting is not specific, (iii) they do not directly target the H_2O_2 levels, (iv) their reactions are irreversible, (v) the nonspecific reactions of the probes may affect their signal with misleading interpretation of the results, and (vi) as they are intensimetric, differences in dye uptake between different cells could be misinterpreted as differences in ROS levels (27, 399). New generations of H_2O_2 probes based on boronate caging of the fluorophore have partially helped to overcome these problems. Such dyes are highly specific toward H_2O_2 , can be targeted to different organelles, and display different fluorescence spectra (73–75, 238), however, reversibility remains a problem.

Redox-sensitive fluorescent proteins have helped to overcome limitations presented by chemical dyes and offer a powerful method for monitoring levels of redox species, as well as investigating the dynamics of protein oxidation and reduction *in vivo*. Fluorescent proteins, such as green

fluorescent protein (GFP), are rendered redox sensitive (roGFP) by the introduction of two cysteine residues onto the surface of the β -barrel. Upon oxidation, the two cysteines form a disulfide bond that induces slight conformational changes, which in turn change the protonation state of the fluorophore, leading to a simultaneous increase of the excitation peak at 400 nm and a decrease of the excitation peak at 480 nm, that is, to a ratiometric response. This property renders the probe readout independent of its concentration or expression levels, which may vary between different cell types and compartments (77,128). Moreover, the fact that the fluorophore is shielded by the β -barrel makes the probe pH independent in the physiological range (328). However, the roGFP probe itself is not specific for the oxidant source and it is not very sensitive or rapidly responding to changes in the cellular redox status.

To overcome these limitations, roGFP is fused, *via* a short peptide linker, to proteins that are more sensitive and selective to the oxidative source. Mostly, the low-molecular-weight (LMW) thiol-specific redoxins (*e.g.*, glutaredoxins, mycoredoxins, and bacilliredoxins) are used to detect changes in the intracellular LMW thiol redox balance or yeast thiol peroxidases that directly detect changes in the H_2O_2 levels (19, 125, 126, 222, 246). The intramolecular disulfide formation in roGFP is slightly different in both fusions: for the LMW thiol redox probes, the LMW thiol is transferred from the LMW thiol redoxin to roGFP, whereas for the thiol peroxidases, a disulfide exchange occurs (oxidant receptor peroxidase 1 [Orp1]) or as an yet-to-be-confirmed mechanism (thiol-specific antioxidant 2 [Tsa2]) (19, 125, 126, 222, 246). Furthermore, another application of roGFP2 fusions is seen in a recent study, in which redox catalysis was investigated (351). Here, roGFP2 was fused to the Prx5-type model enzyme antioxidant protein (AOP) from *Plasmodium falciparum*, and expressed in cells. Kinetic data showed a clear correlation between the roGFP2 readouts and the recombinant $PfAOP$ k_{cat}^{app} values.

Another group of genetically encoded redox probes is fused with a circularly permuted yellow fluorescent protein (cpYFP). By coupling it to Trx and to the stereospecific methionine sulfoxide reductases (MsrA or MsrB), cpYFP is used to detect high stereospecific methionine sulfoxide oxidations (367). The first example of a cpYFP probe (designated HyPer) for detection of changes in intracellular H_2O_2 levels is HyPer (14). In HyPer, the cpYFP probe is integrated into the C-terminal regulatory domain (RD) of the *Escherichia coli* oxidative receptor OxyR, in the region between the peroxidatic and resolving cysteine (C_R) (14). In contrast to the roGFP probes, the modification in the cpYFP fluorometric properties is not due to an intramolecular disulfide in the probe, but to the conformational changes in OxyR upon disulfide formation that induce structural modifications in cpYFP (14). Upon oxidation with HyPer, the 420-nm excitation peak decreases and the 500-nm peak increases (14).

New versions of HyPer have been developed with an improved dynamic range or with a different fluorescent signal, and in some cases combined with a fully genetically encoded system for localized H_2O_2 production by the activation of the yeast D-amino acid oxidase (DAAO) (23, 25, 82). Of note, the HyPer probe was employed as a template for the development of the NADPH probe iNap (366), where cpYFP was inserted into the NAD(H)-binding domain of Rex from

Thermus aquaticus. As NADPH serves as the main source of reducing equivalents for the reduction of redox-regulated proteins, including the genetically-encoded biosensors, monitoring its levels is also important when investigating oxidative signaling.

The probes described in this study are ratiometric, allowing an efficient readout, regardless of the probe expression level. As of today, the roGFP-Tsa2 probe is by far the most sensitive H_2O_2 -sensing probe reported, in agreement with the high reactivity of Prxs ($10^7 M^{-1} \cdot s^{-1}$) (246). HyPer and roGFP-Orp1 have a similar H_2O_2 sensitivity ($10^5 M^{-1} \cdot s^{-1}$), with HyPer responding slightly faster than roGFP-Orp1 (126). In contrast, both HyPer and roGFP-Orp1 return more quickly to the resting state than roGFP-Tsa2 (246). A general disadvantage of HyPer over roGFP-based probes is the intrinsic sensitivity of the cpYFP chromophore to pH changes. A H_2O_2 -insensitive probe, designated SypHer2, has been generated as a control to tackle this problem (233, 336).

A general concern of these probes is their possible interference with H_2O_2 homeostasis, because they possess an antioxidant activity. However, the roGFP2-Tsa2 probe shows a very low H_2O_2 -scavenging capacity, whereas HyPer does not seem to affect redox-dependent physiological reactions (22, 246). Another concern is related to the role of peroxidases, which act as H_2O_2 signal transducers (70, 345, 353), as the introduced thiol peroxidase in roGFP probes might lead to the activation of H_2O_2 -signaling pathways. With HyPer, these secondary effects are less likely to happen in eukaryotic cells, due to the bacterial origin of the OxyR-regulatory region. From a more practical approach, usage of genetically encoded redox biosensors may be limited to cells that are difficult to transfect or transduce, such as primary or suspension cells; in these situations, new generations of H_2O_2 chemical dyes may still be the preferred option of monitoring H_2O_2 levels. Moreover, currently all ratiometric fluorescent probes (roGFP and cpYFP) have similar excitation spectra, which compromises such applications as the simultaneous monitoring of H_2O_2 in different compartments of the same cell, for example. However, the recent development of a red fluorescent H_2O_2 probe, HyPer Red (82) and of a redox-sensitive red fluorescent protein, rxRFP (87) could be employed together with cpYFP and roGFP2-based biosensors for this purpose.

III. Downstream Redox Regulation: Coordinating an Oxidant Response Through Redox-Sensing Transcription Factors

The ability of a cell to survive under oxidative stress conditions depends on its ability to rapidly adapt its transcriptional response to fill itself with antioxidant enzymes (Fig. 1). This responsive capacity is best displayed in prokaryotes and it is the cornerstone that underpins the survival of pathogenic bacteria upon exposure to oxidants released by the mammalian immune system. In this study, we discuss two redox-regulated prokaryotic transcription factors, OxyR and the multiple antibiotic resistance regulator (MarR), which exemplify differing, yet effective, modes of oxidant-induced structural changes that modulate their association to cognate DNA. In comparison, redox-regulated mammalian transcription factors, such as the STAT family, are less effective in their direct response to cellular oxidants and, instead, rely

on redox-sensitive signaling effectors, such as kinases or peroxidases to mediate an oxidative stress reaction.

A. Signal transducer and activator of transcription 3

STAT proteins are transcriptional regulators of signaling factors involved in cell survival and proliferation (64). Upon phosphorylation by kinases, such as Janus kinase 2 (JAK2), the cytoplasmic STAT3 dimerizes and translocates to the nucleus to activate transcription (64). Tyrosine-phosphorylated STAT3 plays a role in the control of ROS production through down-regulation of mitochondrial proteins of the electron transport chain, thereby reducing ROS leakage (71), and upregulation of the expression of ROS-consuming enzymes, such as mitochondrial SOD (257). STAT3 contains multiple oxidant-sensitive cysteines and under oxidative stress, it oligomerizes through intermolecular disulfides at its DNA-binding domain (DBD) and C-terminal transactivation domain (C-TAD) to form dimers, trimers, and tetramers, which inhibit DNA binding (208, 209). As STAT3 also exerts a regulatory effect on the activities of the electron transport chain complexes, STAT3 oxidation has been postulated to promote its regulatory role in mitochondria (333, 359, 360). Mitochondria-localized STAT3 associates with cyclophilin D (CypD) upon oxidation in an interaction proposed to stabilize the oxidized STAT3 form (234). JAK2 is also sensitive to oxidative inactivation, because oxidation of two cysteines in the “N-lobe” of the catalytic site triggers enzymatic inactivation (344). These cysteines are located 9 Å apart, but whether intramolecular disulfide bond formation or sulfenylation alone mediates the reversible inactivation of this kinase is still unknown (79). As a STAT3 activator, the redox sensitivity of JAK2 adds a supplemental, indirect mode of redox regulation to STAT3. A direct redox relay between a Prx and STAT3 has been identified and will be discussed in the Thiol peroxidase redox relays section (345).

B. Oxidative stress regulator OxyR

OxyR is a prokaryotic transcription factor that regulates several genes important for the cellular redox control, such as catalases and ferritins (148, 423). OxyR is considered to be an important resistance factor against the oxidative burst of innate immune responses due to its influence over virulence factors, such as biofilm formation or secretion systems (33, 162, 229, 330, 394). It belongs to the LysR transcription-type regulator family, of which the members contain a conserved N-terminal DBD, and a C-terminal RD. OxyR contains two conserved cysteines in the RD that are separated by nine amino acids within the sequence and ~ 15 Å spatially (53, 161, 315, 358). The N-terminally conserved cysteine is located in a hydrophobic pocket and is highly reactive toward H_2O_2 , with a rate constant of $10^5 \text{ M}^{-1} \cdot \text{s}^{-1}$ (53, 161, 194, 315, 358). Upon oxidation, the nascent sulfenic acid is believed to induce unfolding of the α -helix, allowing the C-terminally conserved cysteine to come in the proximity to form an intramolecular disulfide bond (53, 194, 358). The changes in the tertiary structure of OxyR due to the intramolecular disulfide are translated into the quaternary structure. OxyR is a tetramer in solution and its RDs form tightly associated homodimers (with a buried surface area of $>2000 \text{ Å}^2$) (53, 358). In the *E. coli* OxyR, disulfide bond formation provokes a relative rotation of 30° between the protomeric subunits of the RD homodimer, which

has an allosteric effect on the DBD subunits and so affects the DNA contacts and the interaction with the RNA polymerase, thereby modulating transcription (Fig. 4) (53).

In contrast, the RD of OxyR from *Porphyromonas gingivalis* displays only a very small interprotomer rotation upon disulfide formation and both the reduced and oxidized conformations resemble the oxidized RD of *E. coli* OxyR (358). This demonstrates that the redox-induced conformational changes exerted by the peroxidatic cysteine (C_P)- C_R disulfide switch off are not conserved among OxyRs. Accordingly, the modifications in the OxyR-binding topology and association with RNA polymerase upon oxidation are quite variable among OxyRs from different species. For instance, when binding to the *oxyRS* promoter region, the reduced *E. coli* OxyR binds two pairs of adjacent major grooves separated by one helical turn, provoking DNA bending, whereas the oxidized OxyR binds to four consecutive major grooves, eluding DNA bending (374). A similar mechanism is observed in the *oxyR2-PrxII* promoter region of the bacterium *Vibrio vulnificus* (178). The genes in *oxyRS* in *E. coli* and *oxyR2-PrxII* in *V. vulnificus* are divergently transcribed, possibly because these regions share a redox-related transcriptional control. In the binding to the alkyl hydroperoxide *C* (*ahpC*) promoter region of *E. coli*, the reduced and oxidized *E. coli* OxyR factors have the same DNA contacts, but the oxidized OxyR has an increased affinity and activates *ahpC* transcription (374). In other cases, OxyR acts as a repressor, although the binding mechanism has not been studied in detail (176, 370, 381). In any case, the conformational changes induced by oxidation and intramolecular disulfide bond formation have dramatic effects on the binding affinity of the OxyR toward its cognate promoter regions, resulting in an altered transcriptome (Fig. 4). In all cases, the allosteric control of the DBD is conferred through oxidative modifications of the RD, with the presently held hypothesis that the localized structural changes produced by intramolecular disulfide formation induce a subtle alteration in the quaternary structure that translates into a change in the relative DBD orientation.

C. MarR family

The MarR family comprises a wide variety of species-specific transcriptional regulators, some of which are responsive to intracellular oxidants. MarR family proteins are homodimers that contain a structurally conserved winged helix-turn-helix (wHTH) DNA-binding motif on each protomer (4). Typically, they act as transcription repressors through steric hindrance of the RNA polymerase to a promoter, although some family members act as activators by either competitive exclusion against repressors or by stabilizing the RNA polymerase (377). MarR proteins possess a dimerization region and form relatively compact homodimers through intercalation of the $\alpha 1$, $\alpha 5$, and $\alpha 6$ helices, all while leaving the wHTH free for DNA recognition (29). In MarR-type structures, the dimerization domain functions as a hinge and any movement in the dimeric interface communicates an equivalent shift in the relative distance between the partnered wHTH DNA-binding motifs, triggering a structural synchronicity in which the MarR protein specificity and affinity toward DNA are modulated (186, 424). Despite their origin from a common ancestor, MarR family proteins have evolved multiple distinct redox regulation mechanisms.

FIG. 4. Proposed DNA-binding and transcriptional control mechanisms of OxyR. (A) Under reducing conditions, transcription repression by OxyR of the targeted gene by DNA bending of the promoter region, preventing RNA polymerase binding. OxyR oxidation causes DNA unbending of the DNA. The oxidized OxyR cooperatively associates with the RNA polymerase, activating transcription. (B) Under reducing conditions, lack of promoter region binding by OxyR. Upon oxidation, OxyR binds to the promoter and activates transcription cooperatively with the RNA polymerase. (C) Under reducing conditions, OxyR binding to the promoter region and repression of transcription by preventing binding of the RNA polymerase. Oxidation relaxes the DNA binding and allows the RNA polymerase to transcribe the gene. OxyR, oxidative receptor.

The MarR-type transcription regulator, organic hydroperoxide resistance regulator (OhrR), adopts species-specific modes of redox regulation, depending on the presence of a single (typically N-terminal) cysteine, or two or more cysteines. The 1-Cys variant OhrR of *Bacillus subtilis* is responsive to oxidative modification by LMW thiol conjugation. Sulfenylation of *B. subtilis* OhrR does not weaken its DNA-binding affinity, instead, mixed disulfide formation (with bacillithiol, coenzyme A, or cysteine) or condensation with the amide backbone to a sulfenamide is required for derepression (199). The 2-Cys variant of *Xanthomonas campestris* OhrR also has an oxidant-sensing N-terminal cysteine (Cys22) that, upon oxidation, engages in an intersubunit disulfide bond with a C_R (Cys127) of the neighboring protomer (270). It should be noted that referring to such OhrRs as “2-Cys” does not preclude the existence of more than two cysteines; *X. campestris* OhrR has an additional cysteine in the proximity of Cys127 (Cys131) that is not considered functionally significant. Structural characterization of the reduced and oxidized forms of *X. campestris* OhrR revealed that, upon oxidation of Cys22 to sulfenic acid, a hydrogen bonding network involving neighboring tyrosines is disrupted allowing a 135° rotation and 8.2 Å translation of Cys127 of the partner protomer to engage in a disulfide (258).

This localized structural transition results in a striking reorganization of the dimerization interface, whereby the stacking arrangement of the α6-helices' interface is effectively swapped, reversing the helix polarity while maintaining the pseudo twofold symmetry and overall triangular shape of the homodimer (Fig. 5). The rearrangement of the dimerization region conveys a 28° rotation of the wHTH domains, weakening the

affinity of the oxidized OhrR for its target promoter. Whereas the C-terminal C_R of *X. campestris* OhrR appears to be essential for oxidative regulation of its DNA-binding capacity, the C-terminal cysteines of other 2-Cys orthologs, such as the OhrR of *Pseudomonas aeruginosa* or *Chromobacterium violaceum*, are dispensable for oxidative derepression and seemingly act as a protection against overoxidation (9, 61). In the case of the *P. aeruginosa* OhrR homolog, OspR (bearing 46.5% identity to OhrR), oxidation of the N-terminal cysteine is sufficient to abolish its affinity to the *ohr* promoter, but both N- and C-terminal cysteines are required for oxidative regulation of its affinity to the glutathione peroxidase (*gpx*) promoter (9). The usefulness of a C-terminal C_R in OhrRs in the protection against overoxidation of the N-terminal Cys has been demonstrated in the conversion of the 1-Cys *B. subtilis* OhrR into a 2-Cys regulator through introduction of a C-terminal cysteine either at position 120 (G120C) or 124 (Q124C), placing these C_R at 14.1 or 13.3 Å Sγ-Sγ distance, respectively, from the N-terminal Cys15 of the sister subunit (347). Both C_R variants effectively conferred the reversible oxidative regulation of promoter repression by OhrR. The recently characterized *Staphylococcus epidermidis* MarR family regulator, the aggregation and biofilm formation regulator (AbfR), senses organic peroxides *via* the 2-Cys interprotomer mechanism (220), yet AbfR does not undergo the α6-helix swapping that occurs for the *X. campestris* OhrR, but, instead, a slight “twisting” of the dimer interface is induced (219).

Despite the less dramatic structural changes in the AbfR dimerization region than that of the *X. campestris* OhrR, a much larger rigid-body transition is transmitted to the wHTH

FIG. 5. Oxidation-induced localized structural rearrangement in the transcription factor OhrR. Upon oxidation, intermolecular disulfide formation in *Xanthomonas campestris* OhrR is accompanied by a structural rearrangement at the homodimer interface, with interprotomer swapping of the $\alpha 6$ -helices. Respective protomers are *light gray* and *dark gray* and the structurally perturbed $\alpha 5$ - and $\alpha 6$ -helices are colored *black* for clarity. Structural representation prepared in PyMOL with crystal structures of the reduced (PDB ID, 2PEX) and oxidized (PDB ID, 2PFB) XcOhrR. OhrR, organic hydroperoxide resistance regulator; ROOH, peroxide.

domain of AbfR, with the interprotomer gap between the wHTH DNA-recognition helices ($\alpha 4$) widening by >11 Å in the oxidized form relative to the reduced DNA-bound form due to a marked rigid-body rotation and translation of the wHTH (Fig. 6). AbfR has been shown to sequentially form its cross-subunit disulfides, whereby a population of 1-disulfide crosslinked dimers progresses to a 2-disulfide form with increasing concentrations of organic peroxide (219). AbfR oxidation to a 1-disulfide or 2-disulfide state reduces its affinity toward the *abfR* operator site by 10-fold and 20-fold, respectively, indicative of two allosteric regulation stages that might be related to independent conformational changes occurring separately in each protomer (219).

Among the MarR family, alternate 2-Cys mechanisms of redox regulation have also been detected. The *P. aeruginosa* multidrug efflux regulator (MexR) does not possess the con-

FIG. 6. Regulation of DNA binding by intermolecular disulfide formation in *Staphylococcus epidermidis* AbfR. Native, reduced *S. epidermidis* AbfR (*black*) undergoes a conformational change to adopt a DNA-bound state (*dark gray*). Oxidation induces an intermolecular disulfide bond formation that locks AbfR in a conformation incompatible (*light gray*) with association to its cognate DNA. Structural superposition performed in PyMOL with crystal structures of the reduced (PDB ID, 4HBL), oxidized (PDB ID, 5HLI), and DNA-bound (PDB ID, 5HLG) forms. AbfR, aggregation and biofilm formation regulator.

served N-terminal cysteine of OhrR, but, instead, forms an interprotomer disulfide between a cysteine at the C-terminus of the $\alpha 1$ -helix (Cys30) and a cysteine on a loop following the $\alpha 3$ -helix (Cys62) of the opposing protomer (47). Cys62 is thought to form first a sulfenic acid, which leads to localized disruption of hydrogen bonds, allowing an 8 Å translation toward Cys30. The accompanying structural change is relatively localized, leading to a modest conformational shift in the wHTH regions, without changes in the distance between the DNA major-groove recognition helices. The overall conformations of reduced and oxidized forms of MexR remain largely the same and the structures align with a root-mean-square deviation of 1.6 Å (48). The decrease in DNA affinity is proposed to derive mainly from the steric hindrance introduced by the intermolecular disulfide (48). In addition to the intermolecular 2-Cys mechanisms outlined above, intramolecular disulfides have also been found to regulate some MarR family proteins, as in the case of an OhrR homolog of *Mycobacterium tuberculosis*, oxidation-sensing regulator, MosR.

In MosR, the conserved N-terminal oxidant-sensing cysteine (Cys12) forms an intramolecular disulfide with Cys10 upon oxidation, leading to transcriptional derepression (32). A markedly different mode of oxidation-induced intrasubunit linkage has been suggested for the bleach-sensing MarR family protein of *E. coli*, the N-ethylmaleimide regulator (NemR). NemR is specifically sensitive to both electrophiles and reactive chlorine species (RCS) and regulates its own expression and that of N-ethylmaleimide reductase (NemA) and glyoxalase (GloA) (195). Only one of the six cysteines of NemR, Cys106, is thought to be responsible for RCS sensing and an oxidation-induced sulfenamide bond has been suggested to form between a sulfenylated Cys106 and the Lys175 amino group (118). In addition to the intersubunit and intrasubunit disulfide/sulfenylamide mechanisms of oxidative regulation of the MarR-type transcription factors, an interdimer system of disulfide regulation has also been characterized for the *Burkholderia thailandensis* biofilm regulator (BifR) and the *E. coli* MarR. Both the *E. coli* MarR and BifR undergo interdimer disulfide linkages to form tetramers upon Cu(II) oxidation, but *via* distinct, nonconserved cysteines. Whereas

disulfide (and subsequent tetramer) formation by MarR inhibits DNA binding, oxidation actually increases the affinity of BifR toward DNA (123, 424).

IV. Midstream Redox Signaling by Moonlighting Proteins

Transcription factors sensitive to oxidants typically regulate the expression of antioxidant enzymes or proteins involved in coordinating cell fate (66, 217). Although such transcription factors exert a significant level of control over a cellular response to oxidative stress, they also limit the potential scale and complexity afforded by a signaling network through acting as simple on/off switches. By possessing an additional “midstream” level of constitutively expressed redox-sensing proteins beyond the transcriptional *downstream* effectors, cells can form a more complex and sensitive network of redox regulation, complimentary to the core antioxidant response. By virtue of being constitutively expressed, these midstream effectors bypass the need for protein synthesis before their regulatory effects are exerted. To fulfil the role of a redox-regulated *midstream* effector, the candidate protein should have a cellular function, distinct from any redox-related property, thereby justifying its expression under basal conditions. Furthermore, the candidate protein should be susceptible to reversible oxidation, and oxidation should alter its cellular role or subcellular localization in some way. In this study, we outline candidate proteins which we consider to fulfil this function, and therefore would constitute a level of *midstream* redox signaling, which shapes the cellular response to oxidative stress. The ability of such proteins to adopt a subfunction in response to redox stimuli compels us to refer to them as redox-regulated moonlighting proteins.

A. Human protein deglycase DJ-1

Human protein deglycase DJ-1 (protein deglycanase-1) is a member of the DJ-1/Hsp31/PfpI superfamily with a wide range of predicted cellular roles, including regulation of RNA–protein interactions (139), glyoxalase activity (200), chaperone function (334), cysteine protease activity (49), RNA binding (384), esterase activity (385), and transcriptional coactivation (410). It occurs preferentially as a noncovalent homodimer of 20-kDa protomers and contains an active site consisting of a catalytic triad of Glu18, Cys106, and His126, which confers glyoxalase and cysteine protease activity (49, 200). *In vivo*, DJ-1 is responsive to oxidative stress (242, 243) and Cys106 is considered to be the key redox-sensitive residue (37, 181, 317). A 1.2 Å crystal structure of DJ-1 revealed that in the absence of oxidizing agents, Cys106 is overoxidized to a sulfinic acid (37). DJ-1 displays redox-dependent esterase activity that apparently depends on the Cys106 oxidation state (385). Whereas the esterase function of DJ-1 is abolished by thiol alkylation, it is enhanced by exposure to H₂O₂ or a C106D sulfinylation mimic.

Human DJ-1 contains two additional cysteines, Cys46 and Cys53, and although both are oxidizable upon H₂O₂ treatment, they are significantly less sensitive to oxidants than Cys106 (181). Cys53 is located at the homodimerization interface of DJ-1 and is purported to form an intermolecular Cys53–Cys53 disulfide under oxidizing conditions to facilitate homodimer stabilization (89); there is also evidence suggesting that this disulfide is a reduction target for Trx (103). A further interaction between DJ-1 and another antioxidant enzyme, PrxII,

has been observed, also involving Cys53 of DJ-1 (89). DJ-1 has been demonstrated to form a mixed disulfide with PrxII *via* Cys53 under oxidizing conditions, possibly resolving itself into an intermolecular disulfide-bridged DJ-1 dimer (89).

DJ-1 oxidation has been found to promote its relocalization to mitochondria, an event associated with neuroprotective effects (37). Null mutation of DJ-1 causes an aberrant mitochondrial phenotype in human cells that can be rescued by infection of the human cells by viral particles expressing the wild-type protein or by administering antioxidants, such as *N*-acetyl-L-cysteine, but not by infection with viruses expressing the C106A mutant (150). Furthermore, knockout of DJ-1 in mice has been shown to lead to a twofold increase in mitochondrial H₂O₂ levels (6). The cytoprotective function of mitochondria-localized DJ-1 is suggested to derive from its interaction with the antiapoptotic protein B cell lymphoma-extra large (Bcl-X_L); NMR chemical shift experiments have since confirmed this interaction and it is proposed that binding of DJ-1 to Bcl-X_L inhibits the latter's ubiquitination and subsequent degradation (Fig. 7) (201, 302). Again, Cys106 is critical for this association, as a C106A mutation abolishes the interaction. DJ-1 is also recognized to bind subunits of the electron transport chain Complex I and to help in complex assembly, although the influence of DJ-1 oxidation on this interaction has yet to be explored (132, 135). Intriguingly, a specific Cys106 oxidation state is required for its protective influence in mitochondria. Whereas wild-type DJ-1 preferentially forms a stable sulfinic acid at Cys106, a mutation of the spatially proximal glutamate (E18D), designed to stabilize the sulfinylated Cys106 form, impairs the protective ability of DJ-1 against mitochondrial damage (24).

DJ-1 confers cytoprotectivity to oxidatively stressed cells by interfering with the c-Jun N-terminal kinase (JNK) and p38 mitogen-activated protein kinase (MAPK) apoptotic pathways, through a redox-dependent interaction with an upstream kinase of the JNK/p38 signaling pathway, the apoptosis signal-regulating kinase 1 (ASK1) (373). Under oxidizing conditions, ASK1 oligomerizes *via* intermolecular disulfide bridges to form HMW complexes of >3000 kDa, constituting an active ASK1 signalosome (253). ASK1 contains 23 cysteines, of which none is essential for the covalent complex linking. Only the multiple mutation of seven cysteines across the three ASK1 domains (the N-terminal, kinase, and C-terminal domains) abolished the ability of ASK1 to form disulfide-linked multimers (253). Upon oxidation, DJ-1 can engage in a mixed disulfide with ASK1 through its oxidant-sensitive Cys106 (391). Complex formation between ASK1 and DJ-1 inhibits the kinase function of ASK1, thereby suppressing the JNK/p38 apoptosis pathway under oxidizing conditions (Fig. 7). The peripheral Cys53 of DJ-1 might act as a C_R, driving the dissociation of the mixed disulfide, but this potential role has not been characterized *in vitro*. Crucially, whereas mildly oxidizing conditions promote the ASK1–DJ-1 complex formation, overoxidation of DJ-1 leads to its dissociation from ASK1, implying that this regulatory relationship is highly sensitive to cellular redox conditions (38).

DJ-1 also indirectly suppresses ASK1 by sequestering the death-associated protein Daxx and preventing it from ASK1 activation (166). In addition to its redox-regulated ASK1 modulation, DJ-1 also interacts with and inactivates MAPK kinase kinase 1 (MEKK1), another upstream JNK kinase that protects against UV-induced cell death. However,

FIG. 7. The multiple regulatory roles of DJ-1. Cytosolic DJ-1 is prone to oxidation at Cys106 that enhances its relocalization to the mitochondria and subsequent protection of Bcl-X_L from degradation and simultaneously facilitates mitochondrial complex I assembly. Cys106 oxidation is essential for the interaction with p53 and for the suppression of its transcriptional activity, whereas Cys53 oxidation is necessary for the chaperone function of DJ-1. DJ-1 also stabilizes Nrf2, preventing its redox-dependent degradation. Oxidized DJ-1 inhibits the kinase activity of ASK1, avoiding further downstream signaling through the JNK/p38 pathway. ASK1, apoptosis signal-regulating kinase 1; Bcl-X_L, B cell lymphoma-extra large; DJ-1, protein deglycanase-1; JNK, c-Jun N-terminal kinase; Keap1, Kelch-like ECH-associated protein 1; Nrf2, nuclear factor-erythroid 2 p45-related factor 2.

this interaction does not involve the redox-sensitive Cys106 of DJ-1 (10). DJ-1 intersects with another kinase pathway, the phosphatidylinositol 3' kinase pathway, through its role as a negative regulator of phosphatase and tensin homolog (PTEN) (177). DJ-1 has been suggested to mediate STAT1 dephosphorylation by promoting its interaction with the sarcoma (Src) homology 2 domain-containing protein tyrosine phosphatase 1 (SHP-1) (175).

DJ-1 influences the cellular antioxidant response *via* its association with the transcriptional regulator, nuclear factor-erythroid 2 p45-related factor 2 (Nrf2) (147). Nrf2 is the transcriptional activator of antioxidant response elements and, hence, mediates the expression of various antioxidant enzymes (151, 268, 388). Although the exact mechanism by which DJ-1 positively regulates Nrf2 activity is not well understood, it might prevent the binding of the inhibitor Kelch-like ECH-associated protein 1 (Keap1) to Nrf2 (Fig. 7) (56). Keap1 is a cysteine-rich (27 cysteines in the human isoform), 70-kDa protein that, under basal conditions, binds to Nrf2 and facilitates the association of the adapter component of the cullin 3 (Cul3)-based ubiquitin E3 ligase complex, leading to ubiquitination and proteasomal Nrf2 degradation (81). Astroglial overexpression of DJ-1 results in upregulation of the antioxidant enzymes, PrxII, Trx, and SOD1, although this could purely derive from the positive regulation of Nrf2 by DJ-1 (102).

DJ-1 has many other reported interaction targets, namely androgen receptor (262, 363), polypyrimidine tract-binding protein-associated splicing factor (PSF) (410), and p53 (86, 335). Among these targets, the interaction with p53 seems to be redox regulated, with oxidation of the Cys106 of DJ-1 essential for the interaction with the p53 DBD and subsequent repression of its transcriptional activity (170). DJ-1 also displays a chaperone activity that prevents α -synuclein aggregation, which is connected to Parkinsonism (115). Cys53 has been found necessary for this particular redox-dependent chaperone activity (334). Characterization of the chaperone properties of the prokaryotic DJ-1 homolog, YajL, revealed an extensive capacity for mixed disulfide formation with a wide range of cellular targets, extending to ribosomal proteins, metabolic enzymes, chaperones, transcription factors, and aminoacyl-transfer RNA (tRNA) ligases (192).

The bacterial $\Delta yajL$ mutant displays elevated levels of sulfenylated proteins, further supporting a redox-dependent protective role for YajL (109). Whereas the chaperone function of DJ-1 toward α -synuclein relies on Cys53, the covalent chaperone ability of YajL involves Cys106, consolidating the importance of this residue for the redox-dependent function of DJ-1/YajL. For human DJ-1, Cys106 is probably the predominant residue mediating mixed disulfide complex formation with various cellular partners and the amount of covalently captured complexes is enhanced by a C53A mutation, possibly through promotion of a DJ-1 monomeric state more prone to oxidative interactions (89). One protein identified in separate studies to form a mixed disulfide complex with both human DJ-1 and the bacterial homolog YajL, is another redox-regulated moonlighting protein, glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (89, 192). DJ-1 also interacts with GAPDH in an enzymatic manner, deglycating inactive glycated GAPDH (196, 304), and in human neuroblastoma cell lines was observed to form HMW complexes containing GAPDH and proteins related to RNA metabolism (285).

B. Glyceraldehyde 3-phosphate dehydrogenase

GAPDH is a key enzyme of the glycolytic pathway, catalyzing the reversible phosphorylation of glyceraldehyde 3-phosphate to glyceralate 1,3-bisphosphate with concomitant generation of reduced nicotinamide adenine dinucleotide (NADH). It is one of the most abundant intracellular proteins and, besides its role in glycolysis, performs many autonomous and often unrelated functions in multiple cellular compartments (342, 343, 378). A conserved cysteine at the active site (Cys152 in human GAPDH) acts as a reactive nucleophile and is essential for the catalytic function (31, 277). The Cys152 thiolate is highly sensitive to oxidative modifications, and, for a nonperoxidatic thiol, has a remarkably high rate constant for its reaction with H_2O_2 ($10\text{--}10^3\text{ M}^{-1}\cdot\text{s}^{-1}$) (215, 403, 418). Mathematical models predict that GAPDH is the first protein to become oxidized, once the dedicated thiol peroxidases are inactivated by overoxidation (above $10\text{ }\mu\text{M}$ H_2O_2) (403). Experimental observations are in agreement with these *in silico* predictions, as a $20\text{ }\mu\text{M}$ H_2O_2 bolus applied to Jurkat cells was demonstrated to be sufficient for GAPDH oxidation (12). H_2O_2 -mediated oxidation of GAPDH decreases its glycolytic activity (215, 277), although it is possible to uncouple the redox sensitivity of GAPDH from the oxidative inactivation of the enzymatic activity through mutations that reduce oxidant sensitivity (*e.g.*, C156S and Y314F) (277). Inactivation of GAPDH disrupts the glycolytic pathway, rerouting the carbohydrate flux to generate pentose phosphate pathway metabolites with concomitant generation of NADPH (295). As NADPH is the upstream electron donor of the thiol–disulfide regeneration systems, such as thioredoxin reductase (TR) or glutathione disulfide reductase (GR), increasing NADPH production under oxidative stress conditions is likely to be crucial for cell survival.

As a demonstration, in yeast cells expressing human H_2O_2 -insensitive mutants (C156S and Y314F) of GAPDH, the upregulation of NADPH production under oxidative stress is impaired and susceptibility to cell death is increased relative to wild type (277). In addition to inhibition of the catalytic activity of GAPDH, oxidation also induces aggregation of GAPDH, whereby oxidation of the surface-exposed Met46 of GAPDH is linked to protein misfolding, with subsequent aggregation through intermolecular disulfide formation (318). Neuronal GAPDH aggregation upon oxidative stress has been identified as a marker in neurological diseases (393). Although the Trx/TR system is the main reductant for oxidized GAPDH, tubulin, an abundant neuronal protein, can also engage in thiol–disulfide exchange to reduce GAPDH and, thus, specifically prevent oxidative aggregation/inactivation of neuronal GAPDH (189).

Oxidation or S-nitrosylation of the catalytic cysteine of GAPDH induces its association with Siah1, an E3 ubiquitin ligase protein involved in proteasome-dependent protein degradation (130). Binding of GAPDH to Siah1 promotes localization of GAPDH to the nucleus, where it influences the apoptosis pathway and the enzymatic DNA repair systems. Nuclear GAPDH promotes acetylation and phosphorylation of p53 in the absence of the poly A-binding protein, stimulating translocation of p53 to mitochondria, where it participates in apoptosis initiation (371). GAPDH has also been implicated in mediating the upregulation of p53 expression (420) and, in turn, p53 has been shown to upregulate the mRNA transcription of GAPDH and Siah1 (50, 93). ASK1 augments the

interaction between GAPDH and Siah1 and phosphorylates threonines of Siah1 to activate its nuclear signaling pathway (379), whereas H_2O_2 treatment has been shown to increase the association of ASK1 to GAPDH-Siah1. Nuclear-translocated GAPDH interacts with the base excision DNA repair enzyme, apurinic/apyrimidinic endonuclease 1 (APE1), and reduces the oxidized APE1 in a manner dependent on the redox-sensitive GAPDH cysteines, Cys152 and Cys156 (10). APE1 itself has redox-dependent functions, capable of reactivating the extracellular signal-regulated kinase (ERK) activity *via* its oxidant-sensitive cysteine Cys65 (397) and stimulating p53 DNA binding in a redox-dependent manner (60). Oxidation of human GAPDH induces an interaction with the RNA-binding proteins, PSF and 54-kDa nuclear RNA-binding protein (p54nrb), to form a complex that enhances the topoisomerase I activity (146), and this interaction is dependent on the presence of the catalytic Cys152 site of GAPDH. H_2O_2 treatment of human cell lines has been shown to result in the formation of a mixed disulfide between GAPDH and either PrxI or PrxII (353), with Prx presumably mediating the reversible oxidation of GAPDH thiols, and a proteome-wide study of H_2O_2 -sensitive protein thiols of *Arabidopsis* chloroplasts confirmed that GAPDH undergoes reversible thiol oxidation (250). Among the 16 other oxidant-sensitive stromal proteins identified were 2-Cys Prx and the peptidyl-prolyl *cis-trans* isomerase (PPI) enzyme, cyclophilin 20-3 (Cyp20-3) (250).

C. Peptidyl-prolyl *cis-trans* isomerase

The *Arabidopsis* PPI, Cyp20-3, is a multifaceted protein, and in addition to its PPI activity, plays a putative role in facilitating protein unfolding and degradation under stress (308). Upon oxidation, Cyp20-3 undergoes a large conformational change to form two sets of intramolecular disulfides, Cys54-Cys171 and Cys129-Cys176 (191). Cyp20-3 oxidation inhibits its PPI activity (191, 248) and has been suggested to induce a regulatory function in cysteine synthesis, as under oxidative stress conditions Cyp20-3 interacts with the cysteine synthase complex to activate cysteine synthesis (76). Despite its low redox potential ($E_m = -319$ mV), Cyp20-3 can be maintained in a reduced state by Trx-m ($E_m = -357$ to 368 mV) (57) and might also be a reductant of 2-Cys Prx A and Prx B (E_m of -307 and -322 mV, respectively), although it is unable to enhance the peroxidase activity of 2-Cys Prxs (191, 251).

Mammalian PPIs with redox-responsive functions include the cytosolic cyclophilin A (CypA), cyclophilin B (CypB), which is localized to the endoplasmic reticulum, mitochondrial CypD, and the phosphorylation-dependent parvulin family PPI, Pin1 (peptidyl-prolyl *cis-trans* isomerase NIMA-interacting 1). Similarly to Cyp20-3, human CypA can also act as an electron donor for Prx II and Prx VI and has been demonstrated to enhance their peroxidase activity *in vitro* (203). CypA negatively regulates the JNK/p38 signaling pathway in response to oxidative stress by interacting with ASK1 and suppressing its phosphorylation (174). CypA has been observed to be secreted by vascular smooth muscle cells under conditions of oxidative stress and activates ERK1/2 in a manner dependent on its peptidyl-prolyl isomerase activity (160).

Mammalian CypD is a regulator of the mitochondrial permeability transition pore (mPTP) (329), the activation of which leads to loss of the mitochondrial membrane potential and is linked to necrotic and apoptotic cell death. CypD is

activated by p53 and, in turn, induces p53 aggregation through its isomerase activity (193). CypD activation occurs in response to oxidative stress (324) and its Cys203 has been identified as the redox-sensing cysteine responsible for oxidative regulation and forms an intramolecular disulfide upon H_2O_2 exposure *in vivo* (214, 259). CypD interacts with Trx2 and PrxIII in the mitochondrial matrix (94) and accumulation of oxidized Trx2, either by small interfering RNA (siRNA) knockdown of thioredoxin reductase 2 (TR2), or inhibition of TR2 by auranofin, induces the concomitant oxidation of CypD and PrxIII, implying a level of redox interplay between these enzymes. CypD increases the respiratory activity of Complex III in HEK293 cells by promoting supercomplex formation (85). Overexpression of Cyp22, a CypD homolog of *Trypanosoma cruzi*, led to increased sensitivity to mitochondrial destabilization through loss of membrane potential in response to oxidative stress (34). Under conditions of oxidative stress CypD binds to mitochondrially localized STAT3 in a manner dependent on the N-terminus region of STAT3, and it is proposed that this interaction relates to reduced mitochondrial ROS production (234).

A relationship between STAT3 and CypB has also been demonstrated in cancer cells, with STAT3 repressing transcription of CypB inhibitors, and CypB in turn promoting the activation of STAT3 (210). The parvulin family PPI, Pin1, specifically targets prolines adjacent to phosphorylated Ser/Thr and makes a junction between PPIases and kinase pathways, whereas oxidation of its Cys113 to sulfenic/sulfinic acid inhibits its catalytic activity (46, 149, 411). Pin1 also intersects with cellular stress responses by targeting JNK (272), p53 (120), and p66 Src homologous-collagen homolog (p66Shc) (88). Pin1-mediated translocation of p66Shc to mitochondria leads to increased levels of mitochondrial ROS and induction of apoptosis (113). The adaptor protein, p66Shc, itself contains multiple oxidant-sensitive cysteines, and can undergo a redox-dependent transition between homodimeric and a homotetrameric disulfide-linked oligomer.

D. Transglutaminase 2

Transglutaminases (TGs) are a family of enzymes that catalyze Ca^{2+} -dependent transamidation and (under acidic conditions) deamidation reactions on a wide range of intracellular and extracellular targets (for a comprehensive list of substrates, see the TRANSDAB database, http://genomics.dote.hu/wiki/index.php/Main_Page). Humans possess nine different TGs, of which transglutaminase 2 (TG2) is the most widely distributed and is associated with a plethora of disease phenotypes related to oxidative stress. Links between TGs and cataracts (202), celiac disease (224), cancer (154), inflammation (225), neurodegeneration (11), and diabetes (20), all suggest a ROS-dependent role of TGs in disease response. The role of TG2 in cellular responses to ROS can clearly be linked to its ability to modulate stress-responsive transcription factors, such as NF- κ B and p53 (154, 240), and more recently the redox regulation of TG2 itself is increasingly recognized (28, 349). For their transamidase activity, TGs employ a catalytic triad consisting of a histidine (His335), aspartate (Asp358), and catalytic cysteine (Cys277), and the presence of a GTP-binding site in its β -barrel 1 domain enables TG2 to also act as a G protein (255).

Oxidation of TG2 has been shown to induce intramolecular disulfide formation, either at Cys230-Cys370, or at Cys370-Cys371 (349). Based on site-directed mutagenesis of Cys230,

a disulfide relay might occur, with first the formation of the Cys230-Cys370 disulfide and then with Cys371 presumably acting as a C_R for the generation of the resulting Cys370-Cys371 vicinal disulfide. Oxidation of TG2 inhibits its transamidase activity, and mutation of Cys230 has been shown to render TG2 insensitive to inactivation by oxidation, although this was attributed to an impaired ability to form the Cys370-Cys371 disulfide (349).

TG2 can adopt either a closed ATP/GTP-bound conformation, in which state its transamidation activity is lost, or an open transamidase-active conformation stabilized by Ca²⁺ binding (284). Oxidation appears to influence the conformational preference of TG2, as the Cys230-Cys370 disulfide form preferentially adopts the open configuration even in the presence of GTP (Fig. 8) (349). However, the Cys230-Cys370 disulfide itself is not incompatible with the closed configuration of TG2, as proved by its presence in crystal structures of ATP and GTP-bound closed conformation TG2 (127, 156). Just as GTP inhibits the transamidase activity of TG2 by promoting the closed conformation, Ca²⁺ can decrease the GTPase activity of TG2 (182). TG2 has a binding *K_D* of 1.6 μ M for GTP, compared with a *K_D* of 90 μ M toward Ca²⁺ (15). Considering that typical intracellular Ca²⁺ concentrations are in the range of 100 nM and that free GTP is estimated to be $\sim$ 100 μ M, the majority of intracellular TG2 is expected to be in a closed, GTP/GDP-bound conformation without transamidase activity (183). In this transamidase-inactive conformation, TG2 has been demonstrated to function as a kinase (239–241) and as a protein disulfide isomerase (PDI) (131). As the vicinal Cys370-Cys371 disulfide has a redox potential similar to that of bovine PDI [−184 $\pm$ 4 mV (159) and −190 $\pm$ 10 mV, respectively (226)], this implicates the vicinal disulfide as a potential catalytic site for disulfide isomerization. In terms of an *in vivo* relevance of the PDI action, a role for TG2 in the correction of defective disulfide bonds in the respiratory chain complexes has been proposed (232).

TG2 can be secreted to the extracellular environment, where it serves as a target substrate for extracellular Trx (7, 159). Trx has been shown to be highly selective in its extracellular TG2 targeting in fibroblasts and the small intestine, although its

ability to recognize TG2 in other organ environments has been questioned (7). Surprisingly, chloride intracellular channel protein 3 (CLIC3) has also been proposed to be an extracellular reductant of TG2, utilizing the catalytic cysteine of the CXXC motif of its Trx fold to perform a GSH-dependent reduction of oxidized TG2 (138). CLIC proteins adopt either a soluble globular form, structurally homologous to the glutathione *S*-transferase (GST) family or an oligomeric integral membrane-associated state for ion conductance. Although CLIC3 has not previously been associated with redox-dependent functions, the structure and/or function of CLIC1, CLIC2, and CLIC4 has been previously demonstrated to be altered depending on the oxidation state of the cysteines of its Trx domain (216). More recently, an extracellular oxidizing protein partner for TG2 has been identified. ERp57 (endoplasmic reticulum-resident protein 57 [also known as protein disulfide-isomerase A3, or glucose-regulated protein 58-kDa]) was observed to colocalize with TG2 in cultured human umbilical vein endothelial cells, and was capable of oxidatively inactivating TG2 at a second-order rate constant >700-fold higher than inactivation by oxidants such as H₂O₂ or glutathione disulfide (GSSG) (417). The dual axis of Trx and ERp57 in extracellular redox regulation of TG2 may serve to modulate extracellular transamidation, where Ca²⁺ and nucleotide concentrations favor a transamidase-active form of TG2, and so oxidative inhibition of its activity may have a more significant role.

V. Upstream Redox-Sensing: Prxs as Specific Sensors and Mediators of the Oxidative Stress Response

The high cellular abundance of Prxs and the superior catalytic reactivity of their peroxidatic thiols toward peroxides make them the frontline intracellular oxidant sensor, thereby placing Prx in the *upstream* section of oxidative signaling pathways. The proposed role of Prx as a sensor and transducer of H₂O₂ signaling has been consistently strengthened (125, 126, 136, 246, 277, 345, 353, 354), and in this study we consolidate the view of Prx as an instigator of oxidative signaling, and provide a further overview of the

FIG. 8. TG2 adoption of two functionally distinct conformations. To be catalytically active as a transamidase, TG2 must be in an open conformation (light gray). The closed conformation of TG2 (black) is inactive as a transamidase, but has GTPase and kinase activities. Binding of Ca²⁺ promotes the open conformation, whereas binding of GTP/ATP induces a large-scale conformational change (dashed arrow) to the closed conformation. The transamidase catalytic site and vicinal cysteines Cys370 and Cys371 are indicated. Figure prepared in PyMOL with the crystal structures of GTP-bound (black; PDB ID 4PYG) and inhibitor-bound (gray; PDB ID 3S3J) human TG2. TG2, transglutaminase 2.

potential cellular roles of Prx and its intersection with alternate intracellular signaling pathways.

A. Relationship between redox state and Prx conformation

Prxs catalyze the reduction of peroxide to water/alcohol through a nucleophilic attack by a C_P thiolate that in the process becomes oxidized to a sulfenic acid. In “typical 2-Cys” Prxs, the C_P is spatially conserved in the first turn of α -helix 2, ~14 Å away from the C_R, which is located in the C-terminal region of the neighboring subunit of the Prx homodimer. For a disulfide to form between the C_R and a sulfenylated C_P, localized unfolding is required of both the α 2-helix active site region and the C_R-containing C-terminal tail. This process is defined as a transition from a “fully folded” (FF) to “locally unfolded” (LU) state. Recent consensus in the field of Prxs is that the FF-LU transition is in dynamic conformational equilibrium before C_P oxidation (167, 168, 260, 279). The structural switch from a FF to LU conformation in Prx is an essential protective mechanism, and stabilization of the FF conformation is widely accepted to promote C_P overoxidation to sulfinic or sulfonic acid, inactivating Prx until regeneration of the sulfinylated form by sulfiredoxin (Srx). Prxs that are prone to or resilient to overoxidation are designated “sensitive” and “robust,” respectively. In phylogenetic terms, eukaryotic Prxs are largely sensitive and prokaryotic Prxs robust to overoxidation (408, 415). Eukaryotic Prxs typically contain a conserved YF sequence motif in the C-terminal tail that interacts with the first turn of the α 2-helix to support its packing against the active site, while obstructing the local unfolding of the active-site region (168, 322, 395).

With the exception of the monomeric PrxQ/bacterioferritin comigratory protein (BCP) (158, 212), the dimerization of Prxs is a highly conserved feature, with two distinct dimerization interfaces, designated A-type and B-type. Typical 2-Cys Prxs form homodimers through the B-type interface with protomers interacting in a head-to-tail fashion to bring adjacent β -sheets together in an antiparallel alignment (54), whereas the atypical 2-Cys Prxs dimerize *via* the A-type interface. Through the A-type interface, typical 2-Cys Prxs form higher-order oligomeric structures that occur most commonly as a toroidal decamer of five homodimers and in a few rare cases as dodecamers (121, 327). In the FF conformation, the C_P-containing active-site loop buttresses the decameric interface, thereby stabilizing the oligomeric toroid. C_P sulfenylation from the catalytic reduction of H₂O₂ leads to disulfide formation between the C_P and C_R, locking the active site in an LU conformation and destabilizing the decamer-building interface (407).

The strong relationship between the structural environment of the C_P and the oligomeric state of Prxs has been explored by site-directed mutagenesis, with mutation of the conserved active-site threonine (Thr44 in *S. cerevisiae* thiol-specific antioxidant 1 [Tsa1]) to valine or serine, modulating the oligomeric state toward dimer or decamer, respectively (362). The Thr44-flanking phenylalanines (Phe43 and Phe45) form hydrophobic interactions at the A-type interface and the oxidation-induced switch to the LU conformation is thought to disrupt these stabilizing interactions, leading to decamer dissociation (245, 407). As an exception of such redox-dependent oligomerization in 2-Cys Prxs, PrxIV keeps a stable decameric

arrangement in the absence of a reducing agent (40). The oligomeric structure of PrxIV might be maintained by its distinct N-terminal extension, a feature lacking in other Prx subtypes. In the 2-Cys Prx of *Salmonella typhimurium* (AhpC), a threonine mutation at the decameric interface either stabilizes (T77V) or destabilizes (T77I) the oligomeric structure without direct interference with the active site (275). Destabilization of the AhpC decamer greatly reduces the catalytic efficiency of the enzyme for H₂O₂, but also makes the enzyme more resistant to overoxidation at the C_P. The decreased sensitivity of the decamer-destabilizing mutants to peroxide-mediated inactivation is attributed to an increased flexibility of the active-site region, evidenced by high temperature factors in the crystal structure of AhpC_{T77I} (275). This phenomenon could imply a dynamic mechanism linking redox-dependent oligomerization to the catalytic Prx function, in which the decameric state of the enzyme reacts with peroxide and then dissociates to dimers to allow the rapid formation of a C_P-C_R disulfide bond, thereby preventing enzymatic inactivation by the C_P overoxidation.

Two possible mechanisms for the role of Prxs in oxidative signaling have been proposed; a direct transfer of oxidative equivalents from the oxidized Prx to a reduced target protein *via* a redox relay involving a temporary interprotein mixed disulfide (Fig. 9) (95), or the “floodgate” hypothesis, whereby Prx inactivation by hyperoxidation to a sulfinylated form leads to a localized build-up of H₂O₂ (408).

B. Thiol peroxidase redox relays

The concept of a Prx-based redox relay first gained credence after a ground-breaking study had revealed that the yeast thiol peroxidase Orp1 (previously termed GPx3) transferred oxidative equivalents to the transcription factor Yap1 (70, 354). The critical significance of this redox relay is that Yap1 is not directly responsive to H₂O₂, and oxidation can occur only with peroxidase involvement (228). Upon oxidation by H₂O₂, a sulfenic acid is formed on the C_P of Orp1, Cys36, which can be resolved into a mixed disulfide with the Cys598 of the C-terminal cysteine-rich domain of Yap1. In turn, Cys303 of the N-terminal cysteine-rich domain of Yap1 reacts with Cys598 to produce an intramolecular, interdomain disulfide and to release Orp1. Unless oxidized, Yap1 is rapidly reduced and further sequential interdomain disulfides can occur, resulting in a final triple disulfide oxidation state, involving Cys310-Cys629 and Cys315-Cys620 in addition to the initial Cys303-Cys598. The single Cys303-Cys598 interdomain disulfide is sufficient to mask the C-terminal nuclear export signal of Yap1, leading to nuclear localization and transcriptional activation of oxidative stress response genes (69, 70), but the peak of transcriptional activity is reached only upon generation of the triple disulfide form that has an enhanced resistance to reduction (as evidenced by a decrease in redox potential relative to the single disulfide form) and, thus, can probably confer a more sustained activation in the presence of reductants (266).

Functionally equivalent redox relays have been described for Yap1 homologs, the *Candida albicans* Cap1 that is also oxidized by Orp1, and the *Schizosaccharomyces pombe* (fission yeast) Pap1 that is oxidized by Tpx1, a homolog of *S. cerevisiae* Tsa1 (35). Whereas the redox relay between Orp1 and Yap1 proceeds *via* an Orp1-SOH intermediate, the Tpx1 relay is proposed to proceed *via* a thiol-disulfide exchange, involving the C_P-C_R disulfide of the Tpx1 oxidized form (17,

FIG. 9. Prxs in action: removal of peroxides and transfer of oxidative equivalents. Prxs belong to the first line of cellular defense antioxidants. Primarily they reduce H_2O_2 to H_2O , thereby maintaining the reducing environment of the cell, although, they recently have been shown to transfer oxidative equivalents to target proteins *via* transient mixed disulfide complexes. Thus far, the best described system is the transfer mechanism in which PrxII acts on the transcription factor STAT3 (345). At high H_2O_2 concentrations, Prx is overoxidized and forms a sulfenic acid ($-\text{SO}_2\text{H}$) on its peroxidatic cysteine. Srx is the ATP-dependent return ticket for overoxidized Prx. This overoxidation mechanism has been described as part of the Prx regulation mechanism, also known as the floodgate model (408). STAT3, signal transducer and activator of transcription 3.

35). As proteomic studies have revealed an abundance of proteins with surface-exposed thiolates, along with millimolar concentrations of LMW antioxidants, such as GSH, and submicromolar concentrations of Trx, the question arises of why Orp1 specifically oxidizes Yap1, especially considering the presence of a C_R in Orp1 that should rapidly react to form a C_P - C_R disulfide.

This issue has been addressed by the identification of a Yap1-specific scaffold protein, the Yap1-binding protein 1 (Ybp1) (122, 387). In the absence of Ybp1, Orp1 forms a mixed disulfide with Yap1 at a predicted rate constant of 10^{-3} s^{-1} , too slow for effective Yap1 oxidation in the cellular environment. Ybp1 recruits Orp1 and Yap1 into a ternary complex, effectively increasing their respective local concentrations and the oxidative relay rate constant to 1 s^{-1} (17). The binding of Ybp1 to Orp1 also prevents the C_R of Orp1 (Cys82) from reacting with the sulfenylated C_P before a mixed disulfide can be formed with the Cys598 of Yap1 (17). In yeast strains that produce a truncated Ybp1 gene product, Tsa1 takes over the function of Orp1 (265, 361). Tsa1 ΔC_R is able to form a mixed disulfide with Yap1 *via* its C_P (Cys48), but only the wild-type Tsa1, not the ΔC_R variant, can activate Yap1 in response to oxidative stress. Thus, either a thiol-disulfide exchange mechanism is the preferred mode of oxidant transduction between Tsa1 and Yap1, or a Tsa1 homodimer with only one intermolecular disulfide linkage, leaving the C_P of one subunit free for sulfenic acid condensation, might be the Yap1-activating form (361).

The highly efficient redox relay between Orp1 and Yap1 inspired the development of a fluorescent probe based on roGFP2. By linking Orp1 and roGFP2 together as a fusion

protein construct, the proximity dependence of the relay facilitated by Ybp1 can be enabled artificially. By comparing the redox relay characteristics of a wild-type and an Orp1 C_R mutant (C82S) fusion construct, the roGFP-Orp1 $_{\text{wt}}$ was demonstrated to transduce oxidation by a thiol-disulfide exchange mechanism, whereas the roGFP-Orp1 ΔC_R relays oxidative equivalents directly *via* a sulfenic acid intermediate (126). Besides the redox relay systems described above, the occurrence of oxidant transduction between the yeast Ahp1 and the Yap1 homolog Cad1 has been suggested and an oxidant relay between PrxII and the transcription factor STAT3 has been described. Direct STAT3 oxidation by H_2O_2 is very slow, but in the presence of PrxII, the oxidation rate is significantly enhanced (345). PrxII oxidizes STAT3 *via* a mixed disulfide, although it is not known whether *via* thiol-disulfide exchange, or direct sulfenic acid condensation. STAT3 oxidation leads to the formation of intermolecular disulfide-bonded STAT3 dimers and tetramers, and the involvement of different cysteines of STAT3 in dimer and tetramer formation implies independent PrxII oxidation pathways, resulting in different oligomerization states (345).

Another mammalian transcription factor found to be regulated by Prx-mediated oxidation is forkhead box protein O (FOXO) 3, which has been demonstrated to form mixed disulfides with PrxI *in vivo*, and formation of a PrxI-FOXO3 mixed disulfide appears to induce nuclear localization of FOXO3 (141, 287). PrxI has also been reported to facilitate oxidation of the APE1, and although the mechanism of oxidation is not fully characterized, it appears to involve a PrxI-APE1 mixed disulfide intermediate (256). The resulting oxidized form of APE1

is also unknown, although studies of its oxidation states show that it can form a mixture of disulfide-bonded multimeric species through Cys65, Cys93, Cys99, and Cys138, and that it is also capable of forming a mixed disulfide complex through Cys99 with Trx (227).

Further to its role as a moonlighting redox sensor, DJ-1 is suggested to engage in a thiol–disulfide exchange mechanism with PrxII (72), whereby oxidized PrxII with a C_P–C_R disulfide forms a mixed disulfide with Cys53 of DJ-1, which resolves into an intermolecular disulfide involving the Cys53 of another monomer to generate a covalent homodimer (89). All-in-all, peroxidase-based redox relays have been established to have both a functional relevance and technological applications, but until recently, the number of truly validated redox relays has been exceptionally low, especially relative to the vast number of oxidizable protein thiols identified in proteomic studies. However, the recent findings that the simultaneous depletion of PrxI and PrxII massively decreases the levels of H₂O₂-induced protein thiol oxidation (353) provide compelling evidence in favor of oxidative signaling *via* redox relays.

C. Nonperoxidatic roles for Prx

The floodgate hypothesis was proposed based on the observation that some eukaryotic Prxs appear to have evolved a specific sensitivity to hyperoxidation (408). However, there is presently a lack of experimental support for this mechanism and the proportion of hyperoxidized 2-Cys Prx in mammals is usually quite low, estimated to be <1.6% in mice erythrocytes (52). Accordingly, specific cellular situations might occur in which overoxidized 2-Cys Prx can sufficiently accumulate to facilitate floodgate-based signaling. For example, in the adrenal cortex, 10–20% of PrxIII is found to be hyperoxidized (172). Increased ROS leakage caused by overstimulation of steroid hydroxylation by CYP11B1, a cytochrome P450 family protein, in mitochondria of adrenal gland cells has resulted in PrxIII hyperoxidation and a corresponding increase in p38 MAPK phosphorylation (172). In this study, hyperoxidation was proposed to allow the build-up of mitochondrial H₂O₂ and subsequent overflow to the cytosol, whereby oxidative ASK1 activation, an upstream kinase of p38, leads to p38 phosphorylation (172). However, another study has highlighted a steep concentration gradient of H₂O₂ across the mitochondrial membrane, hence, changes in mitochondrial H₂O₂ levels are expected to have relatively little impact on the cytosolic concentration of H₂O₂ (246). A redox relay mechanism for ASK1 regulation has also been suggested based on the observation that PrxI and PrxIII form a mixed disulfide with ASK1 in mammalian cell lines, thereby inferring a direct ASK1 oxidation mechanism by Prx (157).

An alternative to a hyperoxidation-driven floodgate is the transient inactivation of Prx by reversible phosphorylation that would influence H₂O₂ gradients independently of H₂O₂ itself. The peroxidase activity of PrxI can be inactivated by Tyr194 phosphorylation (which is positioned ~9 Å from the C_P) by Src kinases upon cell stimulation by growth factors (406), Thr90 and Thr183 phosphorylation by the mammalian sterile20-like kinase 1 (MST1) (298), and by Thr90 phosphorylation by the cyclin-dependent kinase 1 (CDK1) (45). The peroxidase activity of PrxII decreases upon Thr89 phosphorylation by the CDK5/p35 complex (290). Conversely, phosphorylation of Ser32 of PrxI by the T-Lymphokine–

Activated Killer (LAK) cell–originated kinase (TOPK) enhances the peroxidase activity (425), thereby demonstrating that location of phosphorylation is critical for its influence on Prx function. Prx susceptibility to phosphorylative inactivation or hyperoxidation can be independent of the isoform, with PrxI relatively insensitive to hyperoxidation, but readily inactivated by phosphorylation, whereas PrxII is typically not susceptible to inactivation by phosphorylation, but more prone to hyperoxidation (406). A number of kinases and phosphatases are becoming regarded as being redox regulated themselves, thereby adding an additional level to phosphorylative Prx regulation. Oxidation of a conserved cysteine of the Src kinases promotes their activation, increasing the likelihood of PrxI phosphorylation and inactivation.

Furthermore, several members of the protein tyrosine phosphatase (PTP) family, which could potentially dephosphorylate (and therefore activate) PrxI, are susceptible to inactivation through the reversible oxidation of cysteine residues (235, 267), culminating in a positive feedback loop that favors PrxI inactivation by phosphorylation and prevention of dephosphorylation. As PTPs have a relatively low reactivity toward endogenous H₂O₂ levels ($k = 9\text{--}18\text{ M}^{-1}\cdot\text{s}^{-1}$) (72, 90), it is tempting to consider the possibility of direct oxidation of PTP by Prx as an additional level of redox-regulatory control. However, in the case of PTP1B and PrxII, a redox relay was not found to occur upon H₂O₂ exposure (62). Although possible redox relay systems between Prx/peroxidases and kinases have been proposed for ASK1 (157) and the *S. pombe* p38/JNK stress-activated protein kinase (SAPK), Sty1 (386), many other regulatory, redox-related interactions with Prx and kinases/phosphatases have been characterized without involvement of a mixed disulfide intermediate. PrxI engages in regulatory interactions with the phosphatases, PTEN, MAPK phosphatase 1 (MKP1), and MKP5, in a manner which is redox dependent, yet does not involve mixed disulfide formation (39, 382). PrxI binds to PTEN and MKP1 with similar affinity ($K_d = 247.3 \pm 33.6$ and $261.0 \pm 21.2\text{ nM}$, respectively) and dissociates upon oxidation, but binds MKP1 with an increased affinity ($2.4 \pm 0.2\text{ nM}$) to form a complex that does not dissociate upon oxidation (382). PTEN, MKP1, and MKP5 reside in a class of PTPs that possess a nucleophilic catalytic cysteine, sensitive to oxidation. PTEN oxidation leads to the formation of an intramolecular disulfide, involving its active-site cysteine, whereas oxidation of MKP1 and MKP5 induces the formation of disulfide-linked oligomers; in both cases, oxidation results in loss of phosphatase activity. PrxI preserves the lipid phosphatase activity of PTEN by protecting it against oxidative inactivation, and also protects MKP5 against oxidation-induced oligomerization and subsequent inactivation.

Conversely, PrxI-associated MKP1 is more susceptible to oxidative inactivation upon exposure to H₂O₂. As MKP1 dephosphorylates both p38 α MAPK and JNK, whereas MKP5 dephosphorylates only p38 α MAPK, the involvement of PrxI indirectly activates JNK signaling while inactivating p38 α (190). A direct regulation of p38 MAPK by PrxI has been suggested based on the coimmunoprecipitation of PrxI with phosphorylated p38 MAPK, but the nature of this interaction has still to be characterized *in vitro*.

The HMW oligomeric form of PrxI has been proposed to promote MST1 activity by association with the MST1 C-terminal kinase-inhibitory domain (247). Significantly, phosphorylation of PrxI at Thr90 by CDK1 or MST1 induces the

formation of HMW oligomers of PrxI (Fig. 10), hinting to a positive feedback loop promoting PrxI phosphorylation by MST1 (155). Phosphorylation is not, however, a general mechanism for the stimulation of Prx oligomerization, because phosphorylation at Tyr194 of PrxI by Src kinases does not influence the oligomeric state of PrxI (406). The conversion of PrxI to a HMW oligomer confers a chaperone function, but reduces the peroxidase activity (179). Prx hyperoxidation is known to trigger the formation of HMW oligomers; therefore, under the conditions to which the floodgate signaling hypothesis applies, a corresponding increase in Prx chaperone action should also be considered (263). The conversion to HMW oligomers involves stacking of decamers *via* “R-interface” interactions to create filamentous structures. Stacking of the decamers into HMW oligomers prevents the proper folding of the C-terminal tails, leaving them as disordered protrusions (314). As the unstructured, hydrophobic regions of other chaperone proteins are known to be involved in substrate binding, these unfolded C-terminal regions of HMW oligomeric Prxs have been postulated to play a role in conferring holdase activity (314).

Currently, the connection between the oligomeric state and regulatory interactions of Prx is unclear and seems to vary depending on the protein substrate. In the case of mitochondrial Prx of the protozoan parasite *Leishmania infantum*, overoxidation and oligomerization to HMW species is not necessary for Prx to act as a holdase on mitochondrial proteins under heat stress (369). PrxI is proposed to bind PTEN as a monomer, as preservation of PTEN phosphatase activity reaches its maximal under equimolar ratios of PrxI and PTEN (39). A relationship between the oligomeric state of PrxI and its regulation of MKP1/MKP5 has not been fully characterized. PrxI binds to Toll-like receptor 4 (TLR4) in the pres-

ence of myeloid differentiation protein 2 (MD2) and cluster of differentiation 14 (CD14) to stimulate cytokine release, in a manner independent of peroxidase activity, but dependent on PrxI oligomerization (305). Nevertheless, whether the decameric form or HMW chaperone form of PrxI is necessary for its interaction with TLR4 is still unclear. Chloroplast 2-Cys Prx can enhance the activity of chloroplast fructose-1,6-bisphosphatase (CFBPase), although not through function as a chaperone, as it is unable to prevent CFBPase denaturation, but *via* protection against CFBPase autooxidation and alteration of its affinity to its substrates (41).

In some cases, a specific physiological function of Prx can require the specifically directed regulation of the balance between hyperoxidized HMW oligomers and the reduced decameric form. Recently, the potential role of the yeast Prx, Tsa1 has been demonstrated in preventing the accumulation of H₂O₂-induced aggregates in stressed yeast cells (129). Decameric Tsa1 was found to physically associate with the Hsp70-type protein, Ssa1, in a redox-independent manner and, upon hyperoxidation, localized along with Ssa1 or Ssa2 to inclusion aggregates, leading to a decrease in aggregate formation by the promotion of proteasomal degradation or refolding of substrate proteins (Fig. 11). Tsa1 was also able to trigger Hsp104- and (Hsp40-type) Sis1-mediated disaggregation in a Srx-dependent manner, implying a necessary conversion of the hyperoxidized form of Tsa1 to its reduced form, possibly driving Tsa1 dissociation from aggregates to allow disaggregation (129), thus providing a eukaryotic example of redox-dependent regulation of the proteostasis machinery through the control of the oxidative state of a Prx.

Considering the significant influence of the oligomeric state of Prx on its cellular function, it is important to note that

FIG. 10. Influence by unrelated enzymes on the oligomeric Prx state. Prx exists in an equilibrium between dimeric and (depending on the isoform) decameric oligomeric states. Association of the oxidized, tetrameric p66Shc shifts the multimeric equilibrium toward a dimeric state, also favored by the oxidized Prx form. Dephosphorylated CDK1 promotes the HMW chaperone Prx state, also preferred by the hyperoxidized Prx. Hyperoxidation or oxidation to the HMW or dimeric forms of Prx can be reversed through reduction by Srx or Trx, respectively. Trx is shown indirectly influencing the oligomeric Prx state by reducing either the oxidized p66Shc or (thereby activating) Cdc25, a CDK1 phosphatase. Depictions of structural models were prepared with the crystal structure of the reduced human PrxII (PDB ID, 1QMV), oxidized human PrxIII (PDB ID, 5K2I), and C_R mutant of human Trx (3E3E), which mimics the reduced form. The HMW chaperone form of Prx is presented [adapted from Yewdall *et al.* (416)]. Cdc25, cell division cycle 25; CDK, cyclin-dependent kinase; C_R, resolving cysteine; HMW, high-molecular-weight; p66Shc, p66 Src homologous-collagen homolog; Trx, thioredoxin.

FIG. 11. Tsa1 as a cochaperone to the Hsp70/Hsp104 system in yeast. The Hsp70 homolog, Ssa1/2, associates with the decameric Tsa1. Under oxidative stress conditions, the hyperoxidized HMW form of Tsa1 ferries Ssa1/2 to protein inclusions, thereby inducing protein refolding or degradation. Tsa1 can then recruit Sis1 and Hsp104 to protein aggregates and, upon Srx-mediated reduction of Tsa1, Hsp104 induces disaggregation of protein aggregates. Hsp, heat shock protein; Tsa1, thiol-specific antioxidant 1.

the presence of fusion tags can affect oligomerization and hyperoxidation sensitivity. A C-terminal Myc-tag on PrxII conferred resistance to hyperoxidation (169), whereas an N-terminal His₆-tag on *Arabidopsis* 2-Cys PrxA increased its sensitivity to hyperoxidation (184) and an N-terminal Myc-tag on yeast Tsa1 destabilized its oligomeric forms (263). The oligomeric state of Prx can also be influenced by the presence of other proteins. In *Arabidopsis*, Cyp20-3, an abundant stromal protein, interacts with 2-Cys Prx with a K_d of $<0.5 \mu\text{M}$, in a pH- and redox-dependent manner. The Cyp20-3 association drives the oligomeric state of 2-Cys Prx toward the dimer. Conversely, titration of chloroplast stromal proteins with isolated *Arabidopsis* 2-Cys Prx shifted the oligomeric equilibrium from dimer toward decamer.

The influence of proteins on the oligomeric Prx state has also been characterized in eukaryotic systems, for instance the interaction between PrxI and p66Shc (110). In a coincidental link, p66Shc is also regulated by a cyclophilin, Pin1 (88). The apoptosis-inducing role of p66Shc is activated by a redox-dependent conversion from homodimers to disulfide-linked tetrameric p66Shc, which constitutes the proapoptotic form (111). The interaction with reduced homodimeric p66Shc induces the dissociation of PrxI decamers to dimers, purportedly due to the enhanced affinity of p66Shc for the dimeric form of PrxI (Fig. 10) (110). The association of oxidized tetrameric p66Shc with the decameric PrxI also generates a decamer-to-dimer conversion, although this time through a thiol–disulfide relay whereby oxidative equivalents are transferred from p66Shc to PrxI (110). It is still unclear whether PrxI can act as a reductant of p66Shc *in vivo*, whereas a role for Trx as a reductant of oxidized p66Shc has been described (111).

Finally, it is important to consider the common characteristics that define the specificity of Prx for its binding partners. Data-mining studies have unearthed recurrent motifs conserved among the interaction partners of human PrxI and PrxII, regardless of the interaction mode or oligomeric state of Prx (18). Besides the redox-active CXXC motif, many binding partners of PrxI/PrxII have been found to possess a LXXLL and/or PXXP motif. Among the functional Prx associations described above, both the LXXLL and PXXP motifs are conserved among STAT3, ASK1, and PTEN, and the PXXP motif alone for p66Shc, and the LXXLL motif alone for TLR4.

D. Complementary roles of Prx and Trx

Significantly, many of the physiologically interacting partners of Prx also interact with Trx. Most typically, the role of Trx in relation to its target proteins is that of a disulfide reductant that “flips the disulfide switch,” by reduction of either an intermolecular disulfide, as for p66Shc, or an intramolecular disulfide, as for example, cell division cycle 25 (Cdc25). Cdc25 is a PTP that is reversibly inactivated by H₂O₂-mediated oxidation, leading to intramolecular disulfide formation between its catalytic Cys and a backdoor Cys (346). Cdc25 participates in the activation of CDK1, a kinase that, as detailed above, phosphophorylates PrxI (45), thereby providing an indirect link between Trx action and phosphorylative Prx regulation. A more direct connection between kinases and Trx and Prx is the mutual interaction with MST1. Trx1 binds to the SARAH domain of MST1 and inhibits autophosphorylation by blocking its homodimerization

(44). Although the association of Trx1 to MST1 does not involve a mixed disulfide, it is still redox sensitive, as mutation of the CXXC motif cysteines to serine abolishes the interaction and Trx1 oxidation also induces the dissociation from MST1 (44). Whereas PrxI is a phosphorylation target of MST1, MST1 does not phosphorylate Trx1.

In addition to its redox-regulated interaction with PrxI, the phosphatase, PTEN, is also regulated in a redox-dependent manner by Trx. Remarkably, Trx can both activate and deactivate PTEN through distinct mechanisms. In its capacity as a disulfide reductase, Trx reduces the intramolecular disulfide that is formed in PTEN between the active-site Cys (Cys124) and the backdoor Cys (Cys71) upon oxidation by H₂O₂, converting PTEN from an inactive to an active state (204). PTEN inactivation by Trx might occur through the formation of a mixed disulfide between the catalytic Cys32 of Trx1 and Cys212 of the C2 domain of PTEN, inhibiting the lipid phosphatase activity (237). The inhibitory interaction between Trx and PTEN could be reversed by TR. Notably, this interaction involves the reduced form of Trx, with the oxidized form unable to interact with PTEN in an inhibitory fashion, thereby precluding a mechanism of thiol–disulfide exchange between Trx_{ox} and PTEN.

A final example of the dual regulation of a kinase by both Prx and Trx is in the case of ASK1. Whereas Prx is proposed to act as an oxidant transducer that oxidizes ASK1 to active, disulfide-linked multimers (180), Trx1 instead suppresses ASK1 activity through binding its N-terminal Trx-binding domain (TBD) and preventing the homophilic interaction of ASK1 required for its functionality (104, 185). Upon oxidation, Trx1 dissociates and ASK1 can convert into its active kinase form and participate in the initiation of the programmed cell death (316). The catalytic cysteine of Trx1 is essential for the maintenance of its affinity for ASK1 and of the seven conserved cysteines of the TBD of ASK1. Cys250 has been identified as the key residue modulating the association between ASK1 and Trx1 (185, 187, 252).

In all the cases of protein coregulation by Trx/Prx outlined above, an antagonistic system of regulation can be seen. Whereas PrxI acts as an activator of ASK1, MST1, and PTEN, Trx conversely acts as an inhibitor. Additionally, the redox stimuli for the interactions of Prx and Trx with these proteins are also in opposition. Under oxidizing conditions, an association of ASK1 and MST1 with PrxI is promoted, whereas under reducing conditions an interaction with Trx is created. A further significant characteristic of these interactions is that they are redox sensitive, but do not involve the oxidoreductase or chaperone functions of either Prx or Trx.

Although the role of Prx as a chaperone is focused on in the Nonperoxidatic roles for Prx section, Trx also has notable cellular chaperoning functions. The molecular chaperone capabilities of Trx and TR were first observed in *E. coli* (171). Since then, a critical role for Trxs as stress-responsive chaperones in plants has been established. Tobacco (*Nicotiana tabacum*) Trx-m and Trx-f and *Arabidopsis* Trx-h act as chaperones depending on their oligomeric states (273, 319). The monomeric form of Trx functions as a disulfide reductase and, under heat stress or oxidative stress, Trx oligomerizes and gains chaperone activity (273). A similar duality of function has been observed for plant NADPH-TR type C (a fusion of TR and Trx) that has a holdase and a disulfide reductase activity in HMW and LMW forms, respectively. Mutation of the active-site cysteines

abolished disulfide reductase and foldase activity, but the effect was less pronounced on the holdase activity (43). Recently, a novel bipartite protein with both designated reductase and chaperone domains has been identified, the *Arabidopsis thaliana* tetratricopeptide repeat domain-containing thiorodoxin (AtTDX). AtTDX has an N-terminal Hsp70-interacting protein (HIP)-like domain that contains three tetratricopeptide repeat domains and a C-terminal Trx domain conferring disulfide reductase activity. In response to heat stress, AtTDX switches from a LMW form with disulfide reductase activity to a HMW form with holdase function (198). AtTDX has been demonstrated to act as a cochaperone of the yeast Hsp70, Ssb2, modulating its ATPase activity and recruiting it to the Hsp90 complex (286). AtTDX associates *via* its HIP-like domain with the ATPase domain of Ssb2 in a redox-dependent interaction. The complex between AtTDX and Ssb2 dissociates under oxidative conditions and mutation of either the Cys20 of Ssb2, which is conserved by many other Hsp70 proteins, or of both cysteines of the catalytic CXXC motif of Trx, renders the association redox insensitive (389). AtTDX can also function as a general chaperone, acting as a foldase in its LMW form and, under heat-shock conditions, oligomerizes to a HMW form with holdase activity (198).

Thus far, beyond the respective peroxidase and reductase activities of Prx and Trx, a functional promiscuity relating to redox-related interactions and chaperoning is exhibited by both enzymes. Although Prx has been demonstrated to participate in oxidative signaling through redox relays (265, 345), its preferred oxidation target is undoubtedly Trx. The reduction of Trx by TR is expected to prevent Trx from further participation in an oxidative relay, but, under stress conditions, sufficient quantities of oxidized Trx may possibly accumulate to allow the Trx-mediated oxidation of target proteins in a relay system mirroring that displayed by Prx. In *S. pombe*, exposure to moderate oxidative stress (0.2 mM H₂O₂) overwhelms the TR system, resulting in high amounts of oxidized Trx1 either in an intramolecular disulfide-bonded form, or in mixed disulfides with various cellular proteins (65). A model of Trx-mediated oxidation of Yap1 has been suggested in *S. cerevisiae* (294) and the action of Trx as an oxidant has also been characterized in Δtrr *E. coli* (68, 352). Proteomic studies of *S. pombe* have revealed extensive thiol oxidation (31% of cysteines oxidized) in Δtrr cells, whereas only minor thiol oxidation (1.5% of cysteines oxidized) occurred in $\Delta trx1$ cells, further confirming the possible role of Trx as a cellular oxidant when the electron flow from the TR/NADPH system is limited (106).

In *Arabidopsis*, a physiologically relevant oxidative relay has been characterized between the atypical chloroplast Trx (AHT) and a cellular substrate protein. AHT has a non-canonical Trx active site and a higher redox potential than that of typical plant Trxs. The redox potentials of AHT1 and AHT4 at pH 7 are -237 and -240 mV, respectively (63), whereas other plant Trxs, such as Trx-m, has an E_m of approximately -300 mV (57). However, AHTs are still able to effectively act as electron donors for 2-Cys PrxA during peroxide reduction, despite the relatively low E_m of PrxA (-307 mV) (63, 142). A specific redox relay in *Arabidopsis* has been defined, whereby PrxA oxidizes AHT4, which in turn oxidizes a redox-sensitive subunit of the starch synthesis enzyme AGPase (APS1). This relay depended on the C-terminal domain of AHT4 and could not be substituted by AHT1, thereby demonstrating that for such an oxidative relay to oc-

cur, regions of the protein outside the redox-active site are required to confer binding specificity. Whereas Trx is kept in a reduced state by the TR/NADPH system, the rate of Prx-mediated oxidation of AHT exceeds its reduction rate under normal growth conditions. Therefore, a significant fraction of AHT can be expected to be oxidized, lending further relevance to its role as a transducer of oxidative signals.

VI. Conclusion

In considering the concept of intracellular redox signaling networks, we impose the simplification of separately defining *downstream*, *midstream*, and *upstream* redox-responsive proteins. Transcription factors of prokaryotes provide an example of direct redox regulation with immediate and measurable positive responses, namely upregulation of antioxidant enzymes. Important efforts have been made to understand the structural determinants that (i) allow some transcription factors to display oxidant sensitivity in the presence of ultrafast scavenging peroxidases, and (ii) modulate conformational changes mediating affinity and specificity of macromolecular associations (in this case toward DNA). A demonstrated asset of understanding the mode of oxidative structural regulation of the prokaryotic transcription factor, OxyR, is the development of a series of genetically encoded ratiometric H₂O₂ sensors, known as HyPer (22). The architecture of oxidative regulation of yeast transcription factors, such as Yap1, has been clearly established (387), with the first identification of a peroxidase redox relay mechanism in lieu of the direct peroxide responsiveness exhibited by prokaryotic transcription factors, such as OxyR or OhrR (258, 374). Our view of redox regulation of mammalian transcription factors can be partially obscured by additional regulatory mechanisms, such as phosphorylation by upstream kinases, which themselves are susceptible to redox regulation; examples being the p38 MAPK/JNK activation of FOXO transcription factors (84), or JAK2 activation of STAT3 (372). The additional axis of phosphorylative regulation of proteins can to some extent complicate the interpretation of oxidative signaling pathways.

Therefore, we have partially attempted to assess this issue through the discussion of some of the interregulatory interactions between antioxidant enzymes and kinases/phosphatases. A clear relationship between phosphorylation pathways and oxidative signaling has been demonstrated in vascular smooth muscle cells, in which a correlation between the H₂O₂ concentration and the tyrosine phosphorylation levels has been observed upon oxidative stimulation (356). As Prxs consume the vast majority (>99%) of the intracellular H₂O₂ (403), the ability of kinases/phosphatases to modulate their activity is a key intersect of the oxidative and phosphorylation signaling pathways and the significance of this has been emphasized previously (271). We have also outlined the redox-dependent noncovalent associations between kinases/phosphatases and Prx/Trx that do not involve a direct oxidation or reduction. It is important to note that Prx and Trx often play opposing regulatory roles in these associations and are differentially responsive to redox conditions. The emerging role of Prx as an intermediary transducer of oxidation through interprotein redox relays derives from the observation that typical redox-sensitive protein thiols are 4–7 orders of magnitude less reactive to peroxides (approximately 1–100 M⁻¹·s⁻¹) than the peroxidatic Prx thiol (~10⁵ to 10⁷ M⁻¹·s⁻¹) (296).

The remarkable complexity and functional promiscuity of mammalian antioxidant enzymes in comparison to their prokaryotic counterparts is evident, and several ground-breaking studies over the past decade have advanced our understanding of the role of eukaryotic antioxidant enzymes in oxidative signaling. In this study, we support the previous description of Prx as a “redox relay hub,” but wish to also draw focus to the role of redox-regulated moonlighting proteins as additional hubs, providing intersects between pathways of oxidative signaling and other cellular processes. In this review, we have touched upon a few of these moonlighting proteins,

It is also useful to highlight cases where a single kinase is targeted by multiple redox-responsive proteins, such as the regulation of ASK1 (the upstream signaler of the JNK/p38 pathway) by DJ-1, CypA, PrxI, and Trx, as this is indicative of a potential redox-regulated hub for kinase signaling (Fig. 12). All

Binding of oxidized DJ-1, reduced Trx, or CypA to ASK1 inhibits its kinase activity. Oxidation of ASK1 by PrxI converts ASK1 to disulfide-linked multimers that participate in the kinase signaling pathway. ASK1 phosphorylates Siah1 and activates the GAPDH-Siah1 nuclear signaling pathway. Reduction of PrxI by Trx or CypA prevents its oxidative activation by ASK1 and oxidation of Trx by PrxI is expected to induce its dissociation from ASK1. Cyp, cyclophilin; GAPDH, glyceraldehyde 3-phosphate dehydrogenase.

four of the moonlighting proteins discussed associate with p53, a tumor suppressor protein regarded as a key coordinator of the response to stress and cellular aging (Fig. 13) (218). DJ-1 engages in a redox-dependent inhibitory interaction with p53 (170), nuclear GAPDH forms a complex with p53 and promotes phosphorylation and expression of p53 (420), TG2 phosphorylates p53 when acting as a kinase (241), and p53 forms a complex with mitochondrial CypD to activate the CypD-dependent apoptotic pathway (21, 289).

Cellular localization is also likely to be a key factor in the cooperative roles of redox-responsive moonlighting proteins and obligate antioxidant enzymes. As a significant intracellular generator of ROS, and an influential coordinator of cell death, mitochondria present a subcellular space, wherein controlled redox regulation of processes is highly beneficial (306). Mitochondria-localized DJ-1 has been proposed to prevent degradation of the antiapoptotic mitochondrial protein, Bcl-X_L, and preserve the respiratory Complex I activity to limit mitochondrial ROS generation. CypD both acts to modulate Complex III, and forms part of mPTP, which mediates the collapse of the mitochondrial membrane potential as part of the cell death pathway. Other components of the mPTP include the voltage-dependent anion channel (VDAC), and adenine nucleotide translocase (ANT), which itself can be redox regulated through cysteine oxidation (58). GAPDH translocates to mitochondria during apoptosis and interacts with the

VDAC to promote pore opening (368). TG2 promotes both the upregulation and mitochondrial translocation of GAPDH, and is also able to act on ANT as both a transaminase and as a PDI (230). Overexpression of TG2 in the mitochondria leads to GSH depletion, increased ROS levels, and membrane hyperpolarization in a mechanism distinct from mPTP opening (282), and TG2 plays a critical role in autophagic clearance of damaged mitochondria in a manner dependent on its transaminase activity (311). Even proteins of this review not labeled as moonlighters appear to adopt a subfunction within mitochondria. STAT3, for example, has been recognized to act in mitochondria to partially suppress electron flow through Complex I and has a cytoprotective role independent of its function as a transcription factor (359).

Due to the limited scope of this review, several elements of cellular redox regulation have not been covered. Discussion of oxidative modifications, such as S-nitrosylation, glutathionylation, and persulfidation, were mostly excluded, although they provide an important spectrum of redox control that influences the fate of oxidative signaling. Furthermore, other posttranslational modifications, such as acetylation, provide a separate level of control over redox pathway enzymes (92), and redox control of deacetylases, such as sirtuins, links this regulatory level to redox conditions (292). Direct oxidative regulation of ion channels has been identified (26, 341, 383), and redox control of ion channels will likely have a marked

FIG. 13. Regulation of the tumor suppressor protein p53 by redox-sensitive moonlighting proteins. The moonlighting proteins, GAPDH, DJ-1, CypD, and TG2, functionally associate with p53. All functional annotations are derived from studies of the human isoforms. The nature of each interaction is annotated, and whether the association activates (by phosphorylation) or deactivates (by steric inhibition) p53 is indicated as well.

influence over the cellular response to oxidative stress, as Ca^{2+} influx for example can influence both redox-dependent and redox-independent signaling pathways alike. Understanding of the molecular mechanisms of ion channel redox regulation is relatively limited, in part due to the inherent difficulties of the structural characterization of membrane proteins. Inevitably, technical and methodological improvements in cryo-EM will facilitate the structural study of more membrane proteins that will translate in a greater access to the molecular mechanisms of redox-regulated ion channels.

The role of redox signaling in disease development is continuously validated (143), and, as recently stated, “An improved understanding of the sophisticated workings of redox biology is imperative to defeating cancer” (51). In this study, we hope to have emphasized that statement and encouraged consideration of moonlighting redox-regulated proteins as key players in deciding cellular fate under oxidative stress. CypA, TG2, CLIC (mentioned within the Transglutaminase 2 section), and DJ-1 are all associated with increased invasion and metastasis of cancer cells (133, 138, 197, 422), with both DJ-1 and CypA exerting their influence through the regulation of kinase pathways (133, 422). As cancer cells tend to block mitochondrial apoptotic induction, developments in cancer therapeutics often target the mitochondrial apoptosis signaling pathway, for example at p53 or the mPTP (105), and a direct link has been established between redox regulation of p53 and tumor progression (323). All of the moonlighting redox sensors described in this review play some role in regulating either p53, the mPTP, or both in response to redox stimuli. It is troubling that the precise molecular mechanisms governing the redox regulation of DJ-1, TG2, CLIC, and cyclophilins are still poorly understood.

Acknowledgments

The authors thank Martine De Cock for help in preparing the article. This work was supported by Vrije Universiteit Brussels (Strategic Research Program 34), the Research Foundation-Flanders (Excellence of Science project no. 30829584 [to F.V.B. and J.M.], the Research Foundation-Flanders (G0D7914N [to F.V.B.] and G0D70149 [to J.M.]), and the European Cooperation in Science and Research (COST Action BM1203/EU-ROS). B.P. and B.D.S. are indebted to the Agency for Innovation by Science and Technology, A.L. to VIB (through the International PhD Program in Life Sciences), and N.B. to Indian Council of Agricultural Research for predoctoral fellowships. M.-A.T. is a predoctoral fellow of the Research Foundation-Flanders.

References

1. Abo M and Weerapana E. A caged electrophilic probe for global analysis of cysteine reactivity in living cells. *J Am Chem Soc* 137: 7087–7090, 2015.
2. Albesa-Jove D, Comino N, Tersa M, Mohorko E, Urresti S, Dainese E, Chiarelli LR, Pasca MR, Manganelli R, Makarov V, Riccardi G, Svergun DI, Glockshuber R, and Guerin ME. The redox state regulates the conformation of Rv2466c to activate the antitubercular prodrug TP053. *J Biol Chem* 290: 31077–31089, 2015.
3. Albright BS, Kosinski A, Szczepaniak R, Cook EA, Stow ND, Conway JF, and Weller SK. The putative herpes simplex virus 1 chaperone protein UL32 modulates disulfide bond formation during infection. *J Virol* 89: 443–453, 2015.

4. Alekshun MN, Kim YS, and Levy SB. Mutational analysis of MarR, the negative regulator of marRAB expression in *Escherichia coli*, suggests the presence of two regions required for DNA binding. *Mol Microbiol* 35: 1394–1404, 2000.
5. Allan EG, Kander MC, Carmichael I, and Garman EF. To scavenge or not to scavenge, that is STILL the question. *J Synchrotron Radiat* 20: 23–36, 2013.
6. Andres-Mateos E, Perier C, Zhang L, Blanchard-Fillion B, Greco TM, Thomas B, Ko HS, Sasaki M, Ischiropoulos H, Przedborski S, Dawson TM, and Dawson VL. DJ-1 gene deletion reveals that DJ-1 is an atypical peroxiredoxin-like peroxidase. *Proc Natl Acad Sci U S A* 104: 14807–14812, 2007.
7. Antonella Nadalutti C, Korponay-Szabo IR, Kaukinen K, Wang Z, Griffin M, Maki M, and Lindfors K. Thioredoxin is involved in endothelial cell extracellular transglutaminase 2 activation mediated by celiac disease patient IgA. *PLoS One* 8: e77277, 2013.
8. Arias DG, Pineyro MD, Iglesias AA, Guerrero SA, and Robello C. Molecular characterization and interactome analysis of *Trypanosoma cruzi* tryparedoxin II. *J Proteomics* 120: 95–104, 2015.
9. Atichartpongkul S, Vattanaviboon P, Wisitkamol R, Jarosuk J, Mongkolsuk S, and Fuangthong M. Regulation of organic hydroperoxide stress response by two OhrR homologs in *Pseudomonas aeruginosa*. *PLoS One* 11: e0161982, 2016.
10. Azam S, Jouvett N, Jilani A, Vongsamphanh R, Yang X, Yang S, and Ramotar D. Human glyceraldehyde-3-phosphate dehydrogenase plays a direct role in reactivating oxidized forms of the DNA repair enzyme APE1. *J Biol Chem* 283: 30632–30641, 2008.
11. Basso M, Berlin J, Xia L, Sleiman SF, Ko B, Haskew-Layton R, Kim E, Antonyak MA, Cerione RA, Iismaa SE, Willis D, Cho S, and Ratan RR. Transglutaminase inhibition protects against oxidative stress-induced neuronal death downstream of pathological ERK activation. *J Neurosci* 32: 6561–6569, 2012.
12. Baty JW, Hampton MB, and Winterbourn CC. Proteomic detection of hydrogen peroxide-sensitive thiol proteins in Jurkat cells. *Biochem J* 389: 785–795, 2005.
13. Bedard K and Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 87: 245–313, 2007.
14. Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Tersikh AV, and Lukyanov S. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Methods* 3: 281–286, 2006.
15. Bergamini CM. GTP modulates calcium binding and cation-induced conformational changes in erythrocyte transglutaminase. *FEBS Lett* 239: 255–258, 1988.
16. Bernal-Bayard J and Ramos-Morales F. *Salmonella* type III secretion effector SlrP is an E3 ubiquitin ligase for mammalian thioredoxin. *J Biol Chem* 284: 27587–27595, 2009.
17. Bersweiler A, D’Autreaux B, Mazon H, Kriznik A, Belli G, Delaunay-Moisan A, Toledano MB, and Rahuel-Clermont S. A scaffold protein that chaperones a cysteine-sulfenic acid in H₂O₂ signaling. *Nat Chem Biol* 13: 909–915, 2017.
18. Bertoldi M. Human peroxiredoxins 1 and 2 and their interacting protein partners; through structure toward functions of biological complexes. *Protein Pept Lett* 23: 69–77, 2016.
19. Bhaskar A, Chawla M, Mehta M, Parikh P, Chandra P, Bhawe D, Kumar D, Carroll KS, and Singh A. Reengineering

- redox sensitive GFP to measure mycothiol redox potential of *Mycobacterium tuberculosis* during infection. *PLoS Pathog* 10: e1003902, 2014.
20. Bhatt MP, Lim YC, Hwang J, Na S, Kim YM, and Ha KS. C-peptide prevents hyperglycemia-induced endothelial apoptosis through inhibition of reactive oxygen species-mediated transglutaminase 2 activation. *Diabetes* 62: 243–253, 2013.
 21. Bigi A, Beltrami E, Trinei M, Stendardo M, Pelicci PG, and Giorgio M. Cyclophilin D counteracts P53-mediated growth arrest and promotes Ras tumorigenesis. *Oncogene* 35: 5132–5143, 2016.
 22. Bilan DS and Belousov VV. HyPer family probes: state of the art. *Antioxid Redox Signal* 24: 731–751, 2016.
 23. Bilan DS, Pase L, Joosen L, Gorokhovatsky AY, Ermakova YG, Gadella TW, Grabher C, Schultz C, Lukyanov S, and Belousov VV. HyPer-3: a genetically encoded H(2)O(2) probe with improved performance for ratio-metric and fluorescence lifetime imaging. *ACS Chem Biol* 8: 535–542, 2013.
 24. Blackinton J, Lakshminarasimhan M, Thomas KJ, Ahmad R, Greggio E, Raza AS, Cookson MR, and Wilson MA. Formation of a stabilized cysteine sulfinic acid is critical for the mitochondrial function of the parkinsonism protein DJ-1. *J Biol Chem* 284: 6476–6485, 2009.
 25. Bogdanova YA, Schultz C, and Belousov VV. Local generation and imaging of hydrogen peroxide in living cells. *Curr Protoc Chem Biol* 9: 117–127, 2017.
 26. Bogeski I, Kappl R, Kummerow C, Gulaboski R, Hoth M, and Niemeyer BA. Redox regulation of calcium ion channels: chemical and physiological aspects. *Cell Calcium* 50: 407–423, 2011.
 27. Bonini MG, Rota C, Tomasi A, and Mason RP. The oxidation of 2',7'-dichlorofluorescein to reactive oxygen species: a self-fulfilling prophesy? *Free Radic Biol Med* 40: 968–975, 2006.
 28. Boothe RL and Folk JE. A reversible, calcium-dependent, copper-catalyzed inactivation of guinea pig liver transglutaminase. *J Biol Chem* 244: 399–405, 1969.
 29. Bordelon T, Wilkinson SP, Grove A, and Newcomer ME. The crystal structure of the transcriptional regulator HucR from *Deinococcus radiodurans* reveals a repressor preconfigured for DNA binding. *J Mol Biol* 360: 168–177, 2006.
 30. Boronat S, Domenech A, and Hidalgo E. Proteomic characterization of reversible thiol oxidations in proteomes and proteins. *Antioxid Redox Signal* 26: 329–344, 2017.
 31. Boschi-Muller S and Branlant G. The active site of phosphorylating glyceraldehyde-3-phosphate dehydrogenase is not designed to increase the nucleophilicity of a serine residue. *Arch Biochem Biophys* 363: 259–266, 1999.
 32. Brugarolas P, Movahedzadeh F, Wang Y, Zhang N, Bartek IL, Gao YN, Voskuil MI, Franzblau SG, and He C. The oxidation-sensing regulator (MosR) is a new redox-dependent transcription factor in *Mycobacterium tuberculosis*. *J Biol Chem* 287: 37703–37712, 2012.
 33. Burbank L and Roper MC. OxyR and SoxR modulate the inducible oxidative stress response and are implicated during different stages of infection for the bacterial phytopathogen *Pantoea stewartii* subsp. *stewartii*. *Mol Plant Microbe Interact* 27: 479–490, 2014.
 34. Bustos PL, Volta BJ, Perrone AE, Milduburger N, and Bua J. A homolog of cyclophilin D is expressed in *Trypanosoma cruzi* and is involved in the oxidative stress-damage response. *Cell Death Discov* 3: 16092, 2017.
 35. Calvo IA, Boronat S, Domenech A, Garcia-Santamarina S, Ayte J, and Hidalgo E. Dissection of a redox relay: H₂O₂-dependent activation of the transcription factor Pap1 through the peroxidatic Tpx1-thioredoxin cycle. *Cell Rep* 5: 1413–1424, 2013.
 36. Campos-Acevedo AA and Rudino-Pinera E. Crystallographic studies evidencing the high energy tolerance to disrupting the interface disulfide bond of thioredoxin 1 from white leg shrimp *Litopenaeus vannamei*. *Molecules* 19: 21113–21126, 2014.
 37. Canet-Aviles RM, Wilson MA, Miller DW, Ahmad R, McLendon C, Bandyopadhyay S, Baptista MJ, Ringe D, Petsko GA, and Cookson MR. The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfinic acid-driven mitochondrial localization. *Proc Natl Acad Sci U S A* 101: 9103–9108, 2004.
 38. Cao J, Ying M, Xie N, Lin G, Dong R, Zhang J, Yan H, Yang X, He Q, and Yang B. The oxidation states of DJ-1 dictate the cell fate in response to oxidative stress triggered by 4-hpr: autophagy or apoptosis? *Antioxid Redox Signal* 21: 1443–1459, 2014.
 39. Cao JX, Schulte J, Knight A, Leslie NR, Zagodzdon A, Bronson R, Manevich Y, Beeson C, and Neumann CA. Prdx1 inhibits tumorigenesis via regulating PTEN/AKT activity. *EMBO J* 28: 1505–1517, 2009.
 40. Cao Z, Tavender TJ, Roszak AW, Cogdell RJ, and Bulleid NJ. Crystal structure of reduced and of oxidized peroxiredoxin IV enzyme reveals a stable oxidized decamer and a non-disulfide-bonded intermediate in the catalytic cycle. *J Biol Chem* 286: 42257–42266, 2011.
 41. Caporaletti D, D'Alessio AC, Rodriguez-Suarez RJ, Senn AM, Duek PD, and Woloskiuk RA. Non-reductive modulation of chloroplast fructose-1,6-bisphosphatase by 2-Cys peroxiredoxin. *Biochem Biophys Res Commun* 355: 722–727, 2007.
 42. Cerveau D, Kraut A, Stotz HU, Mueller MJ, Coute Y, and Rey P. Characterization of the *Arabidopsis thaliana* 2-Cys peroxiredoxin interactome. *Plant Sci* 252: 30–41, 2016.
 43. Chae HB, Moon JC, Shin MR, Chi YH, Jung YJ, Lee SY, Nawkar GM, Jung HS, Hyun JK, Kim WY, Kang CH, Yun DJ, Lee KO, and Lee SY. Thioredoxin reductase Type C (NTRC) orchestrates enhanced thermotolerance to *Arabidopsis* by its redox-dependent holdase chaperone function. *Mol Plant* 6: 323–336, 2013.
 44. Chae JS, Gil Hwang S, Lim DS, and Choi EJ. Thioredoxin-1 functions as a molecular switch regulating the oxidative stress-induced activation of MST1. *Free Radic Biol Med* 53: 2335–2343, 2012.
 45. Chang TS, Jeong W, Choi SY, Yu S, Kang SW, and Rhee SG. Regulation of peroxiredoxin I activity by Cdc2-mediated phosphorylation. *J Biol Chem* 277: 25370–25376, 2002.
 46. Chen CH, Li WZ, Sultana R, You MH, Kondo A, Shahpasand K, Kim BM, Luo ML, Nechama M, Lin YM, Yao YD, Lee TH, Zhou XZ, Swomley AM, Butterfield DA, Zhang Y, and Lu KP. Pin1 cysteine-113 oxidation inhibits its catalytic activity and cellular function in Alzheimer's disease. *Neurobiol Dis* 76: 13–23, 2015.
 47. Chen H, Hu J, Chen PR, Lan L, Li Z, Hicks LM, Dinner AR, and He C. The *Pseudomonas aeruginosa* multidrug efflux regulator MexR uses an oxidation-sensing mechanism. *Proc Natl Acad Sci U S A* 105: 13586–13591, 2008.
 48. Chen H, Yi C, Zhang J, Zhang W, Ge Z, Yang CG, and He C. Structural insight into the oxidation-sensing mechanism of

- the antibiotic resistance of regulator MexR. *EMBO Rep* 11: 685–690, 2010.
49. Chen J, Li L, and Chin LS. Parkinson disease protein DJ-1 converts from a zymogen to a protease by carboxyl-terminal cleavage. *Hum Mol Genet* 19: 2395–2408, 2010.
 50. Chen RW, Saunders PA, Wei HF, Li ZW, Seth P, and Chuang DM. Involvement of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and p53 in neuronal apoptosis: evidence that GAPDH is upregulated by p53. *J Neurosci* 19: 9654–9662, 1999.
 51. Chio IIC and Tuveson DA. ROS in cancer: the burning question. *Trends Mol Med* 23: 411–429, 2017.
 52. Cho CS, Yoon HJ, Kim JY, Woo HA, and Rhee SG. Circadian rhythm of hyperoxidized peroxiredoxin II is determined by hemoglobin autoxidation and the 20S proteasome in red blood cells. *Proc Natl Acad Sci U S A* 111: 12043–12048, 2014.
 53. Choi H, Kim S, Mukhopadhyay P, Cho S, Woo J, Storz G, and Ryu SE. Structural basis of the redox switch in the OxyR transcription factor. *Cell* 105: 103–113, 2001.
 54. Choi H-J, Kang SW, Yang C-H, Rhee SG, and Ryu S-E. Crystal structure of a novel human peroxidase enzyme at 2.0 Å resolution. *Nat Struct Biol* 5: 400–406, 1998.
 55. Choi-Rhee E, Schulman H, and Cronan JE. Promiscuous protein biotinylation by *Escherichia coli* biotin protein ligase. *Protein Sci* 13: 3043–3050, 2004.
 56. Clements CM, McNally RS, Conti BJ, Mak TW, and Ting JP. DJ-1, a cancer- and Parkinson's disease-associated protein, stabilizes the antioxidant transcriptional master regulator Nrf2. *Proc Natl Acad Sci U S A* 103: 15091–15096, 2006.
 57. Collin V, Issakidis-Bourguet E, Marchand C, Hirasawa M, Lancelin JM, Knaff DB, and Miginiac-Maslow M. The *Arabidopsis* plastidial thioredoxins: new functions and new insights into specificity. *J Biol Chem* 278: 23747–23752, 2003.
 58. Costantini P, Belzacq AS, Vieira HL, Larochette N, de Pablo MA, Zamzami N, Susin SA, Brenner C, and Kroemer G. Oxidation of a critical thiol residue of the adenine nucleotide translocator enforces Bcl-2-independent permeability transition pore opening and apoptosis. *Oncogene* 19: 307–314, 2000.
 59. Crane EJ, 3rd, Vervoort J, and Claiborne A. 13C NMR analysis of the cysteine-sulfenic acid redox center of enterococcal NADH peroxidase. *Biochemistry* 36: 8611–8618, 1997.
 60. Cun Y, Dai N, Li M, Xiong C, Zhang Q, Sui J, Qian C, and Wang D. APE1/Ref-1 enhances DNA binding activity of mutant p53 in a redox-dependent manner. *Oncol Rep* 31: 901–909, 2014.
 61. da Silva Neto JF, Negretto CC, and Netto LE. Analysis of the organic hydroperoxide response of *Chromobacterium violaceum* reveals that OhrR is a cys-based redox sensor regulated by thioredoxin. *PLoS One* 7: e47090, 2012.
 62. Dagnell M, Pace PE, Cheng Q, Frijhoff J, Ostman A, Arner ESJ, Hampton MB, and Winterbourn CC. Thioredoxin reductase 1 and NADPH directly protect protein tyrosine phosphatase 1B from inactivation during H₂O₂ exposure. *J Biol Chem* 292: 14371–14380, 2017.
 63. Dangoor I, Peled-Zehavi H, Levitan A, Pasand O, and Danon A. A small family of chloroplast atypical thioredoxins. *Plant Physiol* 149: 1240–1250, 2009.
 64. Darnell JE, Jr., Kerr IM, and Stark GR. Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* 264: 1415–1421, 1994.
 65. Day AM, Brown JD, Taylor SR, Rand JD, Morgan BA, and Veal EA. Inactivation of a peroxiredoxin by hydrogen peroxide is critical for thioredoxin-mediated repair of oxidized proteins and cell survival. *Mol Cell* 45: 398–408, 2012.
 66. de Keizer PL, Burgering BM, and Dansen TB. Forkhead box o as a sensor, mediator, and regulator of redox signaling. *Antioxid Redox Signal* 14: 1093–1106, 2011.
 67. De Munter S, Gornemann J, Derua R, Lesage B, Qian J, Heroes E, Waelkens E, Van Eynde A, Beullens M, and Bollen M. Split-BioID: a proximity biotinylation assay for dimerization-dependent protein interactions. *FEBS Lett* 591: 415–424, 2017.
 68. Debarbieux L and Beckwith J. The reductive enzyme thioredoxin 1 acts as an oxidant when it is exported to the *Escherichia coli* periplasm. *Proc Natl Acad Sci U S A* 95: 10751–10756, 1998.
 69. Delaunay A, Isnard AD, and Toledano MB. H₂O₂ sensing through oxidation of the Yap1 transcription factor. *EMBO J* 19: 5157–5166, 2000.
 70. Delaunay A, Pflieger D, Barrault MB, Vinh J, and Toledano MB. A thiol peroxidase is an H₂O₂ receptor and redox-transducer in gene activation. *Cell* 111: 471–481, 2002.
 71. Demaria M, Giorgi C, Lebedzinska M, Esposito G, D'Angeli L, Bartoli A, Gough DJ, Turkson J, Levy DE, Watson CJ, Wieckowski MR, Provero P, Pinton P, and Poli V. A STAT3-mediated metabolic switch is involved in tumour transformation and STAT3 addiction. *Aging (Albany NY)* 2: 823–842, 2010.
 72. Denu JM and Tanner KG. Specific and reversible inactivation of protein tyrosine phosphatases by hydrogen peroxide: evidence for a sulfenic acid intermediate and implications for redox regulation. *Biochemistry* 37: 5633–5642, 1998.
 73. Dickinson BC and Chang CJ. A targetable fluorescent probe for imaging hydrogen peroxide in the mitochondria of living cells. *J Am Chem Soc* 130: 9638–9639, 2008.
 74. Dickinson BC, Huynh C, and Chang CJ. A palette of fluorescent probes with varying emission colors for imaging hydrogen peroxide signaling in living cells. *J Am Chem Soc* 132: 5906–5915, 2010.
 75. Dickinson BC, Tang Y, Chang Z, and Chang CJ. A nuclear-localized fluorescent hydrogen peroxide probe for monitoring sirtuin-mediated oxidative stress responses in vivo. *Chem Biol* 18: 943–948, 2011.
 76. Dominguez-Solis JR, He Z, Lima A, Ting J, Buchanan BB, and Luan S. A cyclophilin links redox and light signals to cysteine biosynthesis and stress responses in chloroplasts. *Proc Natl Acad Sci U S A* 105: 16386–16391, 2008.
 77. Dooley CT, Dore TM, Hanson GT, Jackson WC, Remington SJ, and Tsien RY. Imaging dynamic redox changes in mammalian cells with green fluorescent protein indicators. *J Biol Chem* 279: 22284–22293, 2004.
 78. Duan X, Kelsen SG, and Merali S. Proteomic analysis of oxidative stress-responsive proteins in human pneumocytes: insight into the regulation of DJ-1 expression. *J Proteome Res* 7: 4955–4961, 2008.
 79. Duhe RJ, Evans GA, Erwin RA, Kirken RA, Cox GW, and Farrar WL. Nitric oxide and thiol redox regulation of Janus kinase activity. *Proc Natl Acad Sci U S A* 95: 126–131, 1998.
 80. Durand A, Papai G, and Schultz P. Structure, assembly and dynamics of macromolecular complexes by single

- particle cryo-electron microscopy. *J Nanobiotechnol* 11(Suppl 1): S4, 2013.
81. Eggler AL, Small E, Hannink M, and Mesecar AD. Cul3-mediated Nrf2 ubiquitination and antioxidant response element (ARE) activation are dependent on the partial molar volume at position 151 of Keap1. *Biochem J* 422: 171–180, 2009.
 82. Ermakova YG, Bilan DS, Matlashov ME, Mishina NM, Markvicheva KN, Subach OM, Subach FV, Bogeski I, Hoth M, Enikolopov G, and Belousov VV. Red fluorescent genetically encoded indicator for intracellular hydrogen peroxide. *Nat Commun* 5: 5222, 2014.
 83. Espinosa A, Garcia A, Hartel S, Hidalgo C, and Jaimovich E. NADPH oxidase and hydrogen peroxide mediate insulin-induced calcium increase in skeletal muscle cells. *J Biol Chem* 284: 2568–2575, 2009.
 84. Essers MA, Weijzen S, de Vries-Smits AM, Saarloos I, de Ruiter ND, Bos JL, and Burgering BM. FOXO transcription factor activation by oxidative stress mediated by the small GTPase Ral and JNK. *EMBO J* 23: 4802–4812, 2004.
 85. Etzler JC, Bollo M, Holstein D, Deng JJ, Perez V, Lin DT, Richardson A, Bai Y, and Lechleiter JD. Cyclophilin D over-expression increases mitochondrial complex III activity and accelerates supercomplex formation. *Arch Biochem Biophys* 613: 61–68, 2017.
 86. Fan J, Ren H, Jia N, Fei E, Zhou T, Jiang P, Wu M, and Wang G. DJ-1 decreases Bax expression through repressing p53 transcriptional activity. *J Biol Chem* 283: 4022–4030, 2008.
 87. Fan Y, Chen Z, and Ai HW. Monitoring redox dynamics in living cells with a redox-sensitive red fluorescent protein. *Anal Chem* 87: 2802–2810, 2015.
 88. Feng D, Yao J, Wang G, Li Z, Zu G, Li Y, Luo F, Ning S, Qasim W, Chen Z, and Tian X. Inhibition of p66Shc-mediated mitochondrial apoptosis via targeting prolyl-isomerase Pin1 attenuates intestinal ischemia/reperfusion injury in rats. *Clin Sci (Lond)* 131: 759–773, 2017.
 89. Fernandez-Caggiano M, Schroder E, Cho HJ, Burgoyne J, Barallobre-Barreiro J, Mayr M, and Eaton P. Oxidant-induced interprotein disulfide formation in cardiac protein DJ-1 occurs via an interaction with peroxiredoxin 2. *J Biol Chem* 291: 10399–10410, 2016.
 90. Ferrer-Sueta G, Manta B, Botti H, Radi R, Trujillo M, and Denicola A. Factors affecting protein thiol reactivity and specificity in peroxide reduction. *Chem Res Toxicol* 24: 434–450, 2011.
 91. Fields S and Song O. A novel genetic system to detect protein-protein interactions. *Nature* 340: 245–246, 1989.
 92. Fiskus W, Coothankandaswamy V, Chen J, Ma H, Ha K, Saenz DT, Krieger SS, Mill CP, Sun B, Huang P, Mumm JS, Melnick AM, and Bhalla KN. SIRT2 deacetylates and inhibits the peroxidase activity of peroxiredoxin-1 to sensitize breast cancer cells to oxidant stress-inducing agents. *Cancer Res* 76: 5467–5478, 2016.
 93. Fiucci G, Beaucourt S, Duflaut D, Lespagnol A, Stumptner-Cuvelette P, Geant A, Buchwalter G, Tuynder M, Susini L, Lassalle JM, Wasylyk C, Wasylyk B, Oren M, Amson R, and Telferman A. Siah-1b is a direct transcriptional target of p53: identification of the functional p53 responsive element in the siah-1b promoter. *Proc Natl Acad Sci U S A* 101: 3510–3515, 2004.
 94. Folda A, Citta A, Scalcon V, Cali T, Zonta F, Scutari G, Bindoli A, and Rigobello MP. Mitochondrial thioredoxin system as a modulator of cyclophilin D redox state. *Sci Rep* 6: 23071, 2016.
 95. Fomenko DE, Koc A, Agisheva N, Jacobsen M, Kaya A, Malinouski M, Rutherford JC, Siu KL, Jin DY, Winge DR, and Gladyshev VN. Thiol peroxidases mediate specific genome-wide regulation of gene expression in response to hydrogen peroxide. *Proc Natl Acad Sci U S A* 108: 2729–2734, 2011.
 96. Forman-Kay JD, Clore GM, and Gronenborn AM. Relationship between electrostatics and redox function in human thioredoxin: characterization of pH titration shifts using two-dimensional homo- and heteronuclear NMR. *Biochemistry* 31: 3442–3452, 1992.
 97. Forman HJ, Davies MJ, Kramer AC, Miotto G, Zaccarin M, Zhang H, and Ursini F. Protein cysteine oxidation in redox signaling: caveats on sulfenic acid detection and quantification. *Arch Biochem Biophys* 617: 26–37, 2017.
 98. Forman HJ and Fridovich I. Superoxide dismutase: a comparison of rate constants. *Arch Biochem Biophys* 158: 396–400, 1973.
 99. Forman HJ, Maiorino M, and Ursini F. Signaling functions of reactive oxygen species. *Biochemistry* 49: 835–842, 2010.
 100. Forman HJ, Ursini F, and Maiorino M. An overview of mechanisms of redox signaling. *J Mol Cell Cardiol* 73: 2–9, 2014.
 101. Forred BJ, Neuhauser S, Kim DI, Amolins MW, Motamedchaboki K, Roux KJ, and Vitiello PF. Identification of redox and glucose-dependent Txnip protein interactions. *Oxid Med Cell Longev* 2016: 5829063, 2016.
 102. Froyset AK, Edson AJ, Gharbi N, Khan EA, Dondorp D, Bai Q, Tiraboschi E, Suster ML, Connolly JB, Burton EA, and Fladmark KE. Astroglial DJ-1 over-expression up-regulates proteins involved in redox regulation and is neuroprotective in vivo. *Redox Biol* 16: 237–247, 2018.
 103. Fu C, Wu C, Liu T, Ago T, Zhai P, Sadoshima J, and Li H. Elucidation of thioredoxin target protein networks in mouse. *Mol Cell Proteomics* 8: 1674–1687, 2009.
 104. Fujino G, Noguchi T, Matsuzawa A, Yamauchi S, Saitoh M, Takeda K, and Ichijo H. Thioredoxin and TRAF family proteins regulate reactive oxygen species-dependent activation of ASK1 through reciprocal modulation of the N-terminal homophilic interaction of ASK1. *Mol Cell Biol* 27: 8152–8163, 2007.
 105. Galluzzi L, Larochette N, Zamzami N, and Kroemer G. Mitochondria as therapeutic targets for cancer chemotherapy. *Oncogene* 25: 4812–4830, 2006.
 106. Garcia-Santamarina S, Boronat S, Calvo IA, Rodriguez-Gabriel M, Ayte J, Molina H, and Hidalgo E. Is oxidized thioredoxin a major trigger for cysteine oxidation? Clues from a redox proteomics approach. *Antioxid Redox Signal* 18: 1549–1556, 2013.
 107. Garcia-Santamarina S, Boronat S, Domenech A, Ayte J, Molina H, and Hidalgo E. Monitoring in vivo reversible cysteine oxidation in proteins using ICAT and mass spectrometry. *Nat Protoc* 9: 1131–1145, 2014.
 108. Garcia-Santamarina S, Boronat S, Espadas G, Ayte J, Molina H, and Hidalgo E. The oxidized thiol proteome in fission yeast—optimization of an ICAT-based method to identify H₂O₂-oxidized proteins. *J Proteomics* 74: 2476–2486, 2011.
 109. Gautier V, Le HT, Malki A, Messaoudi N, Caldas T, Kthiri F, Landoulsi A, and Richarme G. YajL, the prokaryotic homolog of the Parkinsonism-associated protein

- DJ-1, protects cells against protein sulfenylation. *J Mol Biol* 421: 662–670, 2012.
110. Gertz M, Fischer F, Leipelt M, Wolters D, and Steegborn C. Identification of peroxiredoxin 1 as a novel interaction partner for the lifespan regulator protein p66Shc. *Aging (Albany NY)* 1: 254–265, 2009.
 111. Gertz M, Fischer F, Wolters D, and Steegborn C. Activation of the lifespan regulator p66Shc through reversible disulfide bond formation. *Proc Natl Acad Sci U S A* 105: 5705–5709, 2008.
 112. Getz EB, Xiao M, Chakrabarty T, Cooke R, and Selvin PR. A comparison between the sulphydryl reductants tris(2-carboxyethyl)phosphine and dithiothreitol for use in protein biochemistry. *Anal Biochem* 273: 73–80, 1999.
 113. Giorgio M, Migliaccio E, Orsini F, Paolucci D, Moroni M, Contursi C, Pelliccia G, Luzi L, Minucci S, Marcaccio M, Pinton P, Rizzuto R, Bernardi P, Paolucci F, and Pelicci PG. Electron transfer between cytochrome c and p66Shc generates reactive oxygen species that trigger mitochondrial apoptosis. *Cell* 122: 221–233, 2005.
 114. Glaeser RM. How good can cryo-EM become? *Nat Methods* 13: 28–32, 2016.
 115. Goldberg MS and Lansbury PT. Is there a cause-and-effect relationship between alpha-synuclein fibrillization and Parkinson's disease? *Nat Cell Biol* 2: 115–119, 2000.
 116. Gomes A, Fernandes E, and Lima JL. Fluorescence probes used for detection of reactive oxygen species. *J Biochem Biophys Methods* 65: 45–80, 2005.
 117. Graf PC, Martinez-Yamout M, VanHaerents S, Lilie H, Dyson HJ, and Jakob U. Activation of the redox-regulated chaperone Hsp33 by domain unfolding. *J Biol Chem* 279: 20529–20538, 2004.
 118. Gray MJ, Li Y, Leichert LI, Xu Z, and Jakob U. Does the transcription factor NemR use a regulatory sulfenamide bond to sense bleach? *Antioxid Redox Signal* 23: 747–754, 2015.
 119. Griendling KK, Sorescu D, and Ushio-Fukai M. NAD(P)H oxidase: role in cardiovascular biology and disease. *Circ Res* 86: 494–501, 2000.
 120. Grison A, Mantovani F, Comel A, Agostoni E, Gustincich S, Persichetti F, and Del Sal G. Ser46 phosphorylation and prolyl-isomerase Pin1-mediated isomerization of p53 are key events in p53-dependent apoptosis induced by mutant huntingtin. *Proc Natl Acad Sci U S A* 108: 17979–17984, 2011.
 121. Guimaraes BG, Souchon H, Honore N, Saint-Joanis B, Brosch R, Shepard W, Cole ST, and Alzari PM. Structure and mechanism of the alkyl hydroperoxidase AhpC, a key element of the *Mycobacterium tuberculosis* defense system against oxidative stress. *J Biol Chem* 280: 25735–25742, 2005.
 122. Gulshan K, Lee SS, and Moye-Rowley WS. Differential oxidant tolerance determined by the key transcription factor Yap1 is controlled by levels of the Yap1-binding protein, Ybp1. *J Biol Chem* 286: 34071–34081, 2011.
 123. Gupta A, Fuentes SM, and Grove A. Redox-sensitive MarR homologue BifR from *Burkholderia thailandensis* regulates biofilm formation. *Biochemistry* 56: 2315–2327, 2017.
 124. Gupta V and Carroll KS. Sulfenic acid chemistry, detection and cellular lifetime. *Biochim Biophys Acta* 1840: 847–875, 2014.
 125. Gutscher M, Pauleau AL, Marty L, Brach T, Wabnitz GH, Samstag Y, Meyer AJ, and Dick TP. Real-time imaging of the intracellular glutathione redox potential. *Nat Methods* 5: 553–559, 2008.
 126. Gutscher M, Sobotta MC, Wabnitz GH, Ballikaya S, Meyer AJ, Samstag Y, and Dick TP. Proximity-based protein thiol oxidation by H₂O₂-scavenging peroxidases. *J Biol Chem* 284: 31532–31540, 2009.
 127. Han BG, Cho JW, Cho YD, Jeong KC, Kim SY, and Lee BI. Crystal structure of human transglutaminase 2 in complex with adenosine triphosphate. *Int J Biol Macromol* 47: 190–195, 2010. *Erratum* 106: 1330, 2018.
 128. Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, and Remington SJ. Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. *J Biol Chem* 279: 13044–13053, 2004.
 129. Hanzen S, Vielfort K, Yang JS, Roger F, Andersson V, Zamarbide-Fores S, Andersson R, Malm L, Palais G, Biteau B, Liu BD, Toledano MB, Molin M, and Nystrom T. Lifespan control by redox-dependent recruitment of chaperones to misfolded proteins. *Cell* 166: 140–151, 2016.
 130. Hara MR, Agrawal N, Kim SF, Cascio MB, Fujimuro M, Ozeki Y, Takahashi M, Cheah JH, Tankou SK, Hester LD, Ferris CD, Hayward SD, Snyder SH, and Sawa A. S-nitrosylated GAPDH initiates apoptotic cell death by nuclear translocation following Siah1 binding. *Nat Cell Biol* 7: 665–674, 2005.
 131. Hasegawa G, Suwa M, Ichikawa Y, Ohtsuka T, Kumagai S, Kikuchi M, Sato Y, and Saito Y. A novel function of tissue-type transglutaminase: protein disulphide isomerase. *Biochem J* 373: 793–803, 2003.
 132. Hayashi T, Ishimori C, Takahashi-Niki K, Taira T, Kim YC, Maita H, Maita C, Ariga H, and Iguchi-Ariga SM. DJ-1 binds to mitochondrial complex I and maintains its activity. *Biochem Biophys Res Commun* 390: 667–672, 2009.
 133. He X, Zheng Z, Li J, Ben Q, Liu J, Zhang J, Ji J, Yu B, Chen X, Su L, Zhou L, Liu B, and Yuan Y. DJ-1 promotes invasion and metastasis of pancreatic cancer cells by activating SRC/ERK/uPA. *Carcinogenesis* 33: 555–562, 2012.
 134. Hendrickson WA. Atomic-level analysis of membrane-protein structure. *Nat Struct Mol Biol* 23: 464–467, 2016.
 135. Heo JY, Park JH, Kim SJ, Seo KS, Han JS, Lee SH, Kim JM, Park JI, Park SK, Lim K, Hwang BD, Shong M, and Kweon GR. DJ-1 null dopaminergic neuronal cells exhibit defects in mitochondrial function and structure: involvement of mitochondrial complex I assembly. *PLoS One* 7: e32629, 2012.
 136. Heppner DE, Hristova M, Ida T, Mijuskovic A, Dustin CM, Bogdandi V, Fukuto JM, Dick TP, Nagy P, Li J, Akaike T, and van der Vliet A. Cysteine perthiosulfenic acid (Cys-SSOH): a novel intermediate in thiol-based redox signaling? *Redox Biol* 14: 379–385, 2018.
 137. Heppner DE, Janssen-Heininger YMW, and van der Vliet A. The role of sulfenic acids in cellular redox signaling: reconciling chemical kinetics and molecular detection strategies. *Arch Biochem Biophys* 616: 40–46, 2017.
 138. Hernandez-Fernaund JR, Ruengeler E, Casazza A, Neilson LJ, Pulleine E, Santi A, Ismail S, Lilla S, Dhayade S, MacPherson IR, McNeish I, Ennis D, Ali H, Kugerski FG, Al Khamici H, van den Biggelaar M, van den Bergh PV, Cloix C, McDonald L, Millan D, Hoyle A, Kuchnio A, Carmeliet P, Valenzuela SM, Blyth K, Yin H, Mazzone M, Norman JC, and Zanivan S. Secreted CLIC3 drives cancer progression through its glutathione-dependent oxidoreductase activity. *Nat Commun* 8: 14206, 2017.

139. Hod Y, Pentyala SN, Whyard TC, and El-Maghrabi MR. Identification and characterization of a novel protein that regulates RNA-protein interaction. *J Cell Biochem* 72: 435–444, 1999.
140. Holland SM. Chronic granulomatous disease. *Hematol Oncol Clin North Am* 27: 89–99, 2013.
141. Hopkins BL, Nadler M, Skoko JJ, Bertomeu T, Pelosi A, Shafaei PM, Levine K, Schempf A, Pennarun B, Yang B, Datta D, Bucur O, Ndebele K, Oesterreich S, Yang D, Giulia Rizzo M, Khosravi-Far R, and Neumann CA. A peroxidase peroxiredoxin 1-specific redox regulation of the novel FOXO3 microRNA target let-7. *Antioxid Redox Signal* 28: 62–77, 2018.
142. Horling F, Lamkemeyer P, Konig J, Finkemeier I, Kandlbinder A, Baier M, and Dietz KJ. Divergent light-, ascorbate-, and oxidative stress-dependent regulation of expression of the peroxiredoxin gene family in Arabidopsis. *Plant Physiol* 131: 317–325, 2003.
143. Hornsveid M and Dansen TB. The hallmarks of cancer from a redox perspective. *Antioxid Redox Signal* 25: 300–325, 2016.
144. Huang Q, Lan F, Zheng Z, Xie F, Han J, Dong L, Xie Y, and Zheng F. Akt2 kinase suppresses glyceraldehyde-3-phosphate dehydrogenase (GAPDH)-mediated apoptosis in ovarian cancer cells via phosphorylating GAPDH at threonine 237 and decreasing its nuclear translocation. *J Biol Chem* 286: 42211–42220, 2011.
145. Hwang J, Ing MH, Salazar A, Lassegue B, Griendling K, Navab M, Sevanian A, and Hsiai TK. Pulsatile versus oscillatory shear stress regulates NADPH oxidase subunit expression: implication for native LDL oxidation. *Circ Res* 93: 1225–1232, 2003.
146. Hwang NR, Yim SH, Kim YM, Jeong J, Song EJ, Lee Y, Lee JH, Choi S, and Lee KJ. Oxidative modifications of glyceraldehyde-3-phosphate dehydrogenase play a key role in its multiple cellular functions. *Biochem J* 423: 253–264, 2009.
147. Im JY, Lee KW, Woo JM, Junn E, and Mouradian MM. DJ-1 induces thioredoxin 1 expression through the Nrf2 pathway. *Hum Mol Genet* 21: 3013–3024, 2012.
148. Imlay JA. Transcription factors that defend bacteria against reactive oxygen species. *Annu Rev Microbiol* 69: 93–108, 2015.
149. Innes BT, Sowole MA, Gyenis L, Dubinsky M, Konermann L, Litchfield DW, Brandl CJ, and Shilton BH. Peroxide-mediated oxidation and inhibition of the peptidyl-prolyl isomerase Pin1. *Biochim Biophys Acta* 1852: 905–912, 2015.
150. Irrcher I, Aleyasin H, Seifert EL, Hewitt SJ, Chhabra S, Phillips M, Lutz AK, Rousseaux MW, Bevilacqua L, Jahani-Asl A, Callaghan S, MacLaurin JG, Winklhofer KF, Rizzo P, Rippstein P, Kim RH, Chen CX, Fon EA, Slack RS, Harper ME, McBride HM, Mak TW, and Park DS. Loss of the Parkinson's disease-linked gene DJ-1 perturbs mitochondrial dynamics. *Hum Mol Genet* 19: 3734–3746, 2010.
151. Itoh K, Chiba T, Takahashi S, Ishii T, Igarashi K, Katoh Y, Oyake T, Hayashi N, Satoh K, Hatayama I, Yamamoto M, and Nabeshima Y. An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochem Biophys Res Commun* 236: 313–322, 1997.
152. Iwai K, Shibukawa Y, Yamazaki N, and Wada Y. Transglutaminase 2-dependent deamidation of glyceraldehyde-3-phosphate dehydrogenase promotes trophoblastic cell fusion. *J Biol Chem* 289: 4989–4999, 2014.
153. Jaffrey SR, Erdjument-Bromage H, Ferris CD, Tempst P, and Snyder SH. Protein S-nitrosylation: a physiological signal for neuronal nitric oxide. *Nat Cell Biol* 3: 193–197, 2001.
154. Jang GY, Jeon JH, Cho SY, Shin DM, Kim CW, Jeong EM, Bae HC, Kim TW, Lee SH, Choi Y, Lee DS, Park SC, and Kim IG. Transglutaminase 2 suppresses apoptosis by modulating caspase 3 and NF-kappaB activity in hypoxic tumor cells. *Oncogene* 29: 356–367, 2010.
155. Jang HH, Kim SY, Park SK, Jeon HS, Lee YM, Jung JH, Lee SY, Chae HB, Jung YJ, Lee KO, Lim CO, Chung WS, Bahk JD, Yun DJ, Cho MJ, and Lee SY. Phosphorylation and concomitant structural changes in human 2-Cys peroxiredoxin isotype I differentially regulate its peroxidase and molecular chaperone functions. *FEBS Lett* 580: 351–355, 2006.
156. Jang TH, Lee DS, Choi K, Jeong EM, Kim IG, Kim YW, Chun JN, Jeon JH, and Park HH. Crystal structure of transglutaminase 2 with GTP complex and amino acid sequence evidence of evolution of GTP binding site. *PLoS One* 9: e107005, 2014.
157. Jarvis RM, Hughes SM, and Ledgerwood EC. Peroxiredoxin 1 functions as a signal peroxidase to receive, transduce, and transmit peroxide signals in mammalian cells. *Free Radic Biol Med* 53: 1522–1530, 2012.
158. Jeong W, Cha MK, and Kim IH. Thioredoxin-dependent hydroperoxide peroxidase activity of bacterioferritin comigratory protein (BCP) as a new member of the thiol-specific antioxidant protein (TSA)/alkyl hydroperoxide peroxidase C (AhpC) family. *J Biol Chem* 275: 2924–2930, 2000.
159. Jin X, Stamnaes J, Klock C, DiRaimondo TR, Sollid LM, and Khosla C. Activation of extracellular transglutaminase 2 by thioredoxin. *J Biol Chem* 286: 37866–37873, 2011.
160. Jin ZG, Melaragno MG, Liao DF, Yan C, Haendeler J, Suh YA, Lambeth JD, and Berk BC. Cyclophilin A is a secreted growth factor induced by oxidative stress. *Circ Res* 87: 789–796, 2000.
161. Jo I, Chung IY, Bae HW, Kim JS, Song S, Cho YH, and Ha NC. Structural details of the OxyR peroxide-sensing mechanism. *Proc Natl Acad Sci U S A* 112: 6443–6448, 2015.
162. Johnson JR, Russo TA, Drawz SM, Clabots C, Olson R, Kuskowski MA, and Rosen H. OxyR contributes to the virulence of a Clonal Group A *Escherichia coli* strain (O17:K+:H18) in animal models of urinary tract infection, subcutaneous infection, and systemic sepsis. *Microb Pathog* 64: 1–5, 2013.
163. Jones DP and Sies H. The redox code. *Antioxid Redox Signal* 23: 734–746, 2015.
164. Jonic S. Cryo-electron microscopy analysis of structurally heterogeneous macromolecular complexes. *Comput Struct Biotechnol J* 14: 385–390, 2016.
165. Jonic S. Computational methods for analyzing conformational variability of macromolecular complexes from cryo-electron microscopy images. *Curr Opin Struct Biol* 43: 114–121, 2017.
166. Junn E, Taniguchi H, Jeong BS, Zhao X, Ichijo H, and Mouradian MM. Interaction of DJ-1 with Daxx inhibits apoptosis signal-regulating kinase 1 activity and cell death. *Proc Natl Acad Sci U S A* 102: 9691–9696, 2005.
167. Kamariah N, Eisenhaber B, Eisenhaber F, and Gruber G. Active site CP-loop dynamics modulate substrate binding,

- catalysis, oligomerization, stability, over-oxidation and recycling of 2-Cys peroxiredoxins. *Free Radic Biol Med* 118: 59–70, 2018.
168. Kamariah N, Sek MF, Eisenhaber B, Eisenhaber F, and Gruber G. Transition steps in peroxide reduction and a molecular switch for peroxide robustness of prokaryotic peroxiredoxins. *Sci Rep* 6: 37610, 2016.
 169. Kang DH, Lee DJ, Lee S, Lee SY, Jun Y, Kim Y, Kim Y, Lee JS, Lee DK, Lee S, Jho EH, Yu DY, and Kang SW. Interaction of tankyrase and peroxiredoxin II is indispensable for the survival of colorectal cancer cells. *Nat Commun* 8: 40, 2017.
 170. Kato I, Maita H, Takahashi-Niki K, Saito Y, Noguchi N, Iguchi-Ariga SM, and Ariga H. Oxidized DJ-1 inhibits p53 by sequestering p53 from promoters in a DNA-binding affinity-dependent manner. *Mol Cell Biol* 33: 340–359, 2013.
 171. Kern R, Malki A, Holmgren A, and Richarme G. Chaperone properties of *Escherichia coli* thioredoxin and thioredoxin reductase. *Biochem J* 371: 965–972, 2003.
 172. Kil IS, Lee SK, Ryu KW, Woo HA, Hu MC, Bae SH, and Rhee SG. Feedback control of adrenal steroidogenesis via H₂O₂-dependent, reversible inactivation of peroxiredoxin III in mitochondria. *Mol Cell* 46: 584–594, 2012.
 173. Kim DI, Jensen SC, Noble KA, Kc B, Roux KH, Motamedchaboki K, and Roux KJ. An improved smaller biotin ligase for BioID proximity labeling. *Mol Biol Cell* 27: 1188–1196, 2016.
 174. Kim H, Oh Y, Kim K, Jeong S, Chon S, Kim D, Jung MH, Pak YK, Ha J, Kang I, and Choe W. Cyclophilin A regulates JNK/p38-MAPK signaling through its physical interaction with ASK1. *Biochem Biophys Res Commun* 464: 112–117, 2015.
 175. Kim J-H, Choi D-J, Jeong H-K, Kim J, Kim DW, Choi SY, Park S-M, Suh YH, Jou I, and Joe E-H. DJ-1 facilitates the interaction between STAT1 and its phosphatase, SHP-1, in brain microglia and astrocytes: a novel anti-inflammatory function of DJ-1. *Neurobiol Dis* 60: 1–10, 2013.
 176. Kim JS and Holmes RK. Characterization of OxyR as a negative transcriptional regulator that represses catalase production in *Corynebacterium diphtheriae*. *PLoS One* 7: e31709, 2012.
 177. Kim RH, Peters M, Jang YJ, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L, Snow B, Binari RC, Manoukian AS, Bray MR, Liu FF, Tsao MS, and Mak TW. DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* 7: 263–273, 2005.
 178. Kim S, Bang YJ, Kim D, Lim JG, Oh MH, and Choi SH. Distinct characteristics of OxyR2, a new OxyR-type regulator, ensuring expression of peroxiredoxin 2 detoxifying low levels of hydrogen peroxide in *Vibrio vulnificus*. *Mol Microbiol* 93: 992–1009, 2014.
 179. Kim SY, Jang HH, Lee JR, Sung NR, Lee HB, Lee DH, Park DJ, Kang CH, Chung WS, Lim CO, Yun DJ, Kim WY, Lee KO, and Lee SY. Oligomerization and chaperone activity of a plant 2-Cys peroxiredoxin in response to oxidative stress. *Plant Sci* 177: 227–232, 2009.
 180. Kim SY, Kim TJ, and Lee KY. A novel function of peroxiredoxin 1 (Prx-1) in apoptosis signal-regulating kinase 1 (ASK1)-mediated signaling pathway. *FEBS Lett* 582: 1913–1918, 2008.
 181. Kinumi T, Kimata J, Taira T, Ariga H, and Niki E. Cysteine-106 of DJ-1 is the most sensitive cysteine residue due to hydrogen peroxide-mediated oxidation in vivo in human umbilical vein endothelial cells. *Biochem Biophys Res Commun* 317: 722–728, 2004.
 182. Kiraly R, Csosz E, Kurtan T, Antus S, Szigeti K, Simon-Vecsei Z, Korponay-Szabo IR, Keresztessy Z, and Fesus L. Functional significance of five noncanonical Ca²⁺-binding sites of human transglutaminase 2 characterized by site-directed mutagenesis. *FEBS J* 276: 7083–7096, 2009.
 183. Kiraly R, Demeny M, and Fesus L. Protein transamidation by transglutaminase 2 in cells: a disputed Ca²⁺-dependent action of a multifunctional protein. *FEBS J* 278: 4717–4739, 2011.
 184. König J, Galliard H, Jutte P, Schaper S, Dittmann L, and Dietz KJ. The conformational bases for the two functionalities of 2-cysteine peroxiredoxins as peroxidase and chaperone. *J Exp Bot* 64: 3483–3497, 2013.
 185. Kosek D, Kylarova S, Psenakova K, Rezabkova L, Herman P, Vecer J, Obsilova V, and Obsil T. Biophysical and structural characterization of the thioredoxin-binding domain of protein kinase ASK1 and its interaction with reduced thioredoxin. *J Biol Chem* 289: 24463–24474, 2014.
 186. Kumarevel T, Tanaka T, Umehara T, and Yokoyama S. ST1710-DNA complex crystal structure reveals the DNA binding mechanism of the MarR family of regulators. *Nucleic Acids Res* 37: 4723–4735, 2009.
 187. Kylarova S, Kosek D, Petrvalska O, Psenakova K, Man P, Vecer J, Herman P, Obsilova V, and Obsil T. Cysteine residues mediate high-affinity binding of thioredoxin to ASK1. *FEBS J* 283: 3821–3838, 2016.
 188. Lakshminarasimhan M, Madzelan P, Nan R, Milkovic NM, and Wilson MA. Evolution of new enzymatic function by structural modulation of cysteine reactivity in *Pseudomonas fluorescens* isocyanide hydratase. *J Biol Chem* 285: 29651–29661, 2010.
 189. Landino LM, Hagedorn TD, and Kennett KL. Evidence for thiol/disulfide exchange reactions between tubulin and glyceraldehyde-3-phosphate dehydrogenase. *Cytoskeleton* 71: 707–718, 2014.
 190. Latimer HR and Veal EA. Peroxiredoxins in regulation of MAPK signalling pathways; sensors and barriers to signal transduction. *Mol Cells* 39: 40–45, 2016.
 191. Laxa M, König J, Dietz KJ, and Kandlbinder A. Role of the cysteine residues in *Arabidopsis thaliana* cyclophilin CYP20-3 in peptidyl-prolyl cis-trans isomerase and redox-related functions. *Biochem J* 401: 287–297, 2007.
 192. Le HT, Gautier V, Kthiri F, Malki A, Messaoudi N, Mihoub M, Landoulsi A, An YJ, Cha SS, and Richarme G. YajL, prokaryotic homolog of parkinsonism-associated protein DJ-1, functions as a covalent chaperone for thiol proteome. *J Biol Chem* 287: 5861–5870, 2012.
 193. Lebedev I, Nemaierova A, Foda ZH, Kornaj M, Tong M, Moll UM, and Seeliger MA. A novel in vitro CypD-mediated p53 aggregation assay suggests a model for mitochondrial permeability transition by chaperone systems. *J Mol Biol* 428: 4154–4167, 2016.
 194. Lee C, Lee SM, Mukhopadhyay P, Kim SJ, Lee SC, Ahn WS, Yu MH, Storz G, and Ryu SE. Redox regulation of OxyR requires specific disulfide bond formation involving a rapid kinetic reaction path. *Nat Struct Mol Biol* 11: 1179–1185, 2004.
 195. Lee C, Shin J, and Park C. Novel regulatory system nemRA-gloA for electrophile reduction in *Escherichia coli* K-12. *Mol Microbiol* 88: 395–412, 2013.

196. Lee HJ, Howell SK, Sanford RJ, and Beisswenger PJ. Methylglyoxal can modify GAPDH activity and structure. *Ann N Y Acad Sci* 1043: 135–145, 2005.
197. Lee HT, Huang CH, Chen WC, Tsai CS, Chao YL, Liu SH, Chen JH, Wu YY, and Lee YJ. Transglutaminase 2 promotes migration and invasion of lung cancer cells. *Oncol Res* in press; doi: 10.3727/096504018X15149761920868, 2018.
198. Lee JR, Lee SS, Jang HH, Lee YM, Park JH, Park SC, Moon JC, Park SK, Kim SY, Lee SY, Chae HB, Jung YJ, Kim WY, Shin MR, Cheong GW, Kim MG, Kang KR, Lee KO, Yun DJ, and Lee SY. Heat-shock dependent oligomeric status alters the function of a plant-specific thioredoxin-like protein, AtTDX. *Proc Natl Acad Sci U S A* 106: 5978–5983, 2009.
199. Lee JW, Soonsanga S, and Helmann JD. A complex thiolate switch regulates the *Bacillus subtilis* organic peroxide sensor OhrR. *Proc Natl Acad Sci U S A* 104: 8743–8748, 2007.
200. Lee JY, Song J, Kwon K, Jang S, Kim C, Baek K, Kim J, and Park C. Human DJ-1 and its homologs are novel glyoxalases. *Hum Mol Genet* 21: 3215–3225, 2012.
201. Lee MK, Lee MS, Bae DW, Lee DH, Cha SS, and Chi SW. Structural basis for the interaction between DJ-1 and Bcl-X_L. *Biochem Biophys Res Commun* 495: 1067–1073, 2018.
202. Lee SM, Jeong EM, Jeong J, Shin DM, Lee HJ, Kim HJ, Lim J, Lee JH, Cho SY, Kim MK, Wee WR, Lee JH, and Kim IG. Cysteamine prevents the development of lens opacity in a rat model of selenite-induced cataract. *Invest Ophthalmol Vis Sci* 53: 1452–1459, 2012.
203. Lee SP, Hwang YS, Kim YJ, Kwon KS, Kim HJ, Kim K, and Chae HZ. Cyclophilin A binds to peroxiredoxins and activates its peroxidase activity. *J Biol Chem* 276: 29826–29832, 2001.
204. Lee SR, Yang KS, Kwon J, Lee C, Jeong W, and Rhee SG. Reversible inactivation of the tumor suppressor PTEN by H₂O₂. *J Biol Chem* 277: 20336–20342, 2002.
205. Lee YS, Lee J, Ryu KS, Lee Y, Jung TG, Jang JH, Sim DW, Kim EH, Seo MD, Lee KW, and Won HS. Semi-empirical structure determination of *Escherichia coli* Hsp33 and identification of dynamic regulatory elements for the activation process. *J Mol Biol* 427: 3850–3861, 2015.
206. Lemaire SD, Guillon B, Le Marechal P, Keryer E, Miginiac-Maslow M, and Decottignies P. New thioredoxin targets in the unicellular photosynthetic eukaryote *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci U S A* 101: 7475–7480, 2004.
207. Leonard SE, Reddie KG, and Carroll KS. Mining the thiol proteome for sulfenic acid modifications reveals new targets for oxidation in cells. *ACS Chem Biol* 4: 783–799, 2009.
208. Li L, Cheung SH, Evans EL, and Shaw PE. Modulation of gene expression and tumor cell growth by redox modification of STAT3. *Cancer Res* 70: 8222–8232, 2010.
209. Li L and Shaw PE. A STAT3 dimer formed by inter-chain disulphide bridging during oxidative stress. *Biochem Biophys Res Commun* 322: 1005–1011, 2004.
210. Li T, Guo H, Zhao X, Jin J, Zhang L, Li H, Lu Y, Nie Y, Wu K, Shi Y, and Fan D. Gastric cancer cell proliferation and survival is enabled by a cyclophilin B/STAT3/miR-520d-5p signaling feedback loop. *Cancer Res* 77: 1227–1240, 2017.
211. Lievens S, Gerlo S, Lemmens I, De Clercq DJ, Risseuw MD, Vanderroost N, De Smet AS, Ruyssink E, Chevet E, Van Calenbergh S, and Tavernier J. Kinase substrate sensor (KISS), a mammalian in situ protein interaction sensor. *Mol Cell Proteomics* 13: 3332–3342, 2014.
212. Limauro D, D'Ambrosio K, Langella E, De Simone G, Galdi I, Pedone C, Pedone E, and Bartolucci S. Exploring the catalytic mechanism of the first dimeric Bcp: functional, structural and docking analyses of Bcp4 from *Sulfolobus solfataricus*. *Biochimie* 92: 1435–1444, 2010.
213. Lin D, Saleh S, and Liebler DC. Reversibility of covalent electrophile-protein adducts and chemical toxicity. *Chem Res Toxicol* 21: 2361–2369, 2008.
214. Linard D, Kandlbinder A, Degand H, Morsomme P, Dietz KJ, and Knoops B. Redox characterization of human cyclophilin D: identification of a new mammalian mitochondrial redox sensor? *Arch Biochem Biophys* 491: 39–45, 2009.
215. Little C and O'Brien PJ. Mechanism of peroxide-inactivation of the sulphhydryl enzyme glyceraldehyde-3-phosphate dehydrogenase. *Eur J Biochem* 10: 533–538, 1969.
216. Littler DR, Harrop SJ, Fairlie WD, Brown LJ, Pankhurst GJ, Pankhurst S, DeMaere MZ, Campbell TJ, Bauskin AR, Tonini R, Mazzanti M, Breit SN, and Curmi PM. The intracellular chloride ion channel protein CLIC1 undergoes a redox-controlled structural transition. *J Biol Chem* 279: 9298–9305, 2004.
217. Liu B, Chen Y, and St Clair DK. ROS and p53: a versatile partnership. *Free Radic Biol Med* 44: 1529–1535, 2008.
218. Liu D and Xu Y. p53, Oxidative stress, and aging. *Antioxid Redox Signal* 15: 1669–1678, 2011.
219. Liu GJ, Liu X, Xu HJ, Liu XC, Zhou H, Huang Z, Gan JH, Chen H, Lan L, and Yang CG. Structural insights into the redox-sensing mechanism of MarR-type regulator AbfR. *J Am Chem Soc* 139: 1598–1608, 2017.
220. Liu X, Sun X, Wu Y, Xie C, Zhang W, Wang D, Chen X, Qu D, Gan J, Chen H, Jiang H, Lan L, and Yang CG. Oxidation-sensing regulator AbfR regulates oxidative stress responses, bacterial aggregation, and biofilm formation in *Staphylococcus epidermidis*. *J Biol Chem* 288: 3739–3752, 2013.
221. Liu Z, Gong Z, Dong X, and Tang C. Transient protein-protein interactions visualized by solution NMR. *Biochim Biophys Acta* 1864: 115–122, 2016.
222. Loi VV, Harms M, Muller M, Huyen NT, Hamilton CJ, Hochgrafe F, Pane-Farre J, and Antelmann H. Real-time imaging of the bacillithiol redox potential in the human pathogen *Staphylococcus aureus* using a genetically encoded bacilliredoxin-fused redox biosensor. *Antioxid Redox Signal* 26: 835–848, 2017.
223. Love NR, Chen Y, Ishibashi S, Kritsiligkou P, Lea R, Koh Y, Gallop JL, Dorey K, and Amaya E. Amputation-induced reactive oxygen species are required for successful *Xenopus tadpole* tail regeneration. *Nat Cell Biol* 15: 222–228, 2013.
224. Luciani A, Vilella VR, Vasaturo A, Giardino I, Pettoello-Mantovani M, Guido S, Cexus ON, Peake N, Londei M, Quarantino S, and Maiuri L. Lysosomal accumulation of gliadin p31-43 peptide induces oxidative stress and tissue transglutaminase-mediated PPARgamma downregulation in intestinal epithelial cells and coeliac mucosa. *Gut* 59: 311–319, 2010.
225. Luciani A, Vilella VR, Vasaturo A, Giardino I, Raia V, Pettoello-Mantovani M, D'Apolito M, Guido S, Leal T, Quarantino S, and Maiuri L. SUMOylation of tissue

- transglutaminase as link between oxidative stress and inflammation. *J Immunol* 183: 2775–2784, 2009.
226. Lundstrom J and Holmgren A. Determination of the reduction-oxidation potential of the thioredoxin-like domains of protein disulfide-isomerase from the equilibrium with glutathione and thioredoxin. *Biochemistry* 32: 6649–6655, 1993.
 227. Luo M, Zhang J, He H, Su D, Chen Q, Gross ML, Kelley MR, and Georgiadis MM. Characterization of the redox activity and disulfide bond formation in apurinic/aprimidinic endonuclease. *Biochemistry* 51: 695–705, 2012.
 228. Ma LH, Takanishi CL, and Wood MJ. Molecular mechanism of oxidative stress perception by the Orp1 protein. *J Biol Chem* 282: 31429–31436, 2007.
 229. Ma Z, Russo VC, Rabadi SM, Jen Y, Catlett SV, Bakshi CS, and Malik M. Elucidation of a mechanism of oxidative stress regulation in *Francisella tularensis* live vaccine strain. *Mol Microbiol* 101: 856–878, 2016.
 230. Malorni W, Farrace MG, Matarrese P, Tinari A, Ciarlo L, Mousavi-Shafaei P, D'Eletto M, Di Giacomo G, Melino G, Palmieri L, Rodolfo C, and Piacentini M. The adenine nucleotide translocator 1 acts as a type 2 transglutaminase substrate: implications for mitochondrial-dependent apoptosis. *Cell Death Differ* 16: 1480–1492, 2009.
 231. Markvicheva KN, Bilan DS, Mishina NM, Gorokhovatsky AY, Vinokurov LM, Lukyanov S, and Belousov VV. A genetically encoded sensor for H₂O₂ with expanded dynamic range. *Bioorg Med Chem* 19: 1079–1084, 2011.
 232. Mastroberardino PG, Farrace MG, Viti I, Pavone F, Fimia GM, Melino G, Rodolfo C, and Piacentini M. “Tissue” transglutaminase contributes to the formation of disulphide bridges in proteins of mitochondrial respiratory complexes. *Biochim Biophys Acta* 1757: 1357–1365, 2006.
 233. Matlashov ME, Bogdanova YA, Ermakova GV, Mishina NM, Ermakova YG, Nikitin ES, Balaban PM, Okabe S, Lukyanov S, Enikolopov G, Zaraisky AG, and Belousov VV. Fluorescent ratiometric pH indicator SypHer2: applications in neuroscience and regenerative biology. *Biochim Biophys Acta* 1850: 2318–2328, 2015.
 234. Meier JA, Hyun M, Cantwell M, Raza A, Mertens C, Raju V, Sisler J, Tracy E, Torres-Odio S, Gispert S, Shaw PE, Baumann H, Bandyopadhyay D, Takabe K, and Larner AC. Stress-induced dynamic regulation of mitochondrial STAT3 and its association with cyclophilin D reduce mitochondrial ROS production. *Sci Signal* 10: eaag2588, 2017.
 235. Meng TC and Tonks NK. Analysis of the regulation of protein tyrosine phosphatases in vivo by reversible oxidation. *Methods Enzymol* 366: 304–318, 2003.
 236. Merk A, Bartesaghi A, Banerjee S, Falconieri V, Rao P, Davis MI, Pragani R, Boxer MB, Earl LA, Milne JL, and Subramaniam S. Breaking Cryo-EM resolution barriers to facilitate drug discovery. *Cell* 165: 1698–1707, 2016.
 237. Meuillet EJ, Mahadevan D, Berggren M, Coon A, and Powis G. Thioredoxin-1 binds to the C2 domain of PTEN inhibiting PTEN's lipid phosphatase activity and membrane binding: a mechanism for the functional loss of PTEN's tumor suppressor activity. *Arch Biochem Biophys* 429: 123–133, 2004.
 238. Miller EW, Albers AE, Pralle A, Isacoff EY, and Chang CJ. Boronate-based fluorescent probes for imaging cellular hydrogen peroxide. *J Am Chem Soc* 127: 16652–16659, 2005.
 239. Mishra S and Murphy LJ. Tissue transglutaminase has intrinsic kinase activity: identification of transglutaminase 2 as an insulin-like growth factor-binding protein-3 kinase. *J Biol Chem* 279: 23863–23868, 2004.
 240. Mishra S and Murphy LJ. The p53 oncoprotein is a substrate for tissue transglutaminase kinase activity. *Biochem Biophys Res Commun* 339: 726–730, 2006.
 241. Mishra S, Saleh A, Espino PS, Davie JR, and Murphy LJ. Phosphorylation of histones by tissue transglutaminase. *J Biol Chem* 281: 5532–5538, 2006.
 242. Mitsumoto A and Nakagawa Y. DJ-1 is an indicator for endogenous reactive oxygen species elicited by endotoxin. *Free Radic Res* 35: 885–893, 2001.
 243. Mitsumoto A, Nakagawa Y, Takeuchi A, Okawa K, Iwamatsu A, and Takanezawa Y. Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radic Res* 35: 301–310, 2001.
 244. Mo JS, Jung J, Yoon JH, Hong JA, Kim MY, Ann EJ, Seo MS, Choi YH, and Park HS. DJ-1 modulates the p38 mitogen-activated protein kinase pathway through physical interaction with apoptosis signal-regulating kinase 1. *J Cell Biochem* 110: 229–237, 2010.
 245. Morais MA, Giuseppe PO, Souza TA, Alegria TG, Oliveira MA, Netto LE, and Murakami MT. How pH modulates the dimer-decamer interconversion of 2-Cys peroxiredoxins from the Prx1 subfamily. *J Biol Chem* 290: 8582–8590, 2015.
 246. Morgan B, Van Laer K, Owusu TN, Ezerina D, Pastor-Flores D, Amponsah PS, Tursch A, and Dick TP. Real-time monitoring of basal H₂O₂ levels with peroxiredoxin-based probes. *Nat Chem Biol* 12: 437–443, 2016.
 247. Morinaka A, Funato Y, Uesugi K, and Miki H. Oligomeric peroxiredoxin-I is an essential intermediate for p53 to activate MST1 kinase and apoptosis. *Oncogene* 30: 4208–4218, 2011.
 248. Motohashi K, Koyama F, Nakanishi Y, Ueoka-Nakanishi H, and Hisabori T. Chloroplast cyclophilin is a target protein of thioredoxin. Thiol modulation of the peptidyl-prolyl cis-trans isomerase activity. *J Biol Chem* 278: 31848–31852, 2003.
 249. Murray JW, Garman EF, and Ravelli RBG. X-ray absorption by macromolecular crystals: the effects of wavelength and crystal composition on absorbed dose. *J Appl Crystallogr* 37: 513–522, 2004.
 250. Muthuramalingam M, Matros A, Scheibe R, Mock HP, and Dietz KJ. The hydrogen peroxide-sensitive proteome of the chloroplast in vitro and in vivo. *Front Plant Sci* 4: 54, 2013.
 251. Muthuramalingam M, Seidel T, Laxa M, Nunes de Miranda SM, Gartner F, Stroher E, Kandlbinder A, and Dietz KJ. Multiple redox and non-redox interactions define 2-Cys peroxiredoxin as a regulatory hub in the chloroplast. *Mol Plant* 2: 1273–1288, 2009.
 252. Nadeau PJ, Charette SJ, and Landry J. REDOX reaction at ASK1-Cys250 is essential for activation of JNK and induction of apoptosis. *Mol Biol Cell* 20: 3628–3637, 2009.
 253. Nadeau PJ, Charette SJ, Toledano MB, and Landry J. Disulfide bond-mediated multimerization of ASK1 and its reduction by thioredoxin-1 regulate H₂O₂-induced c-Jun NH₂-terminal kinase activation and apoptosis. *Mol Biol Cell* 18: 3903–3913, 2007.
 254. Nakao LS, Everley RA, Marino SM, Lo SM, de Souza LE, Gygi SP, and Gladyshev VN. Mechanism-based proteomic

- screening identifies targets of thioredoxin-like proteins. *J Biol Chem* 290: 5685–5695, 2015.
255. Nakaoka H, Perez DM, Baek KJ, Das T, Husain A, Misono K, Im MJ, and Graham RM. Gh: a GTP-binding protein with transglutaminase activity and receptor signaling function. *Science* 264: 1593–1596, 1994.
 256. Nassour H, Wang Z, Saad A, Papaluca A, Brosseau N, Affar el B, Alaoui-Jamali MA, and Ramotar D. Peroxiredoxin 1 interacts with and blocks the redox factor APE1 from activating interleukin-8 expression. *Sci Rep* 6: 29389, 2016.
 257. Naugler WE and Karin M. NF-kappaB and cancer-identifying targets and mechanisms. *Curr Opin Genet Dev* 18: 19–26, 2008.
 258. Newberry KJ, Fuangthong M, Panmanee W, Mongkolsuk S, and Brennan RG. Structural mechanism of organic hydroperoxide induction of the transcription regulator OhrR. *Mol Cell* 28: 652–664, 2007.
 259. Nguyen TT, Stevens MV, Kohr M, Steenbergen C, Sack MN, and Murphy E. Cysteine 203 of cyclophilin D is critical for cyclophilin D activation of the mitochondrial permeability transition pore. *J Biol Chem* 286: 40184–40192, 2011.
 260. Nielsen MH, Kidmose RT, and Jenner LB. Structure of TSA2 reveals novel features of the active-site loop of peroxiredoxins. *ACTA Crystallogr D Struct Biol* 72: 158–167, 2016.
 261. Niethammer P, Grabher C, Look AT, and Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 459: 996–999, 2009.
 262. Niki T, Takahashi-Niki K, Taira T, Iguchi-Ariga SM, and Ariga H. DJBP: a novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol Cancer Res* 1: 247–261, 2003.
 263. Noichiri Y, Palais G, Ruby V, D'Autreaux B, Delaunay-Moisan A, Nystrom T, Molin M, and Toledano MB. In vivo parameters influencing 2-Cys Prx oligomerization: the role of enzyme sulfinylation. *Redox Biol* 6: 326–333, 2015.
 264. Oger E, Marino D, Guigonis JM, Pauly N, and Puppo A. Sulfenylated proteins in the *Medicago truncatula*-*Sinorhizobium meliloti* symbiosis. *J Proteomics* 75: 4102–4113, 2012.
 265. Okazaki S, Naganuma A, and Kuge S. Peroxiredoxin-mediated redox regulation of the nuclear localization of Yap1, a transcription factor in budding yeast. *Antioxid Redox Signal* 7: 327–334, 2005.
 266. Okazaki S, Tachibana T, Naganuma A, Mano N, and Kuge S. Multistep disulfide bond formation in Yap1 is required for sensing and transduction of H₂O₂ stress signal. *Mol Cell* 27: 675–688, 2007.
 267. Ostman A, Frijhoff J, Sandin A, and Bohmer FD. Regulation of protein tyrosine phosphatases by reversible oxidation. *J Biochem* 150: 345–356, 2011.
 268. Owuor ED and Kong AN. Antioxidants and oxidants regulated signal transduction pathways. *Biochem Pharmacol* 64: 765–770, 2002.
 269. Pan J and Carroll KS. Chemical biology approaches to study protein cysteine sulfinylation. *Biopolymers* 101: 165–172, 2014.
 270. Panmanee W, Vattanaviboon P, Poole LB, and Mongkolsuk S. Novel organic hydroperoxide-sensing and responding mechanisms for OhrR, a major bacterial sensor and regulator of organic hydroperoxide stress. *J Bacteriol* 188: 1389–1395, 2006.
 271. Park J, Lee S, Lee S, and Kang SW. 2-Cys peroxiredoxins: emerging hubs determining redox dependency of mammalian signaling networks. *Int J Cell Biol* 2014: 715867, 2014.
 272. Park JE, Lee JA, Park SG, Lee DH, Kim SJ, Kim HJ, Uchida C, Uchida T, Park BC, and Cho S. A critical step for JNK activation: isomerization by the prolyl isomerase Pin1. *Cell Death Differ* 19: 153–161, 2012.
 273. Park SK, Jung YJ, Lee JR, Lee YM, Jang HH, Lee SS, Park JH, Kim SY, Moon JC, Lee SY, Chae HB, Shin MR, Jung JH, Kim MG, Kim WY, Yun DJ, Lee KO, and Lee SY. Heat-shock and redox-dependent functional switching of an h-type *Arabidopsis* thioredoxin from a disulfide reductase to a molecular chaperone. *Plant Physiol* 150: 552–561, 2009.
 274. Parsonage D, Nelson KJ, Ferrer-Sueta G, Alley S, Karplus PA, Furdul CM, and Poole LB. Dissecting peroxiredoxin catalysis: separating binding, peroxidation, and resolution for a bacterial AhpC. *Biochemistry* 54: 1567–1575, 2015.
 275. Parsonage D, Youngblood DS, Sarma GN, Wood ZA, Karplus PA, and Poole LB. Analysis of the link between enzymatic activity and oligomeric state in AhpC, a bacterial peroxiredoxin. *Biochemistry* 44: 10583–10592, 2005.
 276. Paulsen CE, Truong TH, Garcia FJ, Homann A, Gupta V, Leonard SE, and Carroll KS. Peroxide-dependent sulfinylation of the EGFR catalytic site enhances kinase activity. *Nat Chem Biol* 8: 57–64, 2011.
 277. Peralta D, Bronowska AK, Morgan B, Doka E, Van Laer K, Nagy P, Grater F, and Dick TP. A proton relay enhances H₂O₂ sensitivity of GAPDH to facilitate metabolic adaptation. *Nat Chem Biol* 11: 156–163, 2015.
 278. Perez-Perez ME, Mauries A, Maes A, Tourasse NJ, Hamon M, Lemaire SD, and Marchand CH. The deep thioredoxome in *Chlamydomonas reinhardtii*: new insights into redox regulation. *Mol Plant* 10: 1107–1125, 2017.
 279. Perkins A, Nelson KJ, Williams JR, Parsonage D, Poole LB, and Karplus PA. The sensitive balance between the fully folded and locally unfolded conformations of a model peroxiredoxin. *Biochemistry* 52: 8708–8721, 2013.
 280. Perkins A, Parsonage D, Nelson KJ, Ogbia OM, Cheong PH, Poole LB, and Karplus PA. Peroxiredoxin catalysis at atomic resolution. *Structure* 24: 1668–1678, 2016.
 281. Peskin AV, Dickerhof N, Poynton RA, Paton LN, Pace PE, Hampton MB, and Winterbourn CC. Hyperoxidation of peroxiredoxins 2 and 3: rate constants for the reactions of the sulfenic acid of the peroxidatic cysteine. *J Biol Chem* 288: 14170–14177, 2013.
 282. Piacentini M, Farrace MG, Piredda L, Matarrese P, Ciccosanti F, Falasca L, Rodolfo C, Giammarioli AM, Verderio E, Griffin M, and Malorni W. Transglutaminase overexpression sensitizes neuronal cell lines to apoptosis by increasing mitochondrial membrane potential and cellular oxidative stress. *J Neurochem* 81: 1061–1072, 2002.
 283. Pineyro MD, Parodi-Talice A, Portela M, Arias DG, Guerrero SA, and Robello C. Molecular characterization and interactome analysis of *Trypanosoma cruzi* trypanoxin 1. *J Proteomics* 74: 1683–1692, 2011.
 284. Pinkas DM, Strop P, Brunger AT, and Khosla C. Transglutaminase 2 undergoes a large conformational change upon activation. *PLoS Biol* 5: e327, 2007.

285. Piston D, Alvarez-Erviti L, Bansal V, Gargano D, Yao Z, Szabadkai G, Odell M, Puno MR, Bjorkblom B, Maple-Groden J, Breuer P, Kaut O, Larsen JP, Bonn S, Moller SG, Wullner U, Schapira AHV, and Gegg ME. DJ-1 is a redox sensitive adapter protein for high molecular weight complexes involved in regulation of catecholamine homeostasis. *Hum Mol Genet* 26: 4028–4041, 2017.
286. Prasad BD, Goel S, and Krishna P. In silico identification of carboxylate clamp type tetratricopeptide repeat proteins in *Arabidopsis* and rice as putative co-chaperones of Hsp90/Hsp70. *PLoS One* 5: e12761, 2010.
287. Putker M, Vos HR, van Dorenmalen K, de Ruiter H, Duran AG, Snel B, Burgering BM, Vermeulen M, and Dansen TB. Evolutionary acquisition of cysteines determines FOXO paralogue-specific redox signaling. *Antioxid Redox Signal* 22: 15–28, 2015.
288. Qian J, Wani R, Klomsiri C, Poole LB, Tsang AW, and Furdul CM. A simple and effective strategy for labeling cysteine sulfenic acid in proteins by utilization of beta-ketoesters as cleavable probes. *Chem Commun* 48: 4091–4093, 2012.
289. Qin LS, Jia PF, Zhang ZQ, and Zhang SM. ROS-p53-cyclophilin-D signaling mediates salinomycin-induced glioma cell necrosis. *J Exp Clin Cancer Res* 34: 57, 2015.
290. Qu D, Rashidian J, Mount MP, Aleyasin H, Parsanejad M, Lira A, Haque E, Zhang Y, Callaghan S, Daigle M, Rousseaux MWC, Slack RS, Albert PR, Vincent I, Woulfe JM, and Park DS. Role of Cdk5-mediated phosphorylation of Prx2 in MPTP toxicity and Parkinson's disease. *Neuron* 55: 37–52, 2007.
291. Qu Z, Meng F, Bomgardner RD, Viner RI, Li J, Rogers JC, Cheng J, Greenlief CM, Cui J, Lubahn DB, Sun GY, and Gu Z. Proteomic quantification and site-mapping of S-nitrosylated proteins using isobaric iodoTMT reagents. *J Proteome Res* 13: 3200–3211, 2014.
292. Radak Z, Koltai E, Taylor AW, Higuchi M, Kumagai S, Ohno H, Goto S, and Boldogh I. Redox-regulating sirtuins in aging, caloric restriction, and exercise. *Free Radic Biol Med* 58: 87–97, 2013.
293. Radjainia M, Venugopal H, Desfosses A, Phillips AJ, Yewdall NA, Hampton MB, Gerrard JA, and Mitra AK. Cryo-electron microscopy structure of human peroxiredoxin-3 filament reveals the assembly of a putative chaperone. *Structure* 23: 912–920, 2015.
294. Ragu S, Dardalhon M, Sharma S, Iraqui I, Buhagiar-Labarchede G, Grondin V, Kienda G, Vernis L, Chanet R, Kolodner RD, Huang ME, and Faye G. Loss of the thioredoxin reductase Trx1 suppresses the genomic instability of peroxiredoxin tsal mutants. *PLoS One* 9: e108123, 2014.
295. Ralser M, Wamelink MM, Kowald A, Gerisch B, Heeren G, Struys EA, Klipp E, Jakobs C, Breitenbach M, Lehrach H, and Krobitsch S. Dynamic rerouting of the carbohydrate flux is key to counteracting oxidative stress. *J Biol* 6: 10, 2007.
296. Randall LM, Ferrer-Sueta G, and Denicola A. Peroxiredoxins as preferential targets in H₂O₂-induced signaling. *Methods Enzymol* 527: 41–63, 2013.
297. Ravelli RB and McSweeney SM. The “fingerprint” that X-rays can leave on structures. *Structure* 8: 315–328, 2000.
298. Rawat SJ, Creasy CL, Peterson JR, and Chernoff J. The tumor suppressor Mst1 promotes changes in the cellular redox state by phosphorylation and inactivation of peroxiredoxin-1 protein. *J Biol Chem* 288: 8762–8771, 2013.
299. Rawson S, Davies S, Lippiat JD, and Muench SP. The changing landscape of membrane protein structural biology through developments in electron microscopy. *Mol Membr Biol* 33: 12–22, 2016.
300. Reddie KG, Seo YH, Muse WB, III, Leonard SE, and Carroll KS. A chemical approach for detecting sulfenic acid-modified proteins in living cells. *Mol Biosyst* 4: 521–531, 2008.
301. Reisz JA, Bechtold E, King SB, Poole LB, and Furdul CM. Thiol-blocking electrophiles interfere with labeling and detection of protein sulfenic acids. *FEBS J* 280: 6150–6161, 2013.
302. Ren H, Fu K, Wang D, Mu C, and Wang G. Oxidized DJ-1 interacts with the mitochondrial protein BCL-X_L. *J Biol Chem* 286: 35308–35317, 2011.
303. Rhee SG. Cell signaling. H₂O₂, a necessary evil for cell signaling. *Science* 312: 1882–1883, 2006.
304. Richarme G, Mihoub M, Dairou J, Bui LC, Leger T, and Lamouri A. Parkinsonism-associated protein DJ-1/Park7 is a major protein deglycase that repairs methylglyoxal- and glyoxal-glycated cysteine, arginine, and lysine residues. *J Biol Chem* 290: 1885–1897, 2015.
305. Riddell JR, Wang XY, Minderman H, and Gollnick SO. Peroxiredoxin 1 stimulates secretion of proinflammatory cytokines by binding to TLR4. *J Immunol* 184: 1022–1030, 2010.
306. Rigoulet M, Yoboue ED, and Devin A. Mitochondrial ROS generation and its regulation: mechanisms involved in H(2)O(2) signaling. *Antioxid Redox Signal* 14: 459–468, 2011.
307. Rodacka A. The effect of radiation-induced reactive oxygen species (ROS) on the structural and functional properties of yeast alcohol dehydrogenase (YADH). *Int J Radiat Biol* 92: 11–23, 2016.
308. Romano PG, Horton P, and Gray JE. The *Arabidopsis* cyclophilin gene family. *Plant Physiol* 134: 1268–1282, 2004.
309. Roos G, Fonseca Guerra C, and Bickelhaupt FM. How the disulfide conformation determines the disulfide/thiol redox potential. *J Biomol Struct Dyn* 33: 93–103, 2015.
310. Rosado LA, Wahni K, Degiacomi G, Pedre B, Young D, de la Rubia AG, Boldrin F, Martens E, Marcos-Pascual L, Sancho-Vaello E, Albasa-Jove D, Provvedi R, Martin C, Makarov V, Versees W, Verniest G, Guerin ME, Mateos LM, Manganelli R, and Messens J. The antibacterial prodrug activator Rv2466c is a mycothiol-dependent reductase in the oxidative stress response of *Mycobacterium tuberculosis*. *J Biol Chem* 292: 13097–13110, 2017.
311. Rossin F, D'Eletto M, Falasca L, Sepe S, Cocco S, Fimia GM, Campanella M, Mastroberardino PG, Farrace MG, and Piacentini M. Transglutaminase 2 ablation leads to mitophagy impairment associated with a metabolic shift towards aerobic glycolysis. *Cell Death Differ* 22: 408–418, 2015.
312. Roux KJ, Kim DI, and Burke B. BioID: a screen for protein-protein interactions. *Curr Protoc Protein Sci* 74: Unit 19.23, 2013.
313. Roux KJ, Kim DI, Raida M, and Burke B. A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells. *J Cell Biol* 196: 801–810, 2012.
314. Saccoccia F, Di Micco P, Boumis G, Brunori M, Koutris I, Miele AE, Morea V, Sriratana P, Williams DL, Bellelli A, and Angelucci F. Moonlighting by different stressors: crystal structure of the chaperone species of a 2-Cys peroxiredoxin. *Structure* 20: 429–439, 2012.

315. Sainsbury S, Ren J, Nettleship JE, Saunders NJ, Stuart DI, and Owens RJ. The structure of a reduced form of OxyR from *Neisseria meningitidis*. *BMC Struct Biol* 10: 10, 2010.
316. Saitoh M, Nishitoh H, Fujii M, Takeda K, Tobiume K, Sawada Y, Kawabata M, Miyazono K, and Ichijo H. Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1. *EMBO J* 17: 2596–2606, 1998.
317. Sajjad MU, Green EW, Miller-Fleming L, Hands S, Herrera F, Campesan S, Khoshnan A, Outeiro TF, Giorgini F, and Wytenbach A. DJ-1 modulates aggregation and pathogenesis in models of Huntington's disease. *Hum Mol Genet* 23: 755–766, 2014.
318. Samson AL, Knaupp AS, Kass I, Kleifeld O, Marijanovic EM, Hughes VA, Lupton CJ, Buckle AM, Bottomley SP, and Medcalf RL. Oxidation of an exposed methionine instigates the aggregation of glyceraldehyde-3-phosphate dehydrogenase. *J Biol Chem* 289: 26922–26936, 2014.
319. Sanz-Barrio R, Fernandez-San Millan A, Carballeda J, Corral-Martinez P, Segui-Simarro JM, and Farran I. Chaperone-like properties of tobacco plastid thioredoxins f and m. *J Exp Bot* 63: 365–379, 2012.
320. Sara T, Schwarz TC, Kurzbach D, Wunderlich CH, Kreutz C, and Konrat R. Magnetic resonance access to transiently formed protein complexes. *ChemistryOpen* 3: 115–123, 2014.
321. Saurin AT, Neubert H, Brennan JP, and Eaton P. Widespread sulfenic acid formation in tissues in response to hydrogen peroxide. *Proc Natl Acad Sci U S A* 101: 17982–17987, 2004.
322. Sayed AA and Williams DL. Biochemical characterization of 2-Cys peroxiredoxins from *Schistosoma mansoni*. *J Biol Chem* 279: 26159–26166, 2004.
323. Sayin VI, Ibrahim MX, Larsson E, Nilsson JA, Lindahl P, and Bergo MO. Antioxidants accelerate lung cancer progression in mice. *Sci Transl Med* 6: 221ra15, 2014.
324. Schinzel AC, Takeuchi O, Huang Z, Fisher JK, Zhou Z, Rubens J, Hetz C, Danial NN, Moskowitz MA, and Korsmeyer SJ. Cyclophilin D is a component of mitochondrial permeability transition and mediates neuronal cell death after focal cerebral ischemia. *Proc Natl Acad Sci U S A* 102: 12005–12010, 2005.
325. Schmidt B, Ho L, and Hogg PJ. Allosteric disulfide bonds. *Biochemistry* 45: 7429–7433, 2006.
326. Schopp IM, Amaya Ramirez CC, Debeljak J, Kreibich E, Skribbe M, Wild K, and Béthune J. Split-BioID a conditional proteomics approach to monitor the composition of spatiotemporally defined protein complexes. *Nat Commun* 8: 15690, 2017.
327. Schröder E, Littlechil JA, Lebedev AA, Errington N, Vagin AA, Isupov MN. Crystal structure of decameric 2-Cys peroxiredoxin from human erythrocytes at 1.7 Å resolution. *Structure* 8: 605–615, 2000.
328. Schwarzlander M, Fricker MD, Muller C, Marty L, Brach T, Novak J, Sweetlove LJ, Hell R, and Meyer AJ. Confocal imaging of glutathione redox potential in living plant cells. *J Microsc* 231: 299–316, 2008.
329. Scorrano L, Nicoli A, Basso E, Petronilli V, and Bernardi P. Two modes of activation of the permeability transition pore: the role of mitochondrial cyclophilin. *Mol Cell Biochem* 174: 181–184, 1997.
330. Seib KL, Wu HJ, Srihanta YN, Edwards JL, Falsetta ML, Hamilton AJ, Maguire TL, Grimmond SM, Apicella MA, McEwan AG, and Jennings MP. Characterization of the OxyR regulon of *Neisseria gonorrhoeae*. *Mol Microbiol* 63: 54–68, 2007.
331. Seo YH and Carroll KS. Facile synthesis and biological evaluation of a cell-permeable probe to detect redox-regulated proteins. *Bioorg Med Chem Lett* 19: 356–359, 2009.
332. Sharma D and Rajarathnam K. ¹³C NMR chemical shifts can predict disulfide bond formation. *J Biomol NMR* 18: 165–171, 2000.
333. Shaw PE. Could STAT3 provide a link between respiration and cell cycle progression? *Cell Cycle* 9: 4294–4296, 2010.
334. Shendelman S, Jonason A, Martinat C, Leete T, and Abeliovich A. DJ-1 is a redox-dependent molecular chaperone that inhibits alpha-synuclein aggregate formation. *PLoS Biol* 2: e362, 2004.
335. Shinbo Y, Taira T, Niki T, Iguchi-Ariga SM, and Ariga H. DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3. *Int J Oncol* 26: 641–648, 2005.
336. Shirmanova MV, Druzhkova IN, Lukina MM, Matlashov ME, Belousov VV, Snopova LB, Prodanetz NN, Dudenkova VV, Lukyanov SA, and Zagaynova EV. Intracellular pH imaging in cancer cells in vitro and tumors in vivo using the new genetically encoded sensor Sy-pHer2. *Biochim Biophys Acta* 1850: 1905–1911, 2015.
337. Sies H. Biochemistry of oxidative stress. *Angew Chem Int Ed Engl* 25: 1058–1071, 1986.
338. Sies H. Role of metabolic H₂O₂ generation: redox signaling and oxidative stress. *J Biol Chem* 289: 8735–8741, 2014.
339. Sies H. Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: oxidative eustress. *Redox Biol* 11: 613–619, 2017.
340. Sies H, Berndt C, and Jones DP. Oxidative stress. *Annu Rev Biochem* 86: 715–748, 2017.
341. Singh H and Ashley RH. CLIC4 (p64H1) and its putative transmembrane domain form poorly selective, redox-regulated ion channels. *Mol Membr Biol* 24: 41–52, 2007.
342. Sirover MA. New insights into an old protein: the functional diversity of mammalian glyceraldehyde-3-phosphate dehydrogenase. *Biochim Biophys Acta* 1432: 159–184, 1999.
343. Sirover MA. On the functional diversity of glyceraldehyde-3-phosphate dehydrogenase: biochemical mechanisms and regulatory control. *Biochim Biophys Acta* 1810: 741–751, 2011.
344. Smith JK, Patil CN, Patlolla S, Gunter BW, Booz GW, and Duhe RJ. Identification of a redox-sensitive switch within the JAK2 catalytic domain. *Free Radic Biol Med* 52: 1101–1110, 2012.
345. Sobotta MC, Liou W, Stocker S, Talwar D, Oehler M, Ruppert T, Scharf AN, and Dick TP. Peroxiredoxin-2 and STAT3 form a redox relay for H₂O₂ signaling. *Nat Chem Biol* 11: 64–70, 2015.
346. Sohn J and Rudolph J. Catalytic and chemical competence of regulation of Cdc25 phosphatase by oxidation/reduction. *Biochemistry* 42: 10060–10070, 2003.
347. Soonsanga S, Lee JW, and Helmann JD. Conversion of *Bacillus subtilis* OhrR from a 1-Cys to a 2-Cys peroxide sensor. *J Bacteriol* 190: 5738–5745, 2008.
348. St-Pierre J, Buckingham JA, Roebuck SJ, and Brand MD. Topology of superoxide production from different sites in the mitochondrial electron transport chain. *J Biol Chem* 277: 44784–44790, 2002.
349. Stamnaes J, Pinkas DM, Fleckenstein B, Khosla C, and Sollid LM. Redox regulation of transglutaminase 2 activity. *J Biol Chem* 285: 25402–25409, 2010.

350. Stangherlin A and Reddy AB. Regulation of circadian clocks by redox homeostasis. *J Biol Chem* 288: 26505–26511, 2013.
351. Staudacher V, Trujillo M, Diederichs T, Dick TP, Radi R, Morgan B, and Deponte M. Redox-sensitive GFP fusions for monitoring the catalytic mechanism and inactivation of peroxiredoxins in living cells. *Redox Biol* 14: 549–556, 2018.
352. Stewart EJ, Aslund F, and Beckwith J. Disulfide bond formation in the *Escherichia coli* cytoplasm: an in vivo role reversal for the thioredoxins. *EMBO J* 17: 5543–5550, 1998.
353. Stocker S, Maurer M, Ruppert T, and Dick TP. A role for 2-Cys peroxiredoxins in facilitating cytosolic protein thiol oxidation. *Nat Chem Biol* 14: 148–155, 2018.
354. Stocker S, Van Laer K, Mijuskovic A, and Dick TP. The conundrum of hydrogen peroxide signaling and the emerging role of peroxiredoxins as redox relay hubs. *Antioxid Redox Signal* 28: 558–573, 2018.
355. Su D, Gaffrey MJ, Guo J, Hatchell KE, Chu RK, Clauss TR, Aldrich JT, Wu S, Purvine S, Camp DG, Smith RD, Thrall BD, and Qian WJ. Proteomic identification and quantification of S-glutathionylation in mouse macrophages using resin-assisted enrichment and isobaric labeling. *Free Radic Biol Med* 67: 460–470, 2014.
356. Sundaresan M, Yu ZX, Ferrans VJ, Irani K, and Finkel T. Requirement for generation of H_2O_2 for platelet-derived growth-factor signal-transduction. *Science* 270: 296–299, 1995.
357. Sutton KA, Black PJ, Mercer KR, Garman EF, Owen RL, Snell EH, and Bernhard WA. Insights into the mechanism of X-ray-induced disulfide-bond cleavage in lysozyme crystals based on EPR, optical absorption and X-ray diffraction studies. *Acta Crystallogr D Biol Crystallogr* 69: 2381–2394, 2013.
358. Svintradze DV, Peterson DL, Collazo-Santiago EA, Lewis JP, and Wright HT. Structures of the *Porphyromonas gingivalis* OxyR regulatory domain explain differences in expression of the OxyR regulon in *Escherichia coli* and *P. gingivalis*. *Acta Crystallogr D Biol Crystallogr* 69: 2091–2103, 2013.
359. Szczepanek K, Chen Q, Derecka M, Salloum FN, Zhang Q, Szelag M, Cichy J, Kukreja RC, Dulak J, Lesniewsky EJ, and Lerner AC. Mitochondrial-targeted signal transducer and activator of transcription 3 (STAT3) protects against ischemia-induced changes in the electron transport chain and the generation of reactive oxygen species. *J Biol Chem* 286: 29610–29620, 2011.
360. Szczepanek K, Chen Q, Lerner AC, and Lesniewsky EJ. Cytoprotection by the modulation of mitochondrial electron transport chain: the emerging role of mitochondrial STAT3. *Mitochondrion* 12: 180–189, 2012.
361. Tachibana T, Okazaki S, Murayama A, Naganuma A, Nomoto A, and Kuge S. A major peroxiredoxin-induced activation of Yap1 transcription factor is mediated by reduction-sensitive disulfide bonds and reveals a low level of transcriptional activation. *J Biol Chem* 284: 4464–4472, 2009.
362. Tairum CA, Santos MC, Breyer CA, Geyer RR, Nieves CJ, Portillo-Ledesma S, Ferrer-Sueta G, Toledo JC, Jr., Toyama MH, Augusto O, Netto LE, and de Oliveira MA. Catalytic Thr or Ser residue modulates structural switches in 2-Cys peroxiredoxin by distinct mechanisms. *Sci Rep* 6: 33133, 2016.
363. Takahashi K, Taira T, Niki T, Seino C, Iguchi-Ariga SM, and Ariga H. DJ-1 positively regulates the androgen receptor by impairing the binding of PIAS α to the receptor. *J Biol Chem* 276: 37556–37563, 2001.
364. Takanishi CL, Ma LH, and Wood MJ. A genetically encoded probe for cysteine sulfenic acid protein modification in vivo. *Biochemistry* 46: 14725–14732, 2007.
365. Takanishi CL and Wood MJ. A genetically encoded probe for the identification of proteins that form sulfenic acid in response to H_2O_2 in *Saccharomyces cerevisiae*. *J Proteome Res* 10: 2715–2724, 2011.
366. Tao R, Zhao Y, Chu H, Wang A, Zhu J, Chen X, Zou Y, Shi M, Liu R, Su N, Du J, Zhou H-M, Zhu L, Qian X, Liu H, Loscalzo J, and Yang Y. Genetically encoded fluorescent sensors reveal dynamic regulation of NADPH metabolism. *Nat Methods* 14: 720–728, 2017.
367. Tarrago L, Peterfi Z, Lee BC, Michel T, and Gladyshev VN. Monitoring methionine sulfoxide with stereospecific mechanism-based fluorescent sensors. *Nat Chem Biol* 11: 332–338, 2015.
368. Tarze A, Deniaud A, Le Bras M, Maillier E, Molle D, Larochette N, Zamzami N, Jan G, Kroemer G, and Brenner C. GAPDH, a novel regulator of the pro-apoptotic mitochondrial membrane permeabilization. *Oncogene* 26: 2606–2620, 2007.
369. Teixeira F, Castro H, Cruz T, Tse E, Koldewey P, Southworth DR, Tomas AM, and Jakob U. Mitochondrial peroxiredoxin functions as crucial chaperone reservoir in *Leishmania infantum*. *Proc Natl Acad Sci U S A* 112: E616–E624, 2015.
370. Teramoto H, Inui M, and Yukawa H. OxyR acts as a transcriptional repressor of hydrogen peroxide-inducible antioxidant genes in *Corynebacterium glutamicum* R. *FEBS J* 280: 3298–3312, 2013.
371. Thangima Zannat M, Bhattacharjee RB, and Bag J. In the absence of cellular poly (A) binding protein, the glycolytic enzyme GAPDH translocated to the cell nucleus and activated the GAPDH mediated apoptotic pathway by enhancing acetylation and serine 46 phosphorylation of p53. *Biochem Biophys Res Commun* 409: 171–176, 2011.
372. Tian SS, Lamb P, Seidel HM, Stein RB, and Rosen J. Rapid activation of the STAT3 transcription factor by granulocyte colony-stimulating factor. *Blood* 84: 1760–1764, 1994.
373. Tobiume K, Matsuzawa A, Takahashi T, Nishitoh H, Morita K, Takeda K, Minowa O, Miyazono K, Noda T, and Ichijo H. ASK1 is required for sustained activations of JNK/p38 MAP kinases and apoptosis. *EMBO Rep* 2: 222–228, 2001.
374. Toledano MB, Kullik I, Trinh F, Baird PT, Schneider TD, and Storz G. Redox-dependent shift of OxyR-DNA contacts along an extended DNA-binding site: a mechanism for differential promoter selection. *Cell* 78: 897–909, 1994.
375. Tomalin LE, Day AM, Underwood ZE, Smith GR, Dalle Pezze P, Rallis C, Patel W, Dickinson BC, Bahler J, Brewer TF, Chang CJ, Shanley DP, and Veal EA. Increasing extracellular H_2O_2 produces a bi-phasic response in intracellular H_2O_2 , with peroxiredoxin hyperoxidation only triggered once the cellular H_2O_2 -buffering capacity is overwhelmed. *Free Radic Biol Med* 95: 333–348, 2016.
376. Torres MA, Dangel JL, and Jones JD. Arabidopsis gp91phox homologues AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates in the

- plant defense response. *Proc Natl Acad Sci U S A* 99: 517–522, 2002.
377. Tran HJ, Heroven AK, Winkler L, Spreter T, Beatrix B, and Dersch P. Analysis of RovA, a transcriptional regulator of *Yersinia pseudotuberculosis* virulence that acts through antirepression and direct transcriptional activation. *J Biol Chem* 280: 42423–42432, 2005.
 378. Tristan C, Shahani N, Sedlak TW, and Sawa A. The diverse functions of GAPDH: views from different subcellular compartments. *Cell Signal* 23: 317–323, 2011.
 379. Tristan CA, Ramos A, Shahani N, Emiliani FE, Nakajima H, Noeh CC, Kato Y, Takeuchi T, Noguchi T, Kadowaki H, Sedlak TW, Ishizuka K, Ichijo H, and Sawa A. Role of apoptosis signal-regulating kinase 1 (ASK1) as an activator of the GAPDH-Siah1 stress-signaling cascade. *J Biol Chem* 290: 56–64, 2015.
 380. Trujillo M, Clippe A, Manta B, Ferrer-Sueta G, Smeets A, Declercq JP, Knoop B, and Radi R. Pre-steady state kinetic characterization of human peroxiredoxin 5: taking advantage of Trp84 fluorescence increase upon oxidation. *Arch Biochem Biophys* 467: 95–106, 2007.
 381. Tseng HJ, McEwan AG, Apicella MA, and Jennings MP. OxyR acts as a repressor of catalase expression in *Neisseria gonorrhoeae*. *Infect Immun* 71: 550–556, 2003.
 382. Turner-Ivey B, Manevich Y, Schulte J, Kistner-Griffin E, Jezierska-Drutel A, Liu Y, and Neumann CA. Role for Prdx1 as a specific sensor in redox-regulated senescence in breast cancer. *Oncogene* 32: 5302–5314, 2013.
 383. Ushioda R, Miyamoto A, Inoue M, Watanabe S, Okumura M, Maegawa KI, Uegaki K, Fujii S, Fukuda Y, Umitsu M, Takagi J, Inaba K, Mikoshiba K, and Nagata K. Redox-assisted regulation of Ca²⁺ homeostasis in the endoplasmic reticulum by disulfide reductase ERdj5. *Proc Natl Acad Sci U S A* 113: E6055–E6063, 2016.
 384. van der Brug MP, Blackinton J, Chandran J, Hao LY, Lal A, Mazan-Mamczarz K, Martindale J, Xie C, Ahmad R, Thomas KJ, Beilina A, Gibbs JR, Ding J, Myers AJ, Zhan M, Cai H, Bonini NM, Gorospe M, and Cookson MR. RNA binding activity of the recessive parkinsonism protein DJ-1 supports involvement in multiple cellular pathways. *Proc Natl Acad Sci U S A* 105: 10244–10249, 2008.
 385. Vazquez-Mayorga E, Diaz-Sanchez AG, Dagda RK, Dominguez-Solis CA, Dagda RY, Coronado-Ramirez CK, and Martinez-Martinez A. Novel redox-dependent esterase activity (EC 3.1.1.2) for DJ-1: implications for Parkinson's disease. *Int J Mol Sci* 17: 1346, 2016.
 386. Veal EA, Findlay VJ, Day AM, Bozonet SM, Evans JM, Quinn J, and Morgan BA. A 2-Cys peroxiredoxin regulates peroxide-induced oxidation and activation of a stress-activated MAP kinase. *Mol Cell* 15: 129–139, 2004.
 387. Veal EA, Ross SJ, Malakasi P, Peacock E, and Morgan BA. Ybp1 is required for the hydrogen peroxide-induced oxidation of the Yap1 transcription factor. *J Biol Chem* 278: 30896–30904, 2003.
 388. Venugopal R and Jaiswal AK. Nrf1 and Nrf2 positively and c-Fos and Fra1 negatively regulate the human antioxidant response element-mediated expression of NAD(P)H:quinone oxidoreductase1 gene. *Proc Natl Acad Sci U S A* 93: 14960–14965, 1996.
 389. Vignols F, Mouaheb N, Thomas D, and Meyer Y. Redox control of Hsp70-Co-chaperone interaction revealed by expression of a thioredoxin-like Arabidopsis protein. *J Biol Chem* 278: 4516–4523, 2003.
 390. Vinothkumar KR and Henderson R. Single particle electron cryomicroscopy: trends, issues and future perspective. *Q Rev Biophys* 49: e13, 2016.
 391. Waak J, Weber SS, Gerner K, Schall C, Ichijo H, Stehle T, and Kahle PJ. Oxidizable residues mediating protein stability and cytoprotective interaction of DJ-1 with apoptosis signal-regulating kinase 1. *J Biol Chem* 284: 14245–14257, 2009.
 392. Wallin E and von Heijne G. Genome-wide analysis of integral membrane proteins from eubacterial, archaean, and eukaryotic organisms. *Protein Sci* 7: 1029–1038, 1998.
 393. Wang Q, Woltjer RL, Cimino PJ, Pan C, Montine KS, Zhang J, and Montine TJ. Proteomic analysis of neurofibrillary tangles in Alzheimer disease identifies GAPDH as a detergent-insoluble paired helical filament tau binding protein. *FASEB J* 19: 869–871, 2005.
 394. Wang T, Si M, Song Y, Zhu W, Gao F, Wang Y, Zhang L, Zhang W, Wei G, Luo ZQ, and Shen X. Type VI secretion system transports Zn²⁺ to combat multiple stresses and host immunity. *PLoS Pathog* 11: e1005020, 2015.
 395. Wang X, Wang L, Wang X, Sun F, and Wang CC. Structural insights into the peroxidase activity and inactivation of human peroxiredoxin 4. *Biochem J* 441: 113–118, 2012.
 396. Wang Y, Ande SR, and Mishra S. Phosphorylation of transglutaminase 2 (TG2) at serine-216 has a role in TG2 mediated activation of nuclear factor-kappa B and in the downregulation of PTEN. *BMC Cancer* 12: 277, 2012.
 397. Wang YT, Tzeng DW, Wang CY, Hong JY, and Yang JL. APE1/Ref-1 prevents oxidative inactivation of ERK for G1-to-S progression following lead acetate exposure. *Toxicology* 305: 120–129, 2013.
 398. Wang ZW, Nara M, Wang YX, and Kotlikoff MI. Redox regulation of large conductance Ca(2+)-activated K+ channels in smooth muscle cells. *J Gen Physiol* 110: 35–44, 1997.
 399. Wardman P. Fluorescent and luminescent probes for measurement of oxidative and nitrosative species in cells and tissues: progress, pitfalls, and prospects. *Free Radic Biol Med* 43: 995–1022, 2007.
 400. Waszczak C, Akter S, Eeckhout D, Persiau G, Wahni K, Bodra N, Van Molle I, De Smet B, Vertommen D, Gevaert K, De Jaeger G, Van Montagu M, Messens J, and Van Breusegem F. Sulfenome mining in *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A* 111: 11545–11550, 2014.
 401. Weik M, Berges J, Raves ML, Gros P, McSweeney S, Silman I, Sussman JL, Houee-Levin C, and Ravelli RB. Evidence for the formation of disulfide radicals in protein crystals upon X-ray irradiation. *J Synchrotron Radiat* 9: 342–346, 2002.
 402. Wilke ME, Higaki JN, Craik CS, and Fletterick RJ. Crystal structure of rat trypsin-S195C at –150 degrees C. Analysis of low activity of recombinant and semisynthetic thiol proteases. *J Mol Biol* 219: 511–523, 1991.
 403. Winterbourn CC. Reconciling the chemistry and biology of reactive oxygen species. *Nat Chem Biol* 4: 278–286, 2008.
 404. Winterbourn CC and Hampton MB. Thiol chemistry and specificity in redox signaling. *Free Radic Biol Med* 45: 549–561, 2008.
 405. Wojdyla K, Williamson J, Roepstorff P, and Rogowska-Wrzesinska A. The SNO/SOH TMT strategy for combinatorial analysis of reversible cysteine oxidations. *J Proteomics* 113: 415–434, 2015.
 406. Woo HA, Yim SH, Shin DH, Kang D, Yu DY, and Rhee SG. Inactivation of peroxiredoxin I by phosphorylation

- allows localized H₂O₂ accumulation for cell signaling. *Cell* 140: 517–528, 2010.
407. Wood ZA, Poole LB, Hantgan RR, and Karplus PA. Dimers to doughnuts: redox-sensitive oligomerization of 2-cysteine peroxiredoxins. *Biochemistry* 41: 5493–5504, 2002.
 408. Wood ZA, Poole LB, and Karplus PA. Peroxiredoxin evolution and the regulation of hydrogen peroxide signaling. *Science* 300: 650–653, 2003.
 409. Xu G and Chance MR. Radiolytic modification of sulfur-containing amino acid residues in model peptides: fundamental studies for protein footprinting. *Anal Chem* 77: 2437–2449, 2005.
 410. Xu J, Zhong N, Wang H, Elias JE, Kim CY, Woldman I, Pifl C, Gygi SP, Geula C, and Yankner BA. The Parkinson's disease-associated DJ-1 protein is a transcriptional co-activator that protects against neuronal apoptosis. *Hum Mol Genet* 14: 1231–1241, 2005.
 411. Yaffe MB, Schutkowski M, Shen M, Zhou XZ, Stukenberg PT, Rahfeld JU, Xu J, Kuang J, Kirschner MW, Fischer G, Cantley LC, and Lu KP. Sequence-specific and phosphorylation-dependent proline isomerization: a potential mitotic regulatory mechanism. *Science* 278: 1957–1960, 1997.
 412. Yang J, Carroll KS, and Liebler DC. The expanding landscape of the thiol redox proteome. *Mol Cell Proteomics* 15: 1–11, 2016.
 413. Yang J, Gupta V, Carroll KS, and Liebler DC. Site-specific mapping and quantification of protein S-sulphenylation in cells. *Nat Commun* 5: 4776, 2014.
 414. Yang J, Gupta V, Tallman KA, Porter NA, Carroll KS, and Liebler DC. Global, in situ, site-specific analysis of protein S-sulphenylation. *Nat Protoc* 10: 1022–1037, 2015.
 415. Yang KS, Kang SW, Woo HA, Hwang SC, Chae HZ, Kim K, and Rhee SG. Inactivation of human peroxiredoxin I during catalysis as the result of the oxidation of the catalytic site cysteine to cysteine-sulfinic acid. *J Biol Chem* 277: 38029–38036, 2002.
 416. Yewdall NA, Venugopal H, Desfosses A, Abrishami V, Yosaatmadja Y, Hampton MB, Gerrard JA, Goldstone DC, Mitra AK, and Radjainia M. Structures of human peroxiredoxin 3 suggest self-chaperoning assembly that maintains catalytic state. *Structure* 24: 1120–1129, 2016.
 417. Yi MC, Melkonian AV, Ousey JA, and Khosla C. Endoplasmic reticulum-resident protein 57 (ERp57) oxidatively inactivates human transglutaminase 2. *J Biol Chem* 293: 2640–2649, 2018.
 418. Zaffagnini M, Fermani S, Calvaresi M, Orru R, Iommarini L, Sparla F, Falini G, Bottoni A, and Trost P. Tuning cysteine reactivity and sulfenic acid stability by protein microenvironment in glyceraldehyde-3-phosphate dehydrogenases of *Arabidopsis thaliana*. *Antioxid Redox Signal* 24: 502–517, 2016.
 419. Zeida A, Babbush R, Lebrero MC, Trujillo M, Radi R, and Estrin DA. Molecular basis of the mechanism of thiol oxidation by hydrogen peroxide in aqueous solution: challenging the SN2 paradigm. *Chem Res Toxicol* 25: 741–746, 2012.
 420. Zhai D, Chin K, Wang M, and Liu F. Disruption of the nuclear p53-GAPDH complex protects against ischemia-induced neuronal damage. *Mol Brain* 7: 20, 2014.
 421. Zhang T, Zhu M, Zhu N, Strul JM, Dufresne CP, Schneider JD, Harmon AC, and Chen S. Identification of thioredoxin targets in guard cell enriched epidermal peels using cystTMT proteomics. *J Proteomics* 133: 48–53, 2016.
 422. Zheng J, Koblinski JE, Dutson LV, Feeney YB, and Clevenger CV. Prolyl isomerase cyclophilin A regulation of Janus-activated kinase 2 and the progression of human breast cancer. *Cancer Res* 68: 7769–7778, 2008.
 423. Zheng M, Aslund F, and Storz G. Activation of the OxyR transcription factor by reversible disulfide bond formation. *Science* 279: 1718–1721, 1998.
 424. Zhu R, Hao Z, Lou H, Song Y, Zhao J, Chen Y, Zhu J, and Chen PR. Structural characterization of the DNA-binding mechanism underlying the copper(II)-sensing MarR transcriptional regulator. *J Biol Inorg Chem* 22: 685–693, 2017.
 425. Zykova TA, Zhu F, Vakorina TI, Zhang J, Higgins LA, Urusova DV, Bode AM, and Dong Z. T-LAK cell-originated protein kinase (TOPK) phosphorylation of Prx1 at Ser-32 prevents UVB-induced apoptosis in RPMI7951 melanoma cells through the regulation of Prx1 peroxidase activity. *J Biol Chem* 285: 29138–29146, 2010.

Address correspondence to:

Dr. Joris Messens

Center for Structural Biology

VIB

Brussels Center for Redox Biology

Structural Biology Brussels

Vrije Universiteit Brussel

Pleinlaan 2

1050 Brussels

Belgium

E-mail: joris.messens@vib-vub.be

Date of first submission to ARS Central, January 29, 2017; date of final revised submission, April 5, 2018; date of acceptance, April 8, 2018.

Abbreviations Used

AbfR	= aggregation and biofilm formation regulator
ACHT	= atypical chloroplast thioredoxin
Ahp	= alkyl hydroperoxide
ANT	= adenine nucleotide translocase
AOP	= antioxidant protein
APE1	= apurinic/aprimidinic endonuclease 1
ASK1	= apoptosis signal-regulating kinase 1
AtTDX	= <i>Arabidopsis thaliana</i> tetratricopeptide repeat domain-containing thioredoxin
Bcl-X _L	= B cell lymphoma-extra large
BifR	= biofilm regulator
Cdc25	= cell division cycle 25 (phosphatase)
CDK	= cyclin-dependent kinase
CFBPase	= chloroplast fructose-1,6-bisphosphatase
CLIC	= chloride intracellular channel protein
C _P	= peroxidatic cysteine
cpYFP	= circularly permuted yellow fluorescent protein
C _R	= resolving cysteine
cryo-EM	= cryo-electron microscopy
Cyp	= cyclophilin
DA	= azide-functionalized dimedone
DBD	= DNA-binding domain
DJ-1	= protein deglycanase-1

Abbreviations Used (Cont.)

DTT = dithiothreitol
 DY = alkyne-functionalized dimer
 ERK = extracellular signal-regulated kinase
 ERp57 = endoplasmic reticulum-resident protein 57 (also known as protein disulfide-isomerase A3, or glucose-regulated protein 58-kD)
 FF = fully folded
 FOXO = forkhead box protein O
 GAPDH = glyceraldehyde 3-phosphate dehydrogenase
 GPx = glutathione peroxidase
 H₂O₂ = hydrogen peroxide
 HIP = Hsp70-interacting protein
 HMW = high-molecular-weight
 Hsp = heat shock protein
 IAM = iodoacetamide
 ICATs = isotope-coded affinity tags
 iodo-TMT = iodoacetyl-based tandem mass tag
 JAK2 = Janus kinase 2
 JNK = c-Jun N-terminal kinase
 Keap1 = Kelch-like ECH-associated protein 1
 LMW = low-molecular-weight
 LU = locally unfolded
 MAPK = mitogen-activated protein kinase
 MarR = multiple antibiotic resistance regulator
 MexR = multidrug efflux regulator
 MKP = MAPK phosphatase
 mPTP = mitochondrial permeability transition pore
 MS = mass spectrometry
 Msr = methionine sulfoxide reductase
 MST1 = mammalian sterile20-like kinase 1
 NAD(P)H = nicotinamide adenine dinucleotide phosphate
 NEM = N-ethylmaleimide
 NemR = N-ethylmaleimide regulator
 NMR = nuclear magnetic resonance
 NOX = NADPH oxidase
 Nrf2 = nuclear factor-erythroid 2 p45-related factor 2
 O₂^{•-} = superoxide

•OH = hydroxyl radical
 OhrR = organic hydroperoxide resistance regulator
 Orp1 = oxidant receptor peroxidase 1
 OxyR = oxidative receptor
 p66Shc = p66 Src homologous-collagen homolog
 PDI = protein disulfide isomerase
 Pin1 = peptidyl-prolyl *cis-trans* isomerase NIMA-interacting 1
 PPI = peptidyl-prolyl *cis-trans* isomerase
 Prx = peroxiredoxin
 PrxA = peroxiredoxin A
 PSF = polypyrimidine tract-binding protein-associated splicing factor
 PTEN = phosphatase and tensin homolog
 PTP = protein tyrosine phosphatase
 RCS = reactive chlorine species
 RD = regulatory domain
 roGFP = redox-sensitive green fluorescent protein
 ROS = reactive oxygen species
 ROOH = peroxide
 SOD = superoxide dismutase
 Src = sarcoma
 Srx = sulfiredoxin
 STAT = signal transducer and activator of transcription
 TBD = Trx-binding domain
 TCEP = tris(2-carboxyethyl)phosphine
 TG = transglutaminase (also known as tissue transglutaminase)
 TG2 = transglutaminase 2
 TLR4 = Toll-like receptor 4
 Tpx = Trx peroxidase
 TR = thioredoxin reductase
 Trx = thioredoxin
 Tsa1 = thiol-specific antioxidant 1
 Tsa2 = thiol-specific antioxidant 2
 Txnip = Trx-interacting protein
 VDAC = voltage-dependent anion channel
 wHTH = winged helix-turn-helix
 YAP1 = yeast AP1-like protein
 Ybp1 = Yap1-binding protein 1