REDOX PIONEER #14

Redox Pioneer: Professor Hideo Kimura

David Lefer

Professor Hideo Kimura

Abstract

Dr. Hideo Kimura is recognized as a redox pioneer because he has published an article in the field of antioxidant and redox biology that has been cited >1000 times, and 29 articles that have been cited >100 times. Since the first description of hydrogen sulfide (H_2S) as a toxic gas 300 years ago, most studies have been devoted to its toxicity. In 1996, Dr. Kimura demonstrated a physiological role of H_2S as a mediator of cognitive function and cystathionine β -synthase as an H_2S -producing enzyme. In the following year, he showed H_2S as a vascular smooth muscle relaxant in synergy with nitric oxide and its production by cystathionine γ -lyase in vasculature. Subsequently he reported the cytoprotective effect of H_2S on neurons against oxidative stress. Since then, studies on H_2S have unveiled numerous physiological roles such as the regulation of inflammation, cell growth, oxygen sensing, and senescence. He also discovered polysulfides (H_2S_n), which have a higher number of sulfur atoms than H_2S and are one of the active forms of H_2S , as potent signaling molecules produced by 3-mercaptopyruvate sulfurtransferase. H_2S_n regulate ion channels and transcription factors to upregulate antioxidant genes, tumor suppressors, and protein kinases to, in turn, regulate blood pressure. These findings led to the re-evaluation of other persulfurated molecules such as cysteine persulfide and glutathione persulfide. Dr. Kimura is a pioneer of studies on H_2S and H_2S_n as signaling molecules. *Antioxid. Redox Signal.* 30, 1699–1708.

It is fortunate to come across a secret of nature and pick it up.

—Prof. Hideo Kimura

Keywords: redox pioneer, polysulfides, hydrogen sulfide, signaling, nitric oxide

Background Development and Training

DR. KIMURA GRADUATED FROM THE University of Tokyo, Faculty of Pharmaceutical Sciences, in 1980 and received his PhD from the University of Tokyo in 1985. He studied neurotransmitters in the cerebellum using electrophysiological techniques at the National Defense Medical College (33, 64), and the gene structure of cytochrome P-450 at the Cancer Institute (35, 36). He com-

pleted his postdoctoral studies at the Salk Institute for Biological Studies where he identified a novel growth factor, Schwannoma-derived growth factor, as well as activin (28, 32, 34, 71). He continued working at the Salk Institute as a senior staff scientist, where he identified presenilin-binding protein (PBP), a novel guanine nucleotide exchange factor that activates Rac (27). PBP was later renamed as modifier of cell adhesion (9, 60), and finally as dedicator of cytokines 3 (Dock3) (61).

Reviewing Editors: Philip Aaronson, Martin Diener, Milos Filipovic, Christine Foyer, Gabriel Gojon, and Gregg Semenza

CV Center of Excellence, Louisiana State University Health Sciences Center, New Orleans, Louisiana.

Author note: The author met Dr. Kimura at Experimental Biology 2008 to which the author invited him. Since then, they have enjoyed a decade of academic interactions.

For a list of frequently cited articles published by Prof. Hideo Kimura, see Supplementary Tables S1 and S2, available online.

© David Lefer 2018; Published by Mary Ann Liebert, Inc. This Open Access article is distributed by the terms of the Creative Commons License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

FIG. 1. H_2S modulates the activity of neurons. H_2S facilitates the induction of hippocampal LTP, a synaptic model of memory formation (A), by enhancing the activity of NMDA receptor, leaving another type of glutamate receptor, AMPA receptor, unaffected (B) (1). Although a mechanism for the activation of NMDA receptors had been proposed that the reduction of cysteine disulfide bond located at the hinge of the ligand-binding domain by DTT activates the receptor (2), it is not able to fully explain the activation by a weak reducing molecule such as H_2S . At much lower concentrations, H_2S ($100 \mu M$) induced LTP more efficiently than 1 mM DTT (C) (1). Lower dotted line: control EPSP slope; upper dotted line: EPSP slope in the presence of DTT. This finding led to the identification of H_2S_n as novel signaling molecules (40–43, 55, 59, 66) (see Key Finding 3). AMPA, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; DTT, dithiothreitol; EPSP, excitatory post-synaptic potential; H_2S , hydrogen sulfide; LTP, long-term potentiation; NMDA, *N*-methyl D-aspartate.

Summary of Dr. Kimura's Top Contributions

Dr. Kimura showed the first time that hydrogen sulfide (H_2S) is a novel signaling molecule in multiple body systems. He discovered a physiological role of H_2S as a

mediator of memory formation (1) (Fig. 1), and subsequently identified another role as a vascular smooth muscle relaxant in synergy with nitric oxide (NO) that was the first demonstration of a crosstalk between H_2S and NO (21) (Fig. 2). Prominent neuroscientist Solomon Snyder

FIG. 2. H_2S relaxes vascular smooth muscle in synergy with NO. Synergistic effect of H_2S and NO was discovered on vascular smooth muscle relaxation. The simultaneous application of H_2S and NO donors induced greater relaxation of vascular smooth muscle than either H_2S or NO alone (21). Potentiation of the relaxation effects of Na-nitroprusside (A) and morpholinosydnonimine (B) by $30 \mu M$ NaHS. This finding opened the study of crosstalk between H_2S and NO and led to the identification of H_2S_n generated by the chemical interaction of both molecules (52) (see Key Finding 3). NO, nitric oxide.

FIG. 3. H_2S protects neurons from oxidative stress. Since H_2S is a well-known toxic gas, its cytoprotective effects have been overlooked. Neurons in primary culture were killed by oxidative stress induced by high concentrations of glutamate, while surviving in the presence of NaHS (39).

commented in *Science News*, “They have very impressive evidence that H_2S is a potential neurotransmitter. It’s an exciting paper that should stimulate a lot of people’s interest” (84).

Dr. Kimura discovered the cytoprotective effect of H_2S on neurons from oxidative stress (39) (Fig. 3), and the effect on other tissues and organs such as heart and kidney was followed (14, 85). These findings led to the identification of numerous physiological roles of this molecule, including anti-inflammation, angiogenesis, oxygen sensing, and ATP formation (11, 51, 65, 67, 68, 82, 93, 94).

During further investigation of the physiological roles of H_2S , Dr. Kimura discovered novel signaling molecules, H_2S_n , produced by 3-mercaptopyruvate sulfurtransferase (3MST) (41, 42, 58, 59, 66) (Fig. 4). Since then, studies on H_2S_n have unveiled various physiological roles such as the regulation of ion channels, transcription factors, protein kinases, and tumor suppressors, as well as the production of other per- and polysulfurated molecules (18, 41, 44, 80). Dr. Kimura also identified H_2S_n produced by the chemical interaction of H_2S with NO that may provide one of the mechanisms of the synergy between H_2S and NO (52) (Fig. 5).

FIG. 4. H_2S_n induce Ca^{2+} influx in astrocytes. During the study of the effect of H_2S on astrocytes, Dr. Kimura’s group found that a solution of NaHS, a sodium salt of H_2S whose color is yellowish, activates astrocytes much more efficiently than a colorless solution. The study identified that the yellowish color is derived from H_2S_n generated by the oxidation of H_2S (40–42, 59, 66). H_2S_3 induced Ca^{2+} influx with an EC_{50} value (91 nM) (A, B), approximately 1/1000th of that of H_2S (116 μM) (41). Time required for the induction of Ca^{2+} influx by Na_2S_3 (C). Endogenous H_2S_2 and H_2S_3 were identified in the brain (D) (40–43, 55). The enlarged peaks a and b are shown in the inset. Major peaks for Na_2S_3 and Na_2S_4 are b and c, respectively. Since the peaks a, b, and c were observed in a control sample with monobromobimane alone, it was subtracted from each peak of the sample to estimate the amount of polysulfides in brain homogenates.

FIG. 5. H_2S_n are generated from H_2S and NO. Chemical interaction between H_2S and NO produces H_2S_2 and H_2S_3 (52). It may be one of the mechanisms for the synergistic effects of H_2S and NO on various tissues, including vascular smooth muscle relaxation (12, 21) as shown in Figure 2.

Description of Key Finding 1

In the library at the Salk Institute, Dr. Kimura found a metabolic map showing mammalian enzymes that can produce H_2S . These enzymes were intensively studied from the 1950s to 1970s; cystathionine β -synthase (CBS), cystathionine γ -lyase (CSE), and 3MST together with cysteine amino transferase were found to have the capacity *in vitro* (5, 8, 48, 79), and their full-length cDNAs were cloned in the 1990s (56, 62, 81). However, rather than being recognized as a physiologically active molecule in these early studies, H_2S was thought of simply as a byproduct of metabolic pathways or a marker for the evaluation of enzyme activity.

Three groups discovered endogenous sulfide in mammalian brains when measuring sulfide levels in intoxicated animals (17, 70, 88). Survivors of H_2S poisoning experienced memory loss, and acute intoxication with H_2S caused changes in the levels of neurotransmitters in the brains of animal models. Inspired by these findings, Dr. Kimura began studying H_2S as a signaling molecule in 1993 when carbon

monoxide had just been identified as a gaseous signaling molecule (47, 78, 86, 95), following on NO (4, 16, 23).

In 1996, Dr. Kimura together with his student Dr. Kazuho Abe demonstrated that H_2S , which can be produced by CBS in the brain, facilitates the induction of hippocampal long-term potentiation (LTP), a synaptic model of memory formation (1) (Figs. 1 and 6). H_2S enhances the activity of one type of glutamate receptor, *N*-methyl D-aspartate (NMDA) receptor, leaving another type of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor, which is also activated by glutamate, unaffected. A mechanism for the activation of NMDA receptors had been proposed by Aizenman *et al.* using dithiothreitol (DTT), which reduces cysteine disulfide bond located at the hinge of the ligand-binding domain to activate the receptor (2). However, this mechanism is not able to fully explain the activation by a weak reducing molecule such as H_2S . At much lower concentrations, H_2S induced LTP more efficiently than DTT (Fig. 1) (1). This finding led to the identification of H_2S_n as novel signaling molecules (see Key Finding 3).

FIG. 6. Facilitation of LTP induction by H_2S and H_2S_n . H_2S enhances the activity of NMDA receptors by reducing the cysteine disulfide bond at the hinge of the ligand-binding domain of the receptors (1). H_2S_n activate TRPA1 channels to induce Ca^{2+} influx in astrocytes (41, 59, 66), which, in turn, release gliotransmitters such as D-serine to enhance the activity of NMDA receptors (77). By these integrated mechanisms, LTP may be effectively induced. Modified from Kimura (29). 3MST, 3-mercaptopyruvate sulfurtransferase; CBS, cystathionine β -synthase; TRPA1, transient receptor potential ankyrin 1.

In 1997, Dr. Kimura demonstrated that H_2S can be produced by CSE to relax vasculature in synergy with NO (21) (Fig. 2). This discovery also opened a field of crosstalk between H_2S and NO (see also Key Finding 3). Subsequently, Wang and colleagues identified K_{ATP} channels as one of the targets of H_2S to relax vascular smooth muscle (94), and Cirino and colleagues showed cyclic GMP-dependent protein kinase as another target to mediate H_2S -induced vasorelaxation (6).

Description of Key Finding 2

Since H_2S is a well-known toxic gas, its cytoprotective effects have been overlooked. Dr. Kimura found that H_2S protects neurons from oxidative stress through enhancing the activity of the cystine/glutamate antiporter and cysteine transporter, as well as γ -glutamyl cysteine synthetase or glutamate cysteine ligase, a rate limiting enzyme in the production of glutathione (GSH), which is a major cellular antioxidant (38, 39) (Figs. 3 and 7). Dr. Kimura also showed that H_2S enhances the activity of K_{ATP} and CFTR Cl^- channels to suppress the excessive excitation of neurons by stabilizing membrane potential (37) (Fig. 7). This finding led to the identification of the cytoprotective effect of H_2S on various tissues and organs, including the heart, kidney, retina, pancreas, and intestines, and the regulation of endoplasmic reticulum stress (14, 26, 45, 49, 72, 85, 93). Several H_2S -based therapeutic compounds have been developed, with some undergoing clinical trials (87). H_2S even plays an essential role in the development of bacterial resistance to an-

tibiotics (73). H_2S is a universal cytoprotectant effective in bacteria and mammals.

Dr. Kimura's group also identified the transcription factor specific protein 1 (SP1)-binding site at the 5'-noncoding region of the CSE gene (25) that contributes to the cytoprotection *via* NF- κ B signaling. Snyder and colleagues identified a mechanism for the antiapoptotic actions induced by tumor necrosis factor alpha (TNF α), where TNF α stimulates the binding of SP1 to the CSE promoter that increases the levels of CSE, resulting in the production of H_2S . H_2S S-sulfurates (sulfhydrates) cysteine residues of the p65 subunit of NF- κ B to facilitate the interaction with another subunit RPS3, leading to upregulation of antiapoptotic genes (72).

H_2S -producing pathway from D-cysteine that also contributes to cytoprotection was also identified by Dr. Kimura's group (74). D-Cysteine is metabolized by D-amino acid oxidase (DAO) to 3-mercaptopyruvate (3MP), which is a substrate of 3MST to produce H_2S (75, 76). In the kidney, H_2S production from D-cysteine is ~ 80 times greater than that from L-cysteine, and the administration of D-cysteine protects the kidney from ischemia-reperfusion injury more efficiently than that of L-cysteine. 3MST is a ubiquitous enzyme, whereas DAO is expressed only in restricted tissues, such as the brain and kidney, in mice. This finding may suggest a new therapeutic approach to deliver H_2S to specific tissues such as the brain and kidney.

Description of Key Finding 3

Astrocytes, a type of glia, were thought to merely support and provide nutrients to neurons, but were recently recognized

FIG. 7. Protection of neurons from oxidative stress by H_2S and H_2S_n . H_2S enhances the activity of the cystine/glutamate antiporter to increase the transport of cystine, which is reduced to cysteine in cells (39), and also enhances the activity of the cysteine transporter (38). H_2S upregulates the activity of a rate limiting enzyme for GSH production, γ -GCS, also known as GCL. By these effects, H_2S increases the production of GSH. H_2S also enhances the activity of K_{ATP} channels and CFTR Cl^- channels to suppress excessive excitation by stabilizing the membrane potential (37), while it suppresses the voltage gated Ca^{2+} channels to decrease Ca^{2+} toxicity by enhancing the activity of H^+ -ATPase (49). In contrast, H_2S_n facilitate the release of Nrf2 from Keap1/Nrf2 complex by sulfurating cysteine residues of Keap1, resulting in the transport of Nrf2 to the nucleus where Nrf2 upregulates the transcription of antioxidant genes, including γ -GCS (GCL) (44). By these integrated mechanisms, H_2S and H_2S_n protect neurons from oxidative stress. γ -GCS, γ -glutamyl cysteine synthetase; GCL, glutamate cysteine ligase; GSH, glutathione; Keap1, kelch-like ECH-associated protein 1; Nrf2, nuclear factor-like 2.

as actively regulating neuronal activity by releasing gliotransmitters such as D-serine to synaptic clefts. Dr. Kimura's group found that H_2S induces Ca^{2+} influx in astrocytes (58). However, during this study, it was found that a solution of NaHS, a sodium salt of H_2S whose color is yellowish, activates astrocytes much more efficiently than a colorless solution. The study identified that the yellowish color is derived from H_2S_n generated by the oxidation of H_2S , and it measured the endogenous H_2S_n in the brain (41, 42, 59, 66) (Fig. 4).

H_2S_3 induced Ca^{2+} influx with an EC_{50} value (91 nM), approximately 1/1000th of that of H_2S (116 μM) (Fig. 4). The effect of H_2S_3 was suppressed by inhibitors as well as siRNAs specific to transient receptor potential ankyrin 1 (TRPA1) channels, suggesting that H_2S_3 activates TRPA1 channels (41). In collaboration with Dr. Ohta, Dr. Kimura identified the target of H_2S_3 as two cysteine residues located at the amino terminus of TRPA1 channels (20). This finding suggests the additional mechanism for the facilitation of LTP induction (see Key Finding 1, and Fig. 6). H_2S_n activates TRPA1 channels to induce Ca^{2+} influx in astrocytes (41, 59, 66), which, in turn, release gliotransmitters such as D-serine to enhance the activity of NMDA receptors (29, 77).

Other roles of H_2S_n were subsequently found (30). In collaboration with Dr. Ogasawara, Dr. Kimura found that H_2S_n facilitate the translocation of nuclear factor-like 2 (Nrf2) to the nucleus by modifying its binding partner kelch-like ECH-associated protein 1 (Keap1) to upregulate the transcription of antioxidant genes (44) (Fig. 7). This system was initially reported to be activated by H_2S (7). H_2S_n regulate the activity of the tumor suppressor phosphatase and tensin homolog (18) and reduce blood pressure by dilating vascular smooth muscle through the activation of protein kinase $\text{G}_{1\alpha}$ (80). These studies facilitated the identification of the production pathways for H_2S_n .

Hylin and Wood reported that persulfurated cysteine residues of proteins were produced from 3MP, a substrate of 3MST (22). Bound sulfane sulfur, which is defined as the sulfur species that releases H_2S under reducing conditions, includes H_2S_n , free cysteine persulfide (Cys-SSH), GSH persulfide (GSSH), and persulfurated cysteine residues of proteins (24, 54, 63, 89). Dr. Kimura's group found that cells expressing 3MST contain higher levels of bound sulfane sulfur than cells expressing defective mutants of 3MST as well as cells without 3MST (76). Oral administration of D-cysteine to mice increases the levels of bound sulfane sulfur in the kidney to which DAO is highly localized (74). As predicted, brains of 3MST knockout mice contain less bound sulfane sulfur than those of wild-type mice (40). These observations suggested that 3MST can also produce a free form of persulfurated molecules such as H_2S_n . In collaboration with Dr. Ogasawara and Dr. Nagahara, Dr. Kimura found that 3MST produces H_2S_2 and H_2S_3 as well as H_2S , and determined the endogenous levels of both molecules (40, 42, 43, 55).

During studies on H_2S_n production by 3MST, Dr. Kimura's group noticed that levels of cysteine and GSH were decreased when 3MST produces H_2S_n . Dr. Kimura concluded that H_2S_n readily reacted with Cys-SH and GSH to produce Cys-SSH and GSSH (40). Alternatively, 3MST may transfer sulfur from 3MP to cysteine and GSH to produce these persulfurated species (Fig. 8).

3MST requires a reducing substance to produce H_2S , but an endogenous reducing substance was not known. From the structure of leishmania 3MST, which contains a thioredoxin domain in the molecule, the interaction of thioredoxin with 3MST had been predicted (90). Dr. Kimura's group determined thioredoxin as well as dihydrolipoic acid as potential endogenous reducing molecules necessary for 3MST to produce H_2S (50, 57, 92).

FIG. 8. Production of H_2S , H_2S_n , Cys-SSH, GSSH, and S-sulfurated proteins by 3MST and the effects of these molecules. 3MST produces H_2S_n and H_2S by interacting with thioredoxin (Trx) (40–42, 50). H_2S_n are also generated by the interaction of H_2S with NO or the oxidation of H_2S (52). 3MST can also produce Cys-SSH and GSSH directly or through the generation of H_2S_n , which readily react with Cys-SH and GSH (40). H_2S reduces the cysteine disulfide bond to induce the conformational changes of target proteins, resulting in the regulation of their activity (1). H_2S_n S-sulfurates (sulfhydrylate) cysteine residues of target proteins (54), whereas under oxidative stress or by NO signaling (dotted box) H_2S S-sulfurates the oxidized cysteine residues such as nitrosylated (Cys-SNO) and sulfenic acid (Cys-SOH) to regulate the activity of targets. (Reprinting from Ref. 31). CSE, cystathionine γ -lyase.

3MST thiolates tRNA to maintain the accuracy of the genetic code and stabilize the tRNA structure (46, 91). There are two isoforms of 3MST that localize to the cytosol and mitochondria in humans (15). Cytosolic isoforms thiolate tRNA, whereas the mitochondrial isoform has a dual localization in both mitochondria and the cytosol, and not only thiolates tRNA in the cytosol but also supplies sulfur for iron–sulfur cluster formation in mitochondria (15). It is interesting to note that cysteinyl-tRNA synthetase was recently reported to have the activity to produce Cys-SSH (3).

Eberhardt *et al.* and Cortese-Krott *et al.* reported that the chemical interaction of H₂S with NO produces HNO and SSNO[−], respectively (10, 13). Both groups showed H₂S_n formation, but they did not consider their physiological relevance. Whereas Eberhardt *et al.* concluded that HNO is the chemical species responsible for activating the TRPA1-CGRP neuroendocrine signaling cascade and postulated that this pathway is essential for control of the general vascular tone, Dr. Kimura's group determined that H₂S₂ and H₂S₃ are the molecules that activate the TRPA1 channels of dorsal root ganglion neurons (52) (Fig. 5). Mustafa and Habara reported that H₂S_n must be produced by the interaction of H₂S and NO in mast cells (53). Molecules, which are produced from H₂S and NO and activate TRPA1 channels, and H₂S_n are degraded by cyanide (cyanolysis) and by reduction (52). In contrast, HNO is resistant to cyanolysis and SSNO[−] to reduction (10, 13). Considering these observations, Kimura's group suggested that the production of H₂S_n from H₂S and NO may be one of the mechanisms for the synergistic effects of both molecules on various tissues, including vascular smooth muscle relaxation (12, 21).

Other Achievements

In collaboration with Dr. Nagano, Dr. Urano, and Dr. Hanaoka, Dr. Kimura's group contributed to the development of H₂S- and polysulfide-fluorescence probes (52, 69, 83) as well as inhibitors specific to 3MST (19).

Current Position

Dr. Kimura is a specially appointed researcher, National Institute of Neuroscience, National Center of Neurology and Psychiatry. He was recently appointed as Professor and Faculty of Pharmaceutical Science at the Tokyo University of Science, Yamaguchi (renamed Sanyo-Onoda City University in 2018), Japan.

He received the Promoting Award from the Japanese Pharmacological Society (1988), Human Frontier Science Program (1990), First Award from National Institute of Health (1994), Alzheimer Scholar Award from Alzheimer Association (1994), JB Award from Japanese Biological Society (2010), Research Front Award from Thomson Reuters (2016), Highly Cited Researcher from Clarivate Analytics (2017), and President Award from the National Institute of Neuroscience, National Center of Neurology and Psychiatry (2018). He has served as a nominator of Japan Prize since 2008.

Acknowledgments

Dr. Hideo Kimura would like to thank Professor David Schubert for encouraging and supporting him to promote the study of hydrogen sulfide. He also thanks late Professor Yutaka Sakai, Professors Hideomi Fukuda, Hiroshi Saito, Toshiharu Nagatsu, Sadaaki Iwanaga, Keizo Inoue, and Eijiro Ozawa for

their suggestions and for supporting him. Dr. Kimura thanks all colleagues during their stay in the laboratories for their contribution to the study: Drs. Kazuho Abe, Norihiro Shibuya, Isao Ishii, Yasushi Enokido, Yasuo Nagai, Ken Umemura, Yoshinori Mikami, Kazuhiko Namekata, Mamiko Tsugane, Mari Ishigami, Shin Koike, Noriyuki Akahoshi, Yuriko Kobayashi, Makiko Tanaka, Yukiko Toyofuku, Ryo Miyamoto, and Yuka Kimura. He also thanks Professors Norio Matsuki, Makoto Suematsu, Yuki Ogasawara, Sachio Takashima, Noriyuki Nagahara, David Lefer, Kazuyuki Ishii, Tetsuo Nagano, Yasuteru Urano, Kenjiro Hanaoka, Toshihide Kimura, Yukiko Kaneko, Rumiko Hosoki, Akiko Ichinohe, Takeshi Kanaumi, and late Professor Ichiro Niki for their collaborations.

References

1. Abe K and Kimura H. The possible role of hydrogen sulfide as an endogenous neuromodulator. *J Neurosci* 16: 1066–1071, 1996.
2. Aizenman E, Lipton SA, and Loring RH. Selective modulation of NMDA responses by reduction and oxidation. *Neuron* 2: 1257–1263, 1989.
3. Akaike T, Ida T, Wei F-Y, Nishida M, Kumagai Y, Alam MM, Ihara H, Sawa T, Matsunaga T, Kasamatsu S, Nishimura A, Morita M, Tomizawa K, Nishimura A, Watanabe S, Inaba K, Shima H, Tanuma N, Jung M, Fujii S, Watanabe Y, Ohmuraya M, Nagy P, Feelisch M, Fukuto JM, and Motohashi H. Cysteinyl-tRNA synthetase governs cysteine polysulfidation and mitochondrial bioenergetics. *Nat Commun* 8: 1177, 2017.
4. Arnold WP, Mittal CK, Katsuki S, and Murad F. Nitric oxide activates guanylate cyclase and increases guanosine 3':5'-cyclic monophosphate levels in various tissue preparations. *Proc Natl Acad Sci U S A* 74: 3203–3207, 1977.
5. Braunstein AE, Goryachenkova EV, Tolosa EA, Willhardt IH, and Yefremova LL. Specificity and some other properties of liver serine sulphhydrase: evidence for its identity with cystathionine β -synthase. *Biochim Biophys Acta* 242: 247–260, 1971.
6. Bucci M, Papapetropoulos A, Vellecco V, Zhou Z, Zaid A, Giannogonas P, Cantalupo A, Dhayade S, Karalis KP, Wang R, Feil R, and Cirino G. cGMP-dependent protein kinase contributes to hydrogen sulfide-stimulated vasorelaxation. *PLoS One* 7: e53319, 2012.
7. Calvert JW, Jha S, Gundewar S, Elrod JW, Ramachandran A, Pattillo CB, Kevil CG, and Lefer DJ. Hydrogen sulfide mediates cardioprotection through Nrf2 signaling. *Circ Res* 105: 365–374, 2009.
8. Cavallini D, Mondovi B, De Marco C, and Scioscia-Santoro A. The mechanism of desulphhydration of cysteine. *Enzymologia* 24: 253–266, 1962.
9. Chen Q, Kimura H, and Schubert D. A novel mechanism for the regulation of amyloid precursor protein metabolism. *J Cell Biol* 158: 79–89, 2002.
10. Cortese-Krott MM, Kuhnle GGC, Dyson A, Fernandez BO, Grman M, DuMond JF, Barrow MP, McLeod G, Nakagawa H, Ondrias K, Nagy P, King SB, Saavedra JE, Keefer LK, Singer M, Kelm M, Butler AR, and Feelisch M. Key bioactive reaction products of the NO/H₂S interaction are S/N-hybrid species, polysulfides, and nitroxyl. *Proc Natl Acad Sci U S A* 112: E4651–E4660, 2015.
11. d'Emmanuele di Villa Bianca R, Sorrentino R, Maffia P, Mirone V, Imbimbo C, Fusco F, De Palma R, Ignarro LJ, and Cirino G. Hydrogen sulfide as a mediator of human corpus cavernosum smooth-muscle relaxation. *Proc Natl Acad Sci U S A* 106: 4513–4518, 2009.

12. Earley S. TRPA1 channels in the vasculature. *Br J Pharmacol* 167: 13–22, 2012.
13. Eberhardt M, Dux M, Namer B, Miljkovic J, Cordasic N, Will C, Kichko TI, de la Roche J, Fischer M, Suárez SA, Bikiel D, Dorsch K, Leffler A, Babes A, Lampert A, Lennerz JK, Jacobi J, Martí MA, Doctorovich F, Högestätt ED, Zygmunt PM, Ivanovic-Burmazovic I, Messlinger K, Reeh P, and Filipovic MR. H₂S and NO cooperatively regulate vascular tone by activating a neuroendocrine HNO-TRPA1-CGRP signalling pathway. *Nat Commun* 5: 4381, 2014.
14. Elrod JW, Calvert JW, Morrison J, Doeller JE, Kraus DW, Tao L, Jiao X, Scalia R, Kiss L, Szabo C, Kimura H, Chow CW, and Lefer DJ. Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function. *Proc Natl Acad Sci U S A* 104: 15560–15565, 2007.
15. Fräsdorf B, Radon C, and Leimkühler S. Characterization and interaction studies of two isoforms of the dual localized 3-mercaptopyruvate sulfurtransferase TUM1 from humans. *J Biol Chem* 289: 34543–34556, 2014.
16. Furchtgott RF and Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 288: 373–376, 1980.
17. Goodwin LR, Francom D, Dieken FP, Taylor JD, Warencia MW, Reiffenstein RJ, and Dowling G. Determination of sulfide in brain tissue by gas dialysis/ion chromatography: postmortem studies and two case reports. *J Anal Toxicol* 13: 105–109, 1989.
18. Greiner R, Pálinkás Z, Bässel K, Becher D, Antelmann H, Nagy P, and Dick TP. Polysulfides link H₂S to protein thiol oxidation. *Antioxid Redox Signal* 19: 1749–1765, 2013.
19. Hanaoka K, Sasakura K, Suwanai Y, Toma-Fukai S, Shimamoto K, Takano Y, Shibuya N, Terai T, Komatsu T, Ueno T, Ogasawara Y, Tsuchiya Y, Watanabe Y, Kimura H, Wang C, Uchiyama M, Kojima H, Okabe T, Urano Y, Shimizu T, and Nagano T. Discovery and mechanistic characterization of selective inhibitors of H₂S-producing enzyme: 3-mercaptopyruvate sulfurtransferase (3MST) targeting active-site cysteine persulfide. *Sci Rep* 7: 40227, 2017.
20. Hatakeyama Y, Takahashi K, Tominaga M, Kimura H, and Ohta T. Polysulfide evokes acute pain through the activation of nociceptive TRPA1 in mouse sensory neurons. *Mol Pain* 11: 24, 2015.
21. Hosoki R, Matsuki N, and Kimura H. The possible role of hydrogen sulfide as an endogenous smooth muscle relaxant in synergy with nitric oxide. *Biochem Biophys Res Commun* 237: 527–531, 1997.
22. Hylin JW and Wood JL. Enzymatic formation of polysulfides from mercaptopyruvate. *J Biol Chem* 234: 2141–2144, 1959.
23. Ignarro LJ, Adams JB, Horwitz PM, and Wood KS. Activation of soluble guanylate cyclase by NO-hemoproteins involves NO-heme exchange. Comparison of heme-containing and heme-deficient enzyme forms. *J Biol Chem* 261: 4997–5002, 1986.
24. Ishigami M, Hiraki K, Umemura K, Ogasawara Y, Ishii K, and Kimura H. A source of hydrogen sulfide and a mechanism of its release in the brain. *Antioxid Redox Signal* 11: 205–214, 2009.
25. Ishii I, Akahoshi N, Yu XN, Kobayashi Y, Namekata K, Komaki G, and Kimura H. Murine cystathionine gamma-lyase: complete cDNA and genomic sequences, promoter activity, tissue distribution and developmental expression. *Biochem J* 381: 113–123, 2004.
26. Kaneko Y, Kimura T, Taniguchi S, Souma M, Kojima Y, Kimura Y, Kimura H, and Niki I. Glucose-induced production of hydrogen sulfide may protect the pancreatic beta-cells from apoptotic cell death by high glucose. *FEBS Lett* 583: 377–382, 2009.
27. Kashiwa A, Yoshida H, Lee S, Paladino T, Liu Y, Chen Q, Dargusch R, Schubert D, and Kimura H. Isolation and characterization of novel presenilin binding protein. *J Neurochem* 75: 109–116, 2000.
28. Kimura H. Schwannoma-derived growth factor must be transported into the nucleus to exert its mitogenic activity. *Proc Natl Acad Sci U S A* 90: 2165–2169, 1993.
29. Kimura H. Physiological role of hydrogen sulfide and polysulfide in the central nervous system. *Neurochem Int* 63: 492–497, 2013.
30. Kimura H. Signaling molecules: hydrogen sulfide and polysulfide. *Antioxid Redox Signal* 22: 362–376, 2015.
31. Kimura H. 3MST produces redox regulators Cys-SSH and GSSH as well as signaling molecules H₂S and H₂S_n. *Atlas Sci* 22: 2018. <https://atlasofscience.org/3mst-produces-redox-regulators-cys-ssh-and-gssh-as-well-as-signaling-molecules-h2s-and-h2sn/>
32. Kimura H, Fischer WH, and Schubert D. Structure, expression and function of a schwannoma-derived growth factor. *Nature* 348: 257–260, 1990.
33. Kimura H, Okamoto K, and Sakai Y. Pharmacological evidence for L-aspartate as the neurotransmitter of cerebellar climbing fibers in the guinea-pig. *J Physiol* 365: 103–119, 1985.
34. Kimura H and Schubert D. Schwannoma-derived growth factor promotes the neuronal differentiation and survival of PC12 cells. *J Cell Biol* 116: 777–783, 1992.
35. Kimura H, Sogawa K, Sakai Y, and Fujii-Kuriyama Y. Alternative splicing mechanism in a cytochrome P-450 (P-450PB-1) gene generates the two mRNAs coding for proteins of different functions. *J Biol Chem* 264: 2338–2342, 1989.
36. Kimura H, Yoshioka H, Sogawa K, Sakai Y, and Fujii-Kuriyama Y. Complementary DNA cloning of cytochrome P-450s related to P-450(M-1) from the complementary DNA library of female rat livers. Predicted primary structures for P-450f, PB-1, and PB-1-related protein with a bizarre replacement block and their mode of transcriptional expression. *J Biol Chem* 263: 701–707, 1988.
37. Kimura Y, Dargusch R, Schubert D, and Kimura H. Hydrogen sulfide protects HT22 neuronal cells from oxidative stress. *Antioxid Redox Signal* 8: 661–670, 2006.
38. Kimura Y, Goto Y, and Kimura H. Hydrogen sulfide increases glutathione production and suppresses oxidative stress in mitochondria. *Antioxid Redox Signal* 12: 1–13, 2010.
39. Kimura Y and Kimura H. Hydrogen sulfide protects neurons from oxidative stress. *FASEB J* 18: 1165–1167, 2004.
40. Kimura Y, Koike S, Shibuya N, Lefer D, Ogasawara Y, and Kimura H. 3-Mercaptopyruvate sulfurtransferase produces potential redox regulators cysteine- and glutathione-persulfide (Cys-SSH and GSSH) together with signaling molecules H₂S₂, H₂S₃ and H₂S. *Sci Rep* 7: 10459, 2017.
41. Kimura Y, Mikami Y, Osumi K, Tsugane M, Oka J-I, and Kimura H. Polysulfides are possible H₂S-derived signaling molecules in rat brain. *FASEB J* 27: 2451–2457, 2013.
42. Kimura Y, Toyofuku Y, Koike S, Shibuya N, Nagahara N, Lefer D, Ogasawara Y, and Kimura H. Identification of H₂S₃ and H₂S produced by 3-mercaptopyruvate sulfurtransferase in the brain. *Sci Rep* 5: 14774, 2015.
43. Koike S, Kawamura K, Kimura Y, Shibuya N, Kimura H, and Ogasawara Y. Analysis of endogenous H₂S and H₂S_n

- in mouse brain by high-performance liquid chromatography with fluorescence and tandem mass spectrometric detection. *Free Radic Biol Med* 113: 355–362, 2017.
44. Koike S, Ogasawara Y, Shibuya N, Kimura H, and Ishii K. Polysulfide exerts a protective effect against cytotoxicity caused by t-buthylhydroperoxide through Nrf2 signaling in neuroblastoma cells. *FEBS Lett* 587: 3548–3555, 2013.
45. Krishnan N, Fu C, Pappin DJ, and Tonks NK. H₂S-induced sulfhydration of the phosphatase PTP1B and its role in the endoplasmic reticulum stress response. *Sci Signal* 4: ra86, 2011.
46. Lipsett MN, Norton JS, and Peterkofsky A. A requirement for beta-mercaptopyruvate in the in vitro thiolation of transfer ribonucleic acid. *Biochemistry* 6: 855–860, 1967.
47. Maines MD. Heme oxygenase: function, multiplicity, regulatory mechanisms, and clinical applications. *FASEB J* 2: 2557–2568, 1988.
48. Meister A, Fraser PE, and Tice SV. Enzymatic desulfuration of β -mercaptopyruvate to pyruvate. *J Biol Chem* 206: 561–575, 1954.
49. Mikami Y, Shibuya N, Kimura Y, Nagahara N, Yamada M, and Kimura H. Hydrogen sulfide protects the retina from light-induced degeneration by the modulation of Ca²⁺ influx. *J Biol Chem* 286: 39379–39386, 2011.
50. Mikami Y, Shibuya N, Kimura Y, Nagahara N, Ogasawara Y, and Kimura H. Thioredoxin and dihydrolipoic acid are endogenous reductants required for 3-mercaptopyruvate sulfurtransferase to produce hydrogen sulfide. *Biochem J* 439: 479–485, 2011.
51. Miller DL and Roth MB. Hydrogen sulfide increases thermotolerance and lifespan in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 104: 20618–20622, 2007.
52. Miyamoto R, Koike S, Takano Y, Shibuya N, Kimura Y, Hanaoka K, Urano Y, Ogasawara Y, and Kimura H. Polysulfides (H₂S_n) produced from the interaction of hydrogen sulfide (H₂S) and nitric oxide (NO) activate TRPA1 channels. *Sci Rep* 7: 45995, 2017.
53. Moustafa A and Habara Y. Cross talk between polysulfide and nitric oxide in rat peritoneal mast cells. *Am J Physiol Cell Physiol* 310: C894–C902, 2016.
54. Mustafa AK, Gadalla MM, Sen N, Kim S, Mu W, Gazi SK, Barrow RK, Yang G, Wang R, and Snyder SH. H₂S signals through protein S-sulfhydration. *Sci Signal* 2: ra72, 2009.
55. Nagahara N, Koike S, Nirasawa T, Kimura H, and Ogasawara Y. Alternative pathway of H₂S and polysulfides production from sulfated catalytic-cysteine of reaction intermediates of 3-mercaptopyruvate sulfurtransferase. *Biochem Biophys Res Commun* 496: 648–653, 2018.
56. Nagahara N and Nishino T. Role of amino acid residues in the active site of rat liver mercaptopyruvate sulfurtransferase. cDNA cloning, overexpression, and site-directed mutagenesis. *J Biol Chem* 271: 27395–27401, 1996.
57. Nagahara N, Yoshii T, Abe Y, and Matsumura T. Thioredoxin-dependent enzymatic activation of mercaptopyruvate sulfurtransferase. An intersubunit disulfide bond serves as a redox switch for activation. *J Biol Chem* 282: 1561–1569, 2007.
58. Nagai Y, Tsugane M, Oka J, and Kimura H. Hydrogen sulfide induces calcium waves in astrocytes. *FASEB J* 18: 557–559, 2004.
59. Nagai Y, Tsugane M, Oka J-I, and Kimura H. Polysulfides induce calcium waves in rat hippocampal astrocytes. *J Pharmacol Sci* 100: 200, 2006.
60. Namekata K, Enokido Y, Iwasawa K, and Kimura H. MOCA induces membrane spreading by activating Rac1. *J Biol Chem* 279: 14331–14337, 2004.
61. Namekata K, Harada C, Taya C, Guo X, Kimura H, Parada LF, and Harada T. Dock3 induces axonal outgrowth by stimulating membrane recruitment of the WAVE complex. *Proc Natl Acad Sci U S A* 107: 7586–7591, 2010.
62. Nishi N, Tanabe H, Oya H, Urushihara M, Miyanaka H, and Wada F. Identification of probasin-related antigen as cystathionine gamma-lyase by molecular cloning. *J Biol Chem* 269: 1015–1019, 1994.
63. Ogasawara Y, Ishii K, Togawa T, and Tanabe S. Determination of bound sulfur in serum by gas dialysis/high-performance liquid chromatography. *Anal Biochem* 215: 73–81, 1993.
64. Okamoto K, Kimura H, and Sakai Y. Evidence for taurine as an inhibitory neurotransmitter in cerebellar stellate interneurons: selective antagonism by TAG (6-aminomethyl-3-methyl-4H,1,2,4-benzothiadiazine-1,1-dioxide). *Brain Res* 265: 163–168, 1983.
65. Olson KR, Dombkowski RA, Russell MJ, Doellman MM, Head SK, Whitfield NL, and Madden JA. Hydrogen sulfide as an oxygen sensor/transducer in vertebrate hypoxic vasoconstriction and hypoxic vasodilation. *J Exp Biol* 209: 4011–4023, 2006.
66. Oosumi K, Tsugane M, Ishigami M, Nagai Y, Iwai T, Oka J-I, and Kimura H. Polysulfide activates TRP channels and increases intracellular Ca²⁺ in astrocytes. *Neurosci Res* 68: e109–e222, 2010.
67. Papapetropoulos A, Pyriochou A, Altaany Z, Yang G, Marazioti A, Zhou Z, Jeschke MG, Branski LK, Herndon DN, Wang R, and Szabó C. Hydrogen sulfide is an endogenous stimulator of angiogenesis. *Proc Natl Acad Sci U S A* 106: 21972–21977, 2009.
68. Peng YJ, Nanduri J, Raghuraman G, Souvannakitti D, Gadalla MM, Kumar GK, Snyder SH, and Prabhakar NR. H₂S mediates O₂ sensing in the carotid body. *Proc Natl Acad Sci U S A* 107: 10719–10724, 2010.
69. Sasakura K, Hanaoka K, Shibuya N, Mikami Y, Kimura Y, Komatsu T, Ueno T, Terai T, Kimura H, and Nagano T. Development of a highly selective fluorescence probe for hydrogen sulfide. *J Am Chem Soc* 133: 18003–18005, 2011.
70. Savage JC and Gould DH. Determination of sulfide in brain tissue and rumen fluid by ion-interaction reversed-phase high-performance liquid chromatography. *J Chromatogr* 526: 540–545, 1990.
71. Schubert D, Kimura H, LaCorbiere M, Vaughn J, Karr D, and Fischer W. Activin is a nerve cell survival protein. *Nature* 344: 868–870, 1990.
72. Sen N, Paul BD, Gadalla MM, Mustafa AK, Sen T, Xu R, Kim S, and Snyder SH. Hydrogen sulfide-linked sulfhydration of NF- κ B mediates its antiapoptotic actions. *Mol Cell* 45: 13–24, 2012.
73. Shatalin K, Shatalina E, Mironov A, and Nudler E. H₂S: a universal defense against antibiotics in bacteria. *Science* 334: 986–990, 2011.
74. Shibuya N, Koike S, Tanaka M, Ishigami-Yuasa M, Kimura Y, Ogasawara Y, Fukui K, Nagahara N, and Kimura H. A novel pathway for the production of hydrogen sulfide from D-cysteine in mammalian cells. *Nat Commun* 4: 1366, 2013.
75. Shibuya N, Mikami Y, Kimura Y, Nagahara N, and Kimura H. Vascular endothelium expresses 3-mercaptopyruvate sulfurtransferase and produces hydrogen sulfide. *J Biochem* 146: 623–626, 2009.

76. Shibuya N, Tanaka M, Yoshida M, Ogasawara Y, Togawa T, Ishii K, and Kimura H. 3-Mercaptopyruvate sulfurtransferase produces hydrogen sulfide and bound sulfane sulfur in the brain. *Antioxid Redox Signal* 11: 703–714, 2009.
77. Shigetomi E, Jackson-Weaver O, Huckstepp RT, O'Dell TJ, and Khakh BS. TRPA1 channels are regulators of astrocyte basal calcium levels and long-term potentiation via constitutive D-serine release. *J Neurosci* 33: 10143–10153, 2013.
78. Stevens CF and Wang Y. Reversal of long-term potentiation by inhibitors of haem oxygenase. *Nature* 364: 147–149, 1993.
79. Stipanuk MH and Beck PW. Characterization of the enzymic capacity for cysteine desulphhydration in liver and kidney of the rat. *Biochem J* 206: 267–277, 1982.
80. Stubbett D, Prysyazhna O, Rudyk O, Scotcher J, Burgoyne JR, and Eaton P. Protein kinase G 1 α oxidation paradoxically underlies blood pressure lowering by the reductant hydrogen sulfide. *Hypertension* 64: 1344–1351, 2014.
81. Swaroop M, Bradley K, Ohura T, Tahara T, Roper MD, Rosenberg LE, and Kraus JP. Rat cystathionine beta-synthase. Gene organization and alternative splicing. *J Biol Chem* 267: 11455–11461, 1992.
82. Szabo C. A timeline of hydrogen sulfide (H₂S) research: From environmental toxin to biological mediator. *Biochem Pharmacol* 149: 5–19, 2018.
83. Takano Y, Hanaoka K, Shimamoto K, Miyamoto R, Komatsu T, Ueno T, Terai T, Kimura H, Nagano T, and Urano Y. Development of a reversible fluorescent probe for reactive sulfur species, sulfane sulfur, and its biological application. *Chem Commun (Camb)* 53: 1064–1067, 2017.
84. Travis J. The rotten smell of memory: It's a gas. *Sci News* 149: 116, 1996.
85. Tripatara P, Patel NSA, Collino M, Gallicchio M, Kieswich J, Castiglia S, Benetti E, Stewart KN, Brown PAJ, Yaqoob MM, Fantozzi R, and Thiemermann C. Generation of endogenous hydrogen sulfide by cystathionine γ -lyase limits renal ischemia/reperfusion injury and dysfunction. *Lab Invest* 88: 1038–1048, 2008.
86. Verma A, Hirsch DJ, Glatt CE, Ronnett GV, and Snyder SH. Carbon monoxide: a putative neural messenger. *Science* 259: 381–384, 1993.
87. Wallace JL and Wang R. Hydrogen sulfide-based therapeutics: exploiting a unique but ubiquitous gasotransmitter. *Nat Rev Drug Discov* 14: 329–345, 2015.
88. Warenycia MW, Goodwin LR, Benishin CG, Reiffenstein RJ, Francom DM, Taylor JD, and Dieken FP. Acute hydrogen sulfide poisoning. Demonstration of selective uptake of sulfide by the brainstem by measurement of brain sulfide levels. *Biochem Pharmacol* 38: 973–981, 1989.
89. Warenycia MW, Goodwin LR, Francom DM, Dieken FP, Kombian SB, and Reiffenstein RJ. Dithiothreitol liberates non-acid labile sulfide from brain tissue of H₂S-poisoned animals. *Arch Toxicol* 64: 650–655, 1990.
90. Williams RA, Kelly SM, Mottram JC, and Coombs GH. 3-Mercaptopyruvate sulfurtransferase of *Leishmania* contains an unusual C-terminal extension and is involved in thior-
edoxin and antioxidant metabolism. *J Biol Chem* 278: 1480–1486, 2003.
91. Wong TW, Harris MA, and Jankowicz CA. Transfer ribonucleic acid sulfurtransferase isolated from rat cerebral hemispheres. *Biochemistry* 13: 2805–2812, 1974.
92. Yadav PK, Yamada K, Chiku T, Koutmos M, and Banerjee R. Structure and kinetic analysis of H₂S production by human mercaptopyruvate sulfurtransferase. *J Biol Chem* 288: 20002–20013, 2013.
93. Zanardo RCO, Brancalione V, Distrutti E, Fiorucci S, Cirino G, and Wallace JL. Hydrogen sulfide is an endogenous modulator of leukocyte-mediated inflammation. *FASEB J* 20: 2118–2120, 2006.
94. Zhao W, Zhang J, Lu Y, and Wang R. The vasorelaxant effect of H₂S as a novel endogenous gaseous K_{ATP} channel opener. *EMBO J* 20: 6008–6016, 2001.
95. Zhuo M, Small SA, Kandel ER, and Hawkins RD. Nitric oxide and carbon monoxide produce activity-dependent long-term synaptic enhancement in hippocampus. *Science* 260: 1946–1950, 1993.

Address correspondence to:

Dr. David Lefer

CV Center of Excellence

Louisiana State University Health Sciences Center

New Orleans, LA 70112-2784

E-mail: dlefe1@lsuhsc.edu

Date of first submission to ARS Central, July 23, 2018; date of final revised submission, September 18, 2018; date of acceptance, September 18, 2018.

Abbreviations Used

3MP = 3-mercaptopyruvate
 3MST = 3-mercaptopyruvate sulfurtransferase
 AMPA = α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
 CBS = cystathionine β -synthase
 CSE = cystathionine γ -lyase
 DAO = D-amino acid oxidase
 Dock3 = dedicator of cytokines 3
 DTT = dithiothreitol
 GSH = glutathione
 H₂S = hydrogen sulfide
 Keap1 = kelch-like ECH-associated protein 1
 LTP = long-term potentiation
 NMDA = N-methyl D-aspartate
 NO = nitric oxide
 Nrf2 = nuclear factor-like 2
 PBP = presenilin-binding protein
 SP1 = specific protein 1
 TNF α = tumor necrosis factor alpha
 TRPA1 = transient receptor potential ankyrin 1