REVIEW

Open Access

Review: The Role of Cannabinoids on Esophageal Function—What We Know Thus Far

Jonathan Gotfried,^{1,2} Rahul Kataria,^{1,2} and Ron Schey^{1,2,*}

Abstract

The endocannabinoid system (ECS) primarily consists of cannabinoid receptors (CBRs), endogenous ligands, and enzymes for endocannabinoid biosynthesis and inactivation. Although the presence of CBRs, both CB₁ and CB₂, as well as a third receptor (G-protein receptor 55 [GPR55]), has been established in the gastrointestinal (GI) tract, few studies have focused on the role of cannabinoids on esophageal function. To date, studies have shown their effect on GI motility, inflammation and immunity, intestinal and gastric acid secretion, nociception and emesis pathways, and appetite control. Given the varying and sometimes limited efficacy of current medical therapies for diseases of the esophagus, further understanding and investigation into the interplay of the ECS on esophageal health and disease may present new therapeutic modalities that may help advance current treatment options. In this brief review, the current understanding of the ECS role in various esophageal functions and disorders is presented.

Keywords: cannabinoid receptors; endocannabinoid; esophagus; medicinal marijuana; motility

Overview of Endocannabinoid System

The endocannabinoid system (ECS) primarily consists of cannabinoid receptors (CBRs), endogenous ligands, and enzymes for endocannabinoid biosynthesis and inactivation.¹ The ECS plays an important role in regulation of synaptic transmission in the central and enteric nervous systems (ENS) through both excitatory and inhibitory effects, mediating a variety of physiological processes including pain sensation and modulation, motor function, inflammation, and immunity.²

CBRs belong to the superfamily of G-protein-coupled receptors and are expressed in two main forms, CB₁ and CB₂.³ CB₁ is mainly expressed in central and peripheral neurons, including the ENS, whereas CB₂ is mostly expressed by inflammatory/immune cells.^{4–6} The ubiquitous distribution of CBRs in the ENS and gastrointestinal (GI) tract highlights its role in GI health and disease including motility, inflammation and immunity, intestinal and gastric acid secretion, nociception and emesis pathways, and appetite control^{7–9} (Table 1).

The CB₁ receptor plays a role in intestinal motility by attenuating both large and small bowel muscle tone when activated.^{10–13} Within the upper GI tract, activation of CB₁ decreases intragastric pressure and delays gastric emptying through inhibition of excitatory neurons.^{14,15} As demonstrated in rodent models, CB₁ receptors play a role in energy regulation by driving consumption of food high in dietary fat.¹⁶ After sham feeding in rats with high-fat foods, upregulation of CB₁ receptors was found in rat small intestine, leading to inhibition of neural signaling events of satiety to suppress feeding. These findings suggest that CB₁ acts through a positive feedback loop after dietary fat exposure to stimulate further consumption of higher energy foods. In another study, pharmacological inhibition of CB₁ led to decreased amount of refeeding in rats after food deprivation, highlighting CB₁ receptor's role in appetite regulation.¹⁷ In addition, CB₁ was found to be upregulated in rats fed a high fat, high caloric Western diet (WD). It was also suggested to play a role in hyperphagia

¹Department of Gastroenterology, Temple University Hospital, Philadelphia, Pennsylvania.

²Lewis Katz School of Medicine, Temple University, Philadelphia, Pennsylvania.

*Address correspondence to: Ron Schey, MD, FACP, Department of Gastroenterology, Temple University Hospital, Philadelphia, PA 19140, E-mail: ron.schey@tuhs.temple.edu

Table 1. Known Functions of the Endocannabinoid System Along the Upper Alimentary Tract

Function	Brief summary	Related studies
Appetite regulation	1. Stimulates appetite, especially high energy, fatty foods 2. Induces hyperphagia, potential target for obesity therapy, generates hunger signal	Argueta and DiPatrizio ¹⁸ DiPatrizio et al. ¹⁷ DiPatrizio et al. ¹⁶
Nociception/emesis	1. Stimulation of CB ₁ receptor in CNS leads to nausea, therapeutic target for antiemetic effect of exogenous cannabinoid therapy	Van Sickle et al. ¹⁹ Rock and Parker ⁷¹
Motility	1. Multiple receptors (CB ₁ and CB ₂) thought to influence contractile and relaxant forces in stomach 2. Modulates intestinal propulsion, GPR55 inhibits whole gut transit time, modulation of cholinergic and vagal stimulation of upper GI tract 3. Enhances gut motility in setting of inflammation	Hornby and Prouty ¹⁰ Yuece et al. ¹² Storr et al. ²¹
Gastric acid and intestinal secretions	1. Exhibits antisecretory effects on gastric acid 2. Implicated in mitigating inflammation and mucosal damage in GERD 3. Effect on transient lower esophageal relaxation	Izzo et al. ²⁴ Yang et al. ²⁶ Li et al. ²⁷ Adami et al. ¹⁵ Calabrese et al. ²⁵
Inflammation and immunity	1. Anti-inflammatory effects in esophageal reflux disease 2. CB ₂ receptor downregulates inflammation and associated hypermotility in disease state	Lehmann et al. ³³ Calabrese et al. ²⁵ Izzo ²³
Analgesia	1. Increases pain threshold	Clapper et al. ⁶⁵ Malik et al. ⁶⁶

CNS, central nervous system; GERD, gastroesophageal reflux disease; GI, gastrointestinal; GPR55, G-protein receptor 55.

after antagonism of CB₁ with an inhibitor led to decreased food intake in rats fed a WD for 60 days.¹⁸

CB₂ receptors located in the central nervous system (CNS) have been shown to play a role in the emetic pathway; however, the receptor has also been found in inflammatory tissue and immune cells (plasma cells and macrophages) throughout the GI tract.^{19–22} CB₂ is expressed in the GI tract but less extensively than CB₁ receptors.²³ Evidence that upregulation of CB₂ receptors occurs in patients with inflammatory bowel disease suggests that these receptors play a critical role in colonic inflammation.⁷ Distinct from CB₁ receptors effect on motility, CB₂ receptors may further regulate motility in pathophysiological states with its expression being upregulated in inflammatory disease states.^{5,24}

Presence of CBRs in human esophageal epithelium was first demonstrated in a study comparing patients with nonerosive esophageal reflux disease (NERD) and erosive esophageal reflux disease (ERD) to normal controls.²⁵ The authors found increased expression of CB₁ mRNA in NERD patients compared with erosive esophagitis, but overall less expression compared with normal controls. Interestingly, CB₁ protein expression was similar to patients with ERD, whereas NERD patients showed increased CB₁ receptor levels when compared with healthy controls (Fig. 1). Although other GI inflammatory conditions have increased CB₁ activity, there may be contribution of the inflammatory microenvironment that alters CB₁ gene expression.²⁴

Recently, a potential third CBR has been identified with implications in the GI tract. G-protein receptor 55 (GPR55) shares 13–15% sequence homology with the CB₁ and CB₂ receptors and responds to a multitude of endogenous and exogenous cannabinoid ligands as well as several lipids.²⁶ A recent study found activation of GPR55 after administration of a synthetic agonist slowed down whole-gut transit in mice *in vivo*, suggesting GPR55 may be involved in the regulation of gut motility.²⁷

Endocannabinoids are endogenously produced ligands that exert effects on CBRs. The major ligands, anandamide (AEA) and 2-arachidonylglycerol (2-AG), play a role in maintaining GI homeostasis and have been found at increased levels in GI disease states, including celiac disease, diverticulosis, and colorectal cancer.^{28–30} Exogenous cannabinoids, both plant-derived phytocannabinoids (cannabis sativa) and synthetic cannabinoids, also directly activate CBRs. They have been shown to play a role in both GI pathophysiology (e.g., cannabinoid hyperemesis syndrome) and therapies (e.g., antiemetics and appetite stimulant).³¹

Although both CB₁ and CB₂ receptors have been found in the esophagus, few studies have focused on the role of cannabinoids on esophageal function.²¹ Thus far, the role of GPR55 in esophageal motility has not been established. Given the varying and sometimes limited efficacy of current medical therapies for diseases of the esophagus, further understanding and

FIG. 1. Immunostaining of CB₁ receptor in histological sections of esophageal mucosa. Healthy subjects **(a)** show a weak positive staining localized in mature squamous cells (black arrow) and in connective papillae (red arrows). NERD patients **(b)** show CB₁ receptor expression in mature squamous cells (black arrow), in squamous cells (blue arrow), and in connective papillae (red arrow). ERD patients **(c)** show CB₁ positivity only in mature squamous cells (black arrow) and in squamous cells (blue arrows), whereas connective papillae appear negative (red arrow). CB₁ staining disappeared in esophageal mucosa **(d)** when CB₁ blocking peptide was incubated with CB₁ antibody. ERD, erosive esophageal reflux disease; NERD, nonerosive esophageal reflux disease. Reprinted with permission from Calabrese et al.²⁵

investigation into the interplay of the ECS on esophageal health and disease may present new therapeutic modalities that may help advance current treatment options. In this review, the current understanding of the role of ECS in various esophageal functions and disorders is presented.

Swallowing Mechanism

The swallowing reflex is an important mechanism in controlling acid exposure in the esophagus, whereby increased swallowing decreases stasis and reduces acidic and nonacidic reflux. The role of the swallowing reflex in reducing reflux has been demonstrated after observation that supine sleeping patients experienced a decreased spontaneous swallowing reflex during 24-h pH monitoring, leading to increased acid and nonacid

exposure.³² CB₁ receptor activation has been shown to decrease spontaneous swallows in human and animal studies. One study on dogs found administration of CB₁ agonist suppressed spontaneous swallowing in a dose-dependent manner with an >80% decrease in spontaneous swallows at high doses (57 nmol/kg).³³ The authors noted that although administration of CB₁ agonist decreased transient lower esophageal sphincter relaxations (TLESRs) thereby decreasing reflux events, the concomitant decrease in spontaneous swallows and clearance of refluxate potentially limit the benefits of decreasing TLESRs. The mechanism of action of CBRs on swallowing mechanism is likely centrally located through endogenous cannabinoid action, modifying synaptic neurotransmitter release.³⁴ The swallowing reflex, evoked by repetitive electrical stimulation of the superior

laryngeal nerve in rats, was analyzed with and without combinations of both CB₁ and CB₂ agonists and antagonists. CB₁ receptor antagonist injected directly into the nuclear tractus solitarius blocked the action of intravenous-administered CB₁ agonist. The only study in humans to date found administration of the combination CB₁/CB₂ agonist Δ 9-THC resulted in a significant reduction in the number of swallows wherein high doses (20 mg) led to a reduction in spontaneous swallows by 50%.³⁵

Esophageal Motility

CBRs, specifically CB₁, play a role in GI motility and have mainly been studied in the small intestine and colon.^{8,10} Only two studies have evaluated the effect of ECS on esophageal motility in humans.^{35,36} Administration of Δ 9-THC decreased basal lower esophageal sphincter (LES) pressure in a nondose-dependent manner, with onset occurring 45 min after meal ingestion, maximal effect around 100 min, followed by slow recovery.³⁵ This was confirmed in another human study that administration of rimonabant, a CB₁ receptor antagonist, increased postprandial LES pressures in the first and second postprandial hours.³⁶

Previous studies in animal models did not demonstrate CBR action influenced esophageal motility. Lehman et al. demonstrated that CBR agonism did not influence the extent of LES relaxation. Similarly, CBR antagonists had no influence on esophageal peristalsis.³³ A recent case report suggested that exogenous phytocannabinoid use improved symptoms of dysphagia in a patient with manometric findings, consistent with type 3 achalasia.³⁷ Others have suggested that exogenous cannabinoid use, like chronic opiate use, may produce motor findings similar to type 3 achalasia.³⁸ The conflicting data support the need for further study of endo- and exogenous cannabinoids on esophageal motility.³⁹

Gastroesophageal Reflux Disease and TLESRs

TLESRs are the predominant mechanism seen in gastroesophageal reflux disease (GERD). TLESRs occur after gastric stimuli, mainly distension, to relieve counteracting gastric pressure on the LES.^{40,41} The LES response to gastric distension is vagally mediated through communication of the LES and crural diaphragm with afferent gastric pathways, brainstem integrative centers (nucleus tractus solitarius and dorsal motor nucleus), and efferent inhibitory pathways.⁴¹ The action of CBRs on TLESRs and GERD was first demonstrated in animal models after the observation that administration of a CB₁

agonist (WIN55,212-2) reduced the rate of TLESRs without altering the TLESRs latency or esophageal peristalsis.³³ The authors concluded that the action of the CBRs on TLESRs most likely occurred via central pathways because no effect on the extent of LES relaxation was seen.^{33,42,43} Absence of CB₁ receptors in preganglionic vagal motor neurons projecting to the gastric fundus or LES further supports the assumption that CB₁ does not directly participate in vagal motor output modulation and rather relies on central activation.⁴²

Interestingly, the use of a CB₁ receptor antagonist (rimonabant) in healthy human subjects enhanced postprandial LES pressure but unexpectedly decreased TLESRs.³⁶ This contradicts earlier data in an animal model with dogs that showed rimonabant enhanced the rate of TLESRs and reflux events that received acidified meal and intragastric air insufflation.³⁶ The authors attributed this discrepancy to different dosage, bioavailability, or interspecies differences of rimonabant. A possible explanation for the similar results from administering CBR antagonist and agonists may be a result of rimonabant exerting potent CB₁ receptor-independent pharmacological effects.⁴⁴

Gastric accommodation may affect TLESRs and play an important role in development of GERD.⁴⁵ Previous studies evaluated the effect of CB₁ receptors on pain sensation during intragastric balloon distension, however, they did not find modulation of CB₁ receptors with rimonabant-influenced mechanosensitivity of the proximal stomach.⁴⁶ It should be noted that baclofen has been shown to decrease TLESRs rate and increase basal LES pressure while not affecting meal-induced fundic accommodation, suggesting that the mechanism may not be as simple as binary, an observation previously suggested by Lehmann et al.³³ Other studies have suggested that using Δ 9-THC inhibits gastric insufflation-induced LES relaxations in the decerebrate ferret, highlighting the need for further clarification of the relationship between gastric stimulation and TLESRs.¹⁰

The effect of CBRs on gastric emptying has been studied in humans although the data are limited and conflicting.^{48–51} Although there may be some influence of CBRs, gastric emptying itself does not necessarily correlate with fundus accommodation and activation of TLESRs.

The presence of CBRs in the esophagus, specifically those affecting TLESRs, offers a potential therapeutic target for treating GERD. Using Δ 9-THC, Beaumont et al. demonstrated decreased rate of TLESRs in healthy volunteers who received 10 and 20 mg of Δ 9-THC on

three occasions a week apart.³⁵ Delta(9)-THC significantly reduced the number of TLESRs and caused a nonsignificant reduction of acid reflux episodes in the first postprandial hour. In addition, lower esophageal sphincter pressure and swallowing were significantly reduced by $\Delta(9)$ -THC.³⁵ However, in the high dose of $\Delta 9$ -THC (20 mg) group, central activity led to increased nausea and vomiting. Centrally acting CB₁ receptor agonists produce the psychotropic effects and, therefore, selective targeting of peripheral CB₁ receptors is necessary for effective therapy, with recent efforts to develop higher potency and effective CB₁ agonists.^{52,53} Previous studies have looked at using CB₁ antagonists; however, their therapeutic use is also limited by its side effect of major depression.⁵⁴

Newer methodologies for understanding the pharmacodynamics and pharmacokinetics of medication effects on TLESRs have been developed that may provide more accurate modeling of drug concentrations and their effects.⁵⁵

Bronchoconstriction

In patients with a chronic cough where lung disease, environmental exposure, and medications have been excluded, GERD is often a major cause.⁵⁶ The two main mechanisms implicated in reflux-related cough are microaspiration of refluxate and stimulation of vagal innervation of the esophagus, leading to esophagobronchial reflex arc. The CB₂ receptor has been shown to play a role in inhibiting bronchoconstriction and microvascular leakage in reflux models using guinea pigs. After infusion of intraesophageal HCl, the inhibitory effect of bronchoconstriction by a CB₁ agonist (WIN55,212-2) was blocked after administration of a CB₂ antagonist (SR 144528) but not after CB₁ receptor antagonist, demonstrating CB₂ receptor's role as a downregulatory mechanism of sensory nerve activation.⁵⁷ CB₂ receptor activation in models of direct airway damage has been shown to play an active role in reducing inflammation, potentially supporting protective role of CB₂ in microaspiration of refluxate.⁵⁸

Visceral Hypersensitivity and Pain

Functional esophageal disorders are defined as the presence of esophageal symptoms in the absence of structural, inflammatory, or motor abnormalities.⁵⁹ Proposed pathophysiological mechanisms include alterations in neural processing between peripheral triggers and central perception of esophageal symptoms.⁶⁰ CBRs likely play a role in pain sensation and modulation through visceral

antinociceptive action.⁶¹ Previous studies have demonstrated an analgesic effect of cannabinoids in animal models through both CB₁ and CB₂ activation.^{62–64} In one important study, upregulation of the endocannabinoid AEA through inhibition of its degrading enzyme led to an attenuated behavioral response to noxious stimuli in rodents.⁶⁵ This suggests a central role of CB₁ receptors in mitigating pain-related inputs to the CNS.

There are few studies of visceral sensitivity in the esophagus and the effects of CBR modulation. In a randomized, placebo controlled trial, administration of dronabinol for 1 month increased the threshold for first sensation, pain frequency, and intensity of pain during an esophageal balloon distention test.⁶⁶ Notably, anxiety and depression indices were unchanged after dronabinol administration. The mainstay of functional esophageal disorders relies on tricyclic antidepressants, selective serotonin reuptake inhibitors, and serotonin norepinephrine reuptake inhibitors, all of which have been found to influence psychiatric parameters, highlighting a possible difference in mechanism of action.⁶⁰

Role in Esophageal Neoplasm

Cannabinoids inhibit tumor cell growth and induce apoptosis by modulating cell signaling pathways.⁶⁷ Decreased frequency of CNR-1 gene, the gene encoding CB₁ receptor, expression was found in tissue of esophageal cancer patients (10.8%) compared with controls (60.0%), and its presence is considered an independent predictor of survival.⁶⁸ However, recently a multivariate analysis revealed that CB₁ receptor overexpression was independently associated with poor prognosis ($p=0.019$). Biological analysis of CB₁ receptor overexpression using esophageal squamous carcinoma cell lines revealed that CB₁ receptor activation appeared to promote cell proliferation and invasion.⁶⁹ Thus, the role of CB₁ receptor expression in tumorigenesis needs to be further evaluated.

Conclusion

A more thorough understanding of the ECS in esophageal function and disease is needed. Recently, there has been a push to legalize cannabis for both medicinal and recreational use. There have also been reports of increasing use of both plant-derived cannabis and synthetic cannabinoids in the United States.⁷⁰ This may allow additional studies to be added to the few human studies to date on the effect of cannabinoid use in humans. Further study in this area is imperative for the development of future therapeutic potential of utilizing the ECS.

Authors' Contributions

J.G. was involved in literature review and is the primary author. R.K. was involved in literature review. R.S. was involved in study conceptualization and reviewed the article.

Author Disclosure Statement

No competing financial interests exist.

References

- De Petrocellis L, Di Marzo V. An introduction to the endocannabinoid system: from the early to the latest concepts. *Best Pract Res Clin Endocrinol Metab.* 2009;23:1–15.
- Aizpurua-Olaizola O, Elezgarai I, Rico-Barrio I, et al. Targeting the endocannabinoid system: future therapeutic strategies. *Drug Discov Today.* 2017;22:105–110.
- Coutts AA, Irving AJ, Mackie K, et al. Localisation of cannabinoid CB(1) receptor immunoreactivity in the guinea pig and rat myenteric plexus. *J Comp Neurol.* 2002;448:410–422.
- Matsuda LA, Bonner TI, Lolait SJ. Localization of cannabinoid receptor mRNA in rat brain. *J Comp Neurol.* 1993;327:535–550.
- Duncan M, Mouihate A, Mackie K, et al. Cannabinoid CB2 receptors in the enteric nervous system modulate gastrointestinal contractility in lipopolysaccharide-treated rats. *Am J Physiol Gastrointest Liver Physiol.* 2008;295:G78–G87.
- Coutts AA, Izzo AA. The gastrointestinal pharmacology of cannabinoids: an update. *Curr Opin Pharmacol.* 2004;4:572–579.
- Wright K, Rooney N, Feeney M, et al. Differential expression of cannabinoid receptors in the human colon: cannabinoids promote epithelial wound healing. *Gastroenterology.* 2005;129:437–453.
- Pertwee RG. Cannabinoids and the gastrointestinal tract. *Gut.* 2001;48:859–867.
- Boesmans W, Ameloot K, van den Abbeel V, et al. Cannabinoid receptor 1 signalling dampens activity and mitochondrial transport in networks of enteric neurones. *Neurogastroenterol Motil.* 2009;21:958–e77.
- Hornby PJ, Prouty SM. Involvement of cannabinoid receptors in gut motility and visceral perception. *Br J Pharmacol.* 2004;141:1335–1345.
- Pinto L, Capasso R, Di Carlo G, et al. Endocannabinoids and the gut. *Prostaglandins Leukot Essent Fatty Acids.* 2002;66:333–341.
- Yuece B, Sibaev A, Broedl UC, et al. Cannabinoid type 1 receptor modulates intestinal propulsion by an attenuation of intestinal motor responses within the myenteric part of the peristaltic reflex. *Neurogastroenterol Motil.* 2007;19:744–753.
- Izzo AA, Mascolo N, Pinto L, et al. The role of cannabinoid receptors in intestinal motility, defaecation and diarrhoea in rats. *Eur J Pharmacol.* 1999;384:37–42.
- Storr MA, Yuece B, Andrews CN, et al. The role of the endocannabinoid system in the pathophysiology and treatment of irritable bowel syndrome. *Neurogastroenterol Motil.* 2008;20:857–868.
- Adami M, Zamfirova R, Sotirov E, et al. Gastric antisecretory effects of synthetic cannabinoids after central or peripheral administration in the rat. *Brain Res Bull.* 2004;64:357–361.
- DiPatrizio NV, Astarita G, Schwartz G, et al. Endocannabinoid signal in the gut controls dietary fat intake. *Proc Natl Acad Sci U S A.* 2011;108:12904–12908.
- DiPatrizio NV, Igarashi M, Narayanaswami V, et al. Fasting stimulates 2-AG biosynthesis in the small intestine: role of cholinergic pathways. *Am J Physiol Regul Integr Comp Physiol.* 2015;309:R805–R813.
- Argueta DA, DiPatrizio NV. Peripheral endocannabinoid signaling controls hyperphagia in western diet-induced obesity. *Physiol Behav.* 2017;171:32–39.
- Van Sickle MD, Duncan M, Kingsley PJ, et al. Identification and functional characterization of brainstem cannabinoid CB2 receptors. *Science.* 2005;310:329–332.
- Onaivi ES. Neuropsychobiological evidence for the functional presence and expression of cannabinoid CB2 receptors in the brain. *Neuropsychobiology.* 2006;54:231–246.
- Storr M, Gaffal E, Saur D, et al. Effect of cannabinoids on neural transmission in rat gastric fundus. *Can J Physiol Pharmacol.* 2002;80:67–76.
- Berdyshev EV. Cannabinoid receptors and the regulation of immune response. *Chem Phys Lipids.* 2000;108:169–190.
- Izzo AA. The cannabinoid CB(2) receptor: a good friend in the gut. *Neurogastroenterol Motil.* 2007;19:704–708.
- Izzo AA, Fezza F, Capasso R, et al. Cannabinoid CB1-receptor mediated regulation of gastrointestinal motility in mice in a model of intestinal inflammation. *Br J Pharmacol.* 2001;134:563–570.
- Calabrese C, Spisni E, Liguori G, et al. Potential role of the cannabinoid receptor CB in the pathogenesis of erosive and non-erosive gastro-oesophageal reflux disease. *Aliment Pharmacol Ther.* 2010;32:603–611.
- Yang H, Zhou J, Lehmann C. GPR55—a putative “type 3” cannabinoid receptor in inflammation. *J Basic Clin Physiol Pharmacol.* 2016;27:297–302.
- Li K, Fichna J, Schicho R, et al. A role for O-1602 and G protein-coupled receptor GPR55 in the control of colonic motility in mice. *Neuropharmacology.* 2013;71:255–263.
- D’Argenio G, Petrosino S, Gianfrani C, et al. Overactivity of the intestinal endocannabinoid system in celiac disease and in methotrexate-treated rats. *J Mol Med (Berl).* 2007;85:523–530.
- Guagnini F, Valenti M, Mukenge S, et al. Neural contractions in colonic strips from patients with diverticular disease: role of endocannabinoids and substance P. *Gut.* 2006;55:946–953.
- Ligresti A, Bisogno T, Matias I, et al. Possible endocannabinoid control of colorectal cancer growth. *Gastroenterology.* 2003;125:677–687.
- Pertwee RG. Emerging strategies for exploiting cannabinoid receptor agonists as medicines. *Br J Pharmacol.* 2009;156:397–411.
- Dickman R, Shapiro M, Malagon IB, et al. Assessment of 24-h oesophageal pH monitoring should be divided to awake and asleep rather than upright and supine time periods. *Neurogastroenterol Motil.* 2007;19:709–715.
- Lehmann A, Blackshaw LA, Branden L, et al. Cannabinoid receptor agonism inhibits transient lower esophageal sphincter relaxations and reflux in dogs. *Gastroenterology.* 2002;123:1129–1134.
- Mostafaez RM, Zakir HM, Takatsuji H, et al. Cannabinoids facilitate the swallowing reflex elicited by the superior laryngeal nerve stimulation in rats. *PLoS One.* 2012;7:e50703.
- Beaumont H, Jensen J, Carlsson A, et al. Effect of delta9-tetrahydrocannabinol, a cannabinoid receptor agonist, on the triggering of transient lower esophageal sphincter relaxations in dogs and humans. *Br J Pharmacol.* 2009;156:153–162.
- Scarpellini E, Blondeau K, Boeckstaens V, et al. Effect of rimonabant on oesophageal motor function in man. *Aliment Pharmacol Ther.* 2011;33:730–737.
- Luquians A, Lourenco N, Benyamina A, et al. Self-medication of achalasia with cannabis, complicated by a cannabis use disorder. *World J Gastroenterol.* 2015;21:6381–6383.
- Gyawali CP. Achalasia: new perspectives on an old disease. *Neurogastroenterol Motil.* 2016;28:4–11.
- Malik Z, Baik D, Schey R. The role of cannabinoids in regulation of nausea and vomiting, and visceral pain. *Curr Gastroenterol Rep.* 2015;17:429.
- Dent J, Dodds WJ, Friedman RH, et al. Mechanism of gastroesophageal reflux in recumbent asymptomatic human subjects. *J Clin Invest.* 1980;65:256–267.
- Mittal RK, Holloway RH, Penagini R, et al. Transient lower esophageal sphincter relaxation. *Gastroenterology.* 1995;109:601–610.
- Partosoedarso ER, Abrahams TP, Scullion RT, et al. Cannabinoid1 receptor in the dorsal vagal complex modulates lower esophageal sphincter relaxation in ferrets. *J Physiol.* 2003;550(Pt 1):149–158.
- Van Sickle MD, Oland LD, Ho W, et al. Cannabinoids inhibit emesis through CB1 receptors in the brainstem of the ferret. *Gastroenterology.* 2001;121:767–774.
- Bifulco M, Grimaldi C, Gazziero P, et al. Rimobant: just an antiobesity drug? current evidence on its pleiotropic effects. *Mol Pharmacol.* 2007;71:1445–1456.
- Pauwels A, Altan E, Tack J. The gastric accommodation response to meal intake determines the occurrence of transient lower esophageal sphincter relaxations and reflux events in patients with gastro-oesophageal reflux disease. *Neurogastroenterol Motil.* 2014;26:581–588.
- Ameloot K, Janssen P, Scarpellini E, et al. Endocannabinoid control of gastric sensorimotor function in man. *Aliment Pharmacol Ther.* 2010;31:1123–1131.
- Lee KJ, Vos R, Janssens J, et al. Differential effects of baclofen on lower esophageal sphincter pressure and proximal gastric motility in humans. *Aliment Pharmacol Ther.* 2003;18:199–207.

48. McCallum RW, Soykan I, Sridhar KR, et al. Delta-9-tetrahydrocannabinol delays the gastric emptying of solid food in humans: a double-blind, randomized study. *Aliment Pharmacol Ther.* 1999;13:77–80.
49. Bateman DN. Delta-9-tetrahydrocannabinol and gastric emptying. *Br J Clin Pharmacol.* 1983;15:749–751.
50. Esfandiyari T, Camilleri M, Ferber I, et al. Effect of a cannabinoid agonist on gastrointestinal transit and postprandial satiation in healthy human subjects: a randomized, placebo-controlled study. *Neurogastroenterol Motil.* 2006;18:831–838.
51. Finizia C, Lundell L, Cange L, et al. The effect of cisapride on oesophageal motility and lower sphincter function in patients with gastro-oesophageal reflux disease. *Eur J Gastroenterol Hepatol.* 2002;14:9–14.
52. Ameri A. The effects of cannabinoids on the brain. *Prog Neurobiol.* 1999;58:315–348.
53. Plowright AT, Nilsson K, Antonsson M, et al. Discovery of agonists of cannabinoid receptor 1 with restricted central nervous system penetration aimed for treatment of gastroesophageal reflux disease. *J Med Chem.* 2013;56:220–240.
54. Blondeau K, Boecxstaens V, Rommel N, et al. Baclofen improves symptoms and reduces postprandial flow events in patients with rumination and supragastric belching. *Clin Gastroenterol Hepatol.* 2012;10:379–384.
55. Plan EL, Ma G, Nagard M, et al. Transient lower esophageal sphincter relaxation pharmacokinetic-pharmacodynamic modeling: count model and repeated time-to-event model. *J Pharmacol Exp Ther.* 2011;339: 878–885.
56. Palombini BC, Villanova CA, Araujo E, et al. A pathogenic triad in chronic cough: asthma, postnasal drip syndrome, and gastroesophageal reflux disease. *Chest.* 1999;116:279–284.
57. Cui YY, D'Agostino B, Risse PA, et al. Cannabinoid CB(2) receptor activation prevents bronchoconstriction and airway oedema in a model of gastro-oesophageal reflux. *Eur J Pharmacol.* 2007;573: 206–213.
58. Barnes PJ. The problem of cough and development of novel antitussives. *Pulm Pharmacol Ther.* 2007;20:416–422.
59. Galmiche JP, Clouse RE, Balint A, et al. Functional esophageal disorders. *Gastroenterology.* 2006;130:1459–1465.
60. Aziz Q, Fass R, Gyawali CP, et al. Functional esophageal disorders. *Gastroenterology.* 2016;130:1459–1465.
61. Beltramo M. Cannabinoid type 2 receptor as a target for chronic—pain. *Mini Rev Med Chem.* 2009;9:11–25.
62. Wright KL, Duncan M, Sharkey KA. Cannabinoid CB2 receptors in the gastrointestinal tract: a regulatory system in states of inflammation. *Br J Pharmacol.* 2008;153:263–270.
63. Kikuchi A, Ohashi K, Sugie Y, et al. Pharmacological evaluation of a novel cannabinoid 2 (CB2) ligand, PF-03550096, in vitro and in vivo by using a rat model of visceral hypersensitivity. *J Pharmacol Sci.* 2008;106: 219–224.
64. Bingham B, Jones PG, Uveges AJ, et al. Species-specific in vitro pharmacological effects of the cannabinoid receptor 2 (CB2) selective ligand AM1241 and its resolved enantiomers. *Br J Pharmacol.* 2007;151:1061–1070.
65. Clapper JR, Moreno-Sanz G, Russo R, et al. Anandamide suppresses pain initiation through a peripheral endocannabinoid mechanism. *Nat Neurosci.* 2010;13:1265–1270.
66. Malik Z, Bayman L, Valetin J, et al. Dronabinol increases pain threshold in patients with functional chest pain: a pilot double-blind placebo-controlled trial. *Dis Esophagus.* 2017;30:1–8.
67. Sarfaraz S, Adhami VM, Syed DN, et al. Cannabinoids for cancer treatment: progress and promise. *Cancer Res.* 2008;68:339–342.
68. Bedoya F, Meneu JC, Macias MI, et al. Mutation in CNR1 gene and VEGF expression in esophageal cancer. *Tumori.* 2009;95:68–75.
69. Hijiya N, Shibata T, Daa T, et al. Overexpression of cannabinoid receptor 1 in esophageal squamous cell carcinoma is correlated with metastasis to lymph nodes and distant organs, and poor prognosis. *Pathol Int.* 2017;67:83–90.
70. Azofeifa A, Mattson ME, Schauer G, et al. National estimates of marijuana use and related indicators—national survey on drug use and health, united states, 2002–2014. *MMWR Surveill Summ.* 2016;65:1–28.
71. Rock EM, Parker LA. Cannabinoids as potential treatment for chemotherapy-induced nausea and vomiting. *Front Pharmacol.* 2016;7:221.

Cite this article as: Gotfried J, Kataria R, Schey R (2017) Review: the role of cannabinoids on esophageal function: what we know thus far, *Cannabis and Cannabinoid Research* 2:1, 252–258, DOI: 10.1089/can.2017.0031.

Abbreviations Used

CBRs = cannabinoid receptors
CNS = central nervous system
ECS = endocannabinoid system
ENS = enteric nervous system
ERD = erosive esophageal reflux disease
GERD = gastroesophageal reflux disease
GI = gastrointestinal
GPR55 = G-protein receptor 55
NERD = nonerosive esophageal reflux disease
THC = tetrahydrocannabinol
TLESRs = transient lower esophageal sphincter relaxations
WD = Western diet

Publish in Cannabis and Cannabinoid Research

- Immediate, unrestricted online access
- Rigorous peer review
- Compliance with open access mandates
- Authors retain copyright
- Highly indexed
- Targeted email marketing

liebertpub.com/can