Unraveling the Role of Vegetables in Spreading Antimicrobial-Resistant Bacteria: A Need for Quantitative Risk Assessment

Christina Susanne Hölzel,¹ Julia Louisa Tetens,¹ and Karin Schwaiger²

Abstract

In recent years, vegetables gain consumer attraction due to their reputation of being healthy in combination with low energy density. However, since fresh produce is often eaten raw, it may also be a source for foodborne illness. The presence of antibiotic-resistant bacteria might pose a particular risk to the consumer. Therefore, this review aims to present the current state of knowledge concerning the exposure of humans to antibiotic-resistant bacteria via food of plant origin for quantitative risk assessment purposes. The review provides a critical overview of available information on hazard identification and characterization, exposure assessment, and risk prevention with special respect to potential sources of contamination and infection chains. Several comprehensive studies are accessible regarding major antimicrobial-resistant foodborne pathogens (e.g., *Salmonella* spp., *Listeria* spp., *Bacillus cereus*, *Campylobacter* spp., *Escherichia coli*) and other bacteria (e.g., further Enterobacteriaceae, *Pseudomonas* spp., Gram-positive cocci). These studies revealed vegetables to be a potential—although rare—vector for extended-spectrum beta-lactamase-producing Enterobacteriaceae, *mcr1*-positive *E. coli*, colistin- and carbapenem-resistant *Pseudomonas aeruginosa*, linezolid-resistant enterococci and staphylococci, and vancomycin-resistant enterococci. Even if this provides first clues for assessing the risk related to vegetable-borne antimicrobial-resistant bacteria, the literature research reveals important knowledge gaps affecting almost every part of risk assessment and management. Especially, the need for (comparable) quantitative data as well as data on possible contamination sources other than irrigation water, organic fertilizer, and soil becomes obvious. Most crucially, dose–response studies would be needed to convert a theoretical “risk” (e.g., related to antimicrobial-resistant commensals and opportunistic pathogens) into a quantitative risk estimate.

Keywords: ESBL, antimicrobial resistance, antibiotic resistance, foodborne pathogens, vegetables, risk assessment

Introduction

VEGETABLES ARE ESSENTIAL for healthy nutrition and obesity reduction in Western civilizations, due to a high nutrient density, correlated with low energy density (Darmon *et al.*, 2005). They are defined as “a plant, root, seed, or pod that is used as food, especially in dishes that are not sweet” in the Cambridge dictionary. In 2003, WHO and FAO started an initiative to promote fruit and vegetable consumption for health worldwide, with a recommended minimum intake of 400 g fruits and vegetables per day (www.who.int).

However, vegetables are also implicated in foodborne outbreaks due to various—most often unknown—sources of contamination, the most recent familiar case in Europe being

the enterohemorrhagic *Escherichia coli* (EHEC) outbreak of 2011, with 4321 reported cases, of which at least 50 had been fatal (Robert-Koch-Institute, Germany, cited from Buchholz *et al.*, 2011). Besides other unusual features, the outbreak strain was characterized as an extended-spectrum beta-lactamase (ESBL) producer due to the presence of *bla*_{CTX-M15} genes (Mellmann *et al.*, 2011). The press reported extensively on antimicrobial resistance (AMR), but not on the fact that this resistance was meaningless for therapy, due to contraindication of antibiotics in EHEC infections (see Hazard Identification and Characterization section).

According to the FAO definition, quantitative risk assessment is based on four steps: hazard identification, hazard characterization, exposure assessment, and risk characterization

¹Animal Hygiene and Animal Health Management, Faculty of Agricultural and Nutritional Sciences, Christian-Albrechts-University Kiel (CAU), Kiel, Germany.

²Department of Veterinary Sciences, Institute of Food Safety, Ludwig-Maximilians-University Munich (LMU), Munich, Germany.

© Christina Susanne Hölzel et al. 2018; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are cited.

(www.fao.org). In this definition, hazard identification means to identify impacts on human health and the circumstances under which the danger is present, whereas hazard characterization aims to quantitatively or qualitatively evaluate the adverse effects on human health (e.g., using dose–response relationship). Exposure assessment provides the likely degree of consumption or intake of the hazardous agent, and the risk characterization tries to offer an estimate of the likely adverse effect in the target population by integrating the first three steps.

There have been well-recognized approaches for risk assessment related to general microbial hazards imposed by vegetable consumption (Hamilton *et al.*, 2006; Franz *et al.*, 2010; Danyluk and Schaffner, 2011; Pang *et al.*, 2017). However, no specific quantitative risk assessment has been published for hazards related to antimicrobial-resistant bacteria (ARB) in vegetables, although AMR is seen as an obligate “risk amplifier” in public awareness. Steps to assess specific risks due to AMR would involve all steps of quantitative risk assessment. This starts from hazard identification (i.e., is a known adverse health effect aggravated by AMR features of the causative agent? Does an adverse health effect arise from spread of antibiotic resistance features in commensals, which otherwise would not be considered as microbial hazards?) but also includes hazard characterization as well as exposure assessment.

This literature survey aims to initiate a discussion on quantitative risk assessment related to the spread of antibiotic resistance in the vegetable food chain. For this purpose, we tried to provide an overview over current screening data starting from 2007. To allow risk prevention, sources of contamination were highlighted as well. Finally, important knowledge gaps were identified, which should be addressed in the future to facilitate quantitative risk assessment within a reasonable term.

Literature Search and Exclusion Criteria

Our PubMed search combined the keywords “antimicrobial resistance” or “antibiotic resistance” with “vegetables” or “fresh produce” in May 2018. We aimed to provide a systematic overview on recent screening data for antibiotic-resistant bacteria isolated from vegetables, so we covered the period from January 2007 to April 2018 and found 169 studies, after excluding double hits. The search result comprised 93 studies that reported screening data for antibiotic resistance in vegetable-borne isolates (Tables 1–7). We excluded 22 obvious mishits (one study dealing with the inhibition of bladder cancer cell proliferation by mustard oil, for example) and 13 studies that did not investigate vegetable-borne isolates (instead, these studies reported on, e.g., reptiles). Eight studies were excluded since they did not attribute resistance rates from different sources, e.g., meat and vegetables, as one common value). We excluded opinion reports, reviews (unless they comprised relevant screening data), and publications that described modeling or method implementation or whole genome sequencing data of single isolates (in total 14 studies). We also excluded three studies in a language other than English (Polish, Ukrainian, and Chinese). One study that would not have met the exclusion criteria was excluded afterward since it presented mainly resistance data for *Escherichia coli* and vancomycin, a substance that is not

effective in Enterobacteriaceae. All other studies presenting intrinsic resistance data (e.g., erythromycin in *Salmonella* spp.) were kept since significant results for acquired resistance were presented as well.

All 13 experimental studies returned by the keywords were included in the text (e.g., experimental application of manure or biosolids to soil), as were two studies that reported meta-genomic screening data.

Besides our systematic approach, we used targeted searches with individual keywords to add details (e.g., on vegetable consumption).

Hazard Identification and Characterization

A recent outbreak due to vegetables was related to a multiresistant strain of EHEC O104:H4, which produced ESBL enzymes (Buchholz *et al.*, 2011). Did this fact add adverse effects to the outbreak? In general, antibiotics are not the first-line agents for treatment of diarrhea (Guerrant *et al.*, 2001). On the contrary, for EHEC, as compiled by Goldwater and Bettelheim (2012), antibiotics—and especially beta-lactams—are even contraindicated due to the fact that cell-wall-impaired dead bacteria release more toxin (Tarr *et al.*, 2005; Smith *et al.*, 2012). Even subinhibitory concentrations are thought to increase toxin production and/or toxin release (Grif *et al.*, 1998). Thus, in the absence of treatment indication, AMR could not lead to treatment failure. However, for other bacterial infections such as systemic listeriosis, antibiotic therapy is the treatment of choice (Safdar and Armstrong, 2003), and clinical resistance against first-line antibiotics is likely to impose additional health hazards. Centers for Disease Control and Prevention report recent cases of listeriosis in the United States due to vegetable consumption, which led to several deaths (www.cdc.gov); it is unknown whether these deaths were related to treatment failure or other reasons such as delayed diagnosis.

Providing comprehensive hazard identification of all food-borne pathogens associated with vegetables goes beyond the scope of this review. Thus, three more general situations will be exemplarily illustrated and connected to AMR: (1) primary foodborne infectious disease due to obligatory or opportunistic pathogens, (2) foodborne microbial intoxication, (3) foodborne colonization, maybe followed by opportunistic disease after a considerable time-shift (Fig. 1a–c).

Primary foodborne infectious disease is seen after infection with pathogenic serovars of, for example, *Salmonella enterica* subsp. *enterica*, *Campylobacter* spp., *Listeria monocytogenes*, *Bacillus cereus*, and *E. coli* (e.g., EHEC, Shiga-toxin producing *E. coli* [STEC]) (Butler *et al.*, 2015). In case of systemic infection, as already mentioned, antimicrobial treatment is often indicated (except for EHEC), and AMR to clinically relevant antimicrobial drugs will lower the number of therapeutic options (Fig. 1a). In contrast, foodborne *Staphylococcus aureus* exclusively causes food poisoning due to the presence of preformed heat-stable toxins independent of viable bacteria (Kadariya *et al.*, 2014). Thus, this kind of food poisoning is self-limiting, and antibiotic treatment is never indicated (Fig. 1b). However, *S. aureus* might cause opportunistic infections of wounds like cuts (van de Sande-Bruinsma *et al.*, 2015), which might occur during food preparation and might call for topical treatment. Only a limited number of *B. cereus* lineages (Ehling-Schulz *et al.*,

TABLE 1. STUDIES ON MAJOR ANTIMICROBIAL-RESISTANT FOODBORNE PATHOGENS (*SALMONELLA ENTERICA* SUBSP. *ENTERICA*, *LISTERIA* SPP., *BACILLUS CEREUS*, *CAMPYLOBACTER* SPP.) IN VEGETABLES

<i>Species/serovar</i>	<i>No. of isolates from vegetables</i>	<i>Specification of vegetables</i>	<i>Tested antimicrobials (method)</i>	<i>Region</i>	<i>Source</i>
<i>Salmonella</i> Cubana, 50 others	14, 138	Cantaloupes, celery, cilantro, green onions, hot peppers, lettuce, parsley, spinach, alfalfa sprouts, and tomatoes	NARMS gram negative panel amoxicillin, ampicillin, cefoxitin, <u>cephalothin</u> , chloramphenicol, <u>kanamycin</u> , nalidixic acid, <u>streptomycin</u> , sulfamethoxazole, sulfisoxazole, <u>tetracycline</u> , trimethoprim (<i>microdilution, sensitivity</i>)	United States	Reddy <i>et al.</i> (2016)
<i>Salmonella</i> Weltevreden, <i>Salmonella</i> Agona, others	33, 24, 77	Indigenous vegetables	Ampicillin , cefotaxime, ceftriaxone, chloramphenicol, erythromycin, gentamicin, nalidixic acid, streptomycin, tetracycline, trimethoprim-sulfamethoxazole (<i>disk diffusion</i>)	Malaysia	Yoke-Kqueen <i>et al.</i> (2008)
<i>S. enterica</i> subsp. <i>enterica</i> —serovar not specified	26	Tomato, spinach, carrot, radish, cantaloupe, and cucumber	Amikacin, amoxicillin-clavulanic acid, ampicillin , aztreonam, azithromycin, cefixime, cefoxitin, chloramphenicol , ciprofloxacin, colistin , cotrimoxazole, doxycycline, erythromycin, gatifloxacin, gentamicin, imipenem, kanamycin, meropenem, moxifloxacin, nalidixic acid, netilmicin, norfloxacin, ofloxacin, piperacillin, polymyxin B, sparfloxacin, streptomycin, tetracycline, tobramycin (<i>disk diffusion</i>)	India	Verma <i>et al.</i> (2018)
<i>B. cereus</i>	110	Fermented soybean products	Ampicillin , cefepime, cefotetan, ciprofloxacin, chloramphenicol, clindamycin, erythromycin, gentamicin, imipenem, oxacillin , penicillin, rifampin, sulfamethoxazole, tetracycline, trimethoprim, vancomycin (<i>disk diffusion</i>)	Korea	Kim <i>et al.</i> (2015)
<i>B. cereus</i>	39	Sunsik (a Korean ready-to-eat food prepared from grains, fruits, and vegetables)	Ampicillin , cefepime, cefoxitin, chloramphenicol, gentamicin, imipenem, oxacillin , penicillin, sulfamethoxazole, tetracycline, vancomycin (<i>disk diffusion</i>)	Korea	Lee <i>et al.</i> (2012)
<i>B. cereus</i>	87	Fermented soybean products	Ampicillin , cefepime, cefotetan, ciprofloxacin, chloramphenicol, clindamycin, erythromycin, gentamicin, imipenem, oxacillin , penicillin, rifampin, sulfamethoxazole/trimethoprim, tetracycline, vancomycin (<i>disk diffusion</i>)	Korea	Yim <i>et al.</i> (2015)

(continued)

TABLE 1. (CONTINUED)

Species/serovar	No. of isolates from vegetables	Specification of vegetables	Tested antimicrobials (method)	Region	Source
<i>Listeria monocytogenes</i> , <i>Listeria grayi</i> , <i>Listeria welshimeri</i> , <i>Listeria ivanovii</i> <i>L. monocytogenes</i>	30, 42, 46, 28 144	Ready-to-eat salads Fresh and frozen vegetables	Amikacin, ampicillin, cefotaxime, cephalothin, chloramphenicol, clindamycin, erythromycin, gentamicin, neomycin, oxacillin, penicillin, streptomycin, tetracycline, vancomycin^a (disk diffusion) Ampicillin, amoxicillin, chloramphenicol, ciprofloxacin, erythromycin, gentamicin, minocycline, norfloxacin, rifampicin, sulfamethoxazole, tetracycline, trimethoprim, vancomycin (<i>microdilution, E-test</i>)	Turkey Poland	Gurler <i>et al.</i> (2015) Korsak <i>et al.</i> (2012)
<i>Campylobacter jejuni</i>	33	Salad style vegetables (ulam)	Ampicillin, amikacin , ciprofloxacin, enrofloxacin, erythromycin , gentamicin, norfloxacin, penicillin G, tetracycline, vancomycin (disk diffusion) n.d.	Malaysia n.d.	Khalid <i>et al.</i> (2015) Singh <i>et al.</i> (2007); Miranda <i>et al.</i> (2009); Guchi and Ashenafi (2010); Yan <i>et al.</i> (2010); Raufu <i>et al.</i> (2013); Taban <i>et al.</i> (2013); Yu <i>et al.</i> (2014); Chen <i>et al.</i> (2015); Gurler <i>et al.</i> (2015); Mäka <i>et al.</i> (2015); Lee <i>et al.</i> (2016); Niyomdechach <i>et al.</i> (2016); de Vasconcelos Byrne <i>et al.</i> (2016); Liu and Kilongo-Nthenge (2017)
	Each study <25	n.d.	n.d.		

Font normal: resistance tested but not found; underlined: resistance in one single isolate; bold face: resistance in two or more isolates. Resistance defined as in the original study.

^aIn *L. ivanovii* and *L. welshimeri*.

TABLE 2. STUDIES ON ANTIMICROBIAL-RESISTANT *ESCHERICHIA COLI* IN VEGETABLES

No. of isolates from vegetables	Specification of vegetables	Tested antimicrobials (method)	Region	Source
70	Leafy green vegetables	Amoxicillin–clavulanic acid, ampicillin, cefoxitin, ceftazidime, ceftriaxone, co-trimoxazole, chloramphenicol, ciprofloxacin, gentamicin, nalidixic acid, streptomycin, tetracycline (disk diffusion)	South Africa	Jongman and Korsten (2016)
208	Leek, radish, basil, parsley, spinach, and lettuce, commercial and traditional	Ampicillin, cephalothin, chloramphenicol, ciprofloxacin, enrofloxacin, gentamicin, imipenem, lincomycin, nitrofurantoin, streptomycin, sulfamethoxazole, tetracycline, trimethoprim (disk diffusion)	Iran	Shakerian <i>et al.</i> (2016)
Approximately 230 (STEC)	whole salad Leafy vegetables and other vegetables from private produce farms	<i>mcr1</i> , <i>mcr2</i> (PCR)	California	Mavrici <i>et al.</i> (2017)
60	Salad vegetables, cucumber, parsley, tomato, mint, and lettuce, salad (restaurant)	Ampicillin, cephalothin, colistin, co-trimoxazole, gentamicin, streptomycin, sulphatriad, tetracycline (disk diffusion)	Lebanon	Faour-Klingbeil <i>et al.</i> (2016)
72	Lettuce	Amikacin, amoxicillin–clavulanic acid, ampicillin, cefotaxime, cephalothin, chloramphenicol, ciprofloxacin, gentamicin, kanamycin, nalidixic acid, streptomycin, sulfonamide, tetracycline, trimethoprim (<i>disk diffusion</i>)	Belgium	Holvoet <i>et al.</i> (2013)
82 (Diarrheogenic pathotypes)	Cactus salad (nopalitos)	Amikacin, amoxicillin–clavulanic acid, ampicillin, ceftriaxone, chloramphenicol, ciprofloxacin, colistin, co-trimoxazole, erythromycin, gentamicin, kanamycin, nalidixic acid, neomycin, streptomycin, tetracycline (<i>disk diffusion</i>)	Mexico	Gómez-Aldapa <i>et al.</i> (2016)
78 (from 13 samples)	Packaged ready-to-eat salad	Amoxicillin–clavulanic acid, ciprofloxacin, chloramphenicol, co-trimoxazole, gentamicin, kanamycin, nalidixic acid, streptomycin, tetracycline (disk diffusion, E-test)	78 (from 13 samples)	Campos <i>et al.</i> (2013)
73	Tomato, spinach, carrot, radish, cantaloupe, and cucumber	Amikacin, amoxicillin–clavulanic acid, ampicillin, aztreonam, azithromycin, cefixime, cefoxitin, chloramphenicol, ciprofloxacin, colistin, co-trimoxazole, doxycycline, erythromycin, gatifloxacin, gentamicin, imipenem, kanamycin, meropenem, moxifloxacin, nalidixic acid, netilmicin, norfloxacin, ofloxacin, piperacillin, polymyxin B, sparfloxacin, streptomycin, tetracycline, tobramycin (disk diffusion)	India	Verma <i>et al.</i> (2018)
239	Collard, cucumber, lettuce, tomato, and spinach	Amoxicillin–clavulanic acid, ampicillin, piperacillin, piperacillin–tazobactam, cefepime, ceftazidime, cefotaxime, imipenem, aztreonam, gentamicin, streptomycin, nalidixic acid, ciprofloxacin, tetracycline, chloramphenicol, co-trimoxazole (disk diffusion)	Portugal	Araújo <i>et al.</i> (2017)
29	Spinach, cabbage	Amikacin, amoxicillin–clavulanic acid, ampicillin, aztreonam, cefepime, cefotaxime, cefoxitin, ceftazidime, ceftriaxone, chloramphenicol, ciprofloxacin, enrofloxacin, florfenicol, gentamicin, kanamycin, nalidixic acid, neomycin, streptomycin, tetracycline, trimethoprim–sulfamethoxazole	South Africa	Du Plessis <i>et al.</i> (2017)
Each study <25	n.d.	n.d.	n.d.	Hassan <i>et al.</i> (2011); Skočková <i>et al.</i> (2013); Kim and Woo (2014); Rasheed <i>et al.</i> (2014); Jones-Dias <i>et al.</i> (2016); Zurhuf <i>et al.</i> (2016); Lima <i>et al.</i> (2017); Randall <i>et al.</i> (2017); Wang <i>et al.</i> (2017)

Font normal: resistance tested but not found; underlined: resistance in one single isolate; bold face: resistance in two or more isolates. Resistance defined as in the original study. STEC, Shiga-toxin producing *E. coli*.

TABLE 3. STUDIES ON ANTIMICROBIAL-RESISTANT PSEUDOMONADS IN VEGETABLES

Species	No. of isolates from vegetables	Specification of vegetables	Tested antimicrobials (method)	Region	Source
<i>Pseudomonas aeruginosa</i> , <i>Pseudomonas putida</i>	295, 106	Fruit, root, bulbous vegetables, salads, and cereals	<u>Amikacin</u> , apramycin , cefepime , ceftazidime , ciprofloxacin , colistin , doxycycline (only in <i>P. putida</i>), enrofloxacin , gentamicin , imipenem , neomycin , netilmicin , piperacillin+tazobactam , streptomycin , tobramycin , (<i>microdilution</i>)	Germany	Schwaiger <i>et al.</i> (2011a)
<i>P. aeruginosa</i>	88	Lettuce, white cabbage, red cabbage, carrots, sweet pepper, cucumber, and tomatoes, mixed	Ampicillin , aztreonam , ceftazidime , chloramphenicol, ciprofloxacin, <u>gentamicin</u> , imipenem, sulfamethoxazole-trimethoprim , tetracycline (<i>disk diffusion</i>)	Jamaica	Allydice-Francis and Brown (2012)
<i>Pseudomonas</i> spp.	35	Leaf lettuces, tomatoes, and carrots	<u>Amikacin</u> , cefepime , cefotaxime , ceftazidime , ciprofloxacin, imipenem , <u>gentamicin</u> (<i>disk diffusion</i>)	Portugal	Jones-Dias <i>et al.</i> (2016)
	Each study <25	n.d.	n.d.	n.d.	Bezanson <i>et al.</i> (2008); Hassan <i>et al.</i> (2011); Estepa <i>et al.</i> (2015)

Font normal: resistance tested but not found; underlined: resistance in one single isolate; bold face: resistance in two or more isolates. Resistance defined as in the original study.

2005) are able to cause mere food poisoning due to the presence of cereulide (see Rosenquist *et al.*, 2005, for example). Instead, in most cases, *B. cereus* needs to infect the organism to cause gastrointestinal disease (due to synthesis of Nhe and other enterotoxins in the intestine; EFSA, 2016). This is similar to the situation seen in *Clostridium perfringens* (Uzal *et al.*, 2014). Antibiotic treatment of *B. cereus* infections is recommended in severe disease, whereas it is unreasonable in self-limited food poisoning (Spiliopoulou *et al.*, 2014). *C. perfringens* might contribute to antibiotic-associated diarrhea (Borriello *et al.*, 1984; Kim *et al.*, 2017b); treatment of choice is discontinued primary antibiotics; only severe cases might be treated with glycopeptides or metronidazole (Bergogne-Bérézin, 2000).

A particular situation is found for organisms that are able to colonize the human gastrointestinal tract (Fig. 1c), such as *E. coli* or enterococci. These bacteria are useful indicators for AMR (Franklin *et al.*, 2001; Schwaiger *et al.*, 2011b) and were included in a recent EFSA recommendation to provide harmonized European monitoring of AMR (EFSA, 2012), as authorized by Directive 2003/99/EC. In addition, *E. coli* might produce ESBL enzymes, and this ESBL production (or other resistance phenotypes) might complicate the treatment of opportunistic infections. However, it is impossible to relate such colonization to a particular foodborne source, such as it is done for foodborne outbreaks: the onset of opportunistic

infectious disease (e.g., urinary tract infection due to ESBL *E. coli*) might sporadically happen years after ingestion of the contaminated food. Thus, according to Depoorter *et al.* (2012), the human health risk posed by a given exposure to foodborne ESBL *E. coli* cannot be estimated yet.

In general, the situation described in Figure 1a–c applies for all foods, not only vegetables. However, the risk to encounter infection or colonization is elevated in vegetables, due to a high share of raw consumption.

Opportunistic pathogens (and apathogenic commensals) might further serve as a vehicle for the transfer of antimicrobial resistance genes (ARGs) to pathogens. This might happen in the intestine, but also before consumption as observed on lettuce with a resistance gene of clinical relevance in humans (*bla*_{SHV18}, Jung and Matthews, 2016). DNA is partly degraded by heat, so raw consumed food is also more likely to pass high concentrations of ARGs into the human intestine.

In terms of hazard identification, additional hazard due to AMR is identified in systemic foodborne infection and in opportunistic colonization, but not in self-limiting diarrhea and intoxication.

In terms of hazard characterization, dose–response relationships are the most critical point for quantitative risk assessment. In case of obligate pathogens that would be antibiotically treated (Fig. 1a), the dose–response relationship

TABLE 4. STUDIES ON OTHER ANTIMICROBIAL-RESISTANT GRAM NEGATIVES IN VEGETABLES

Genus/species	No. of isolates from vegetables	Specification of vegetables	Tested antimicrobials (method)		Region	Source
<i>Acinetobacter</i> spp.	89	Leaf lettuces, tomatoes, and carrots	<u>Amikacin</u> , <u>cefepime</u> , <u>cefotaxime</u> , <u>ceftazidime</u> , <u>ciprofloxacin</u> , <u>gentamicin</u> , <u>imipenem</u> , <u>tetracycline</u> (<i>disk diffusion</i>)		Portugal	Jones-Dias <i>et al.</i> (2016)
<i>Cronobacter</i> spp.	34	Carrot, cucumber, leafy greens, sprouts, and others	Ampicillin, amoxicillin+clavulanic acid, apramycin, aztreonam, cefotaxime, chloramphenicol, ciprofloxacin, colistin, enrofloxacin, gentamicin, kanamycin, meropenem, nalidixic acid, neomycin, streptomycin, sulfonamides , co-trimoxazole, tetracycline, trimethoprim (<i>disk diffusion</i>)		Czech Republic	Vojkowska <i>et al.</i> (2016)
<i>Enterobacter cloacae</i> , <i>Enterobacter geroviae</i> , <i>Pantoea agglomerans</i>	172, 92, 96	Fruit, root, bulbous vegetables, salads, and cereals	Amikacin, amoxicillin+clavulanate (not tested in <i>E. cloacae</i>), ampicillin (<i>E. geroviae</i> only), apramycin, cefador (not tested in <i>E. cloacae</i>), cefepime, ceftazidime , ciprofloxacin, colistin , ^a <u>doxycycline</u> , enrofloxacin, imipenem , mezlocillin , neomycin, netilmicin, piperacillin , piperacillin+tazobactam, gentamicin, streptomycin , co-trimoxazole , tobramycin (<i>microdilution</i>)		Germany	Schwaiger <i>et al.</i> (2011a)
<i>Helicobacter pylori</i>	52	Leek, radish, basil, parsley, spinach, lettuce, cabbage, carrot, scallion, chive, fenugreek, coriander, pepper, turnip, beet, garlic, maize, broccoli, and cucumber, commercial and traditional salad samples	Amoxicillin , ampicillin , cefsulodin , clarithromycin , erythromycin , furazolidone , levofloxacin , metronidazole , rifampin , spiramycin , streptomycin , tetracycline , trimethoprim (<i>disk diffusion</i>)		Iran	Yahaghi <i>et al.</i> (2014)
Enterobacteriaceae, <i>Helicobacter pylori</i> , Coliforms, <i>Acinetobacter baumannii</i> , <i>Sphingobacterium multivorum</i> , <i>Cronobacter</i> spp.	Each study <25 or studies, which mix/sum up resistance data of different genera	n.d.	n.d.		n.d.	Bezanson <i>et al.</i> (2008); Hassan <i>et al.</i> (2011); Mokhtari <i>et al.</i> (2012); Campos <i>et al.</i> (2013); Hemmatinezhad <i>et al.</i> (2016); Jones-Dias <i>et al.</i> (2016); Mesbah Zekar <i>et al.</i> (2017); Fei <i>et al.</i> (2018)

Font normal: resistance tested but not found; underlined: resistance in one single isolate; bold face: resistance in two or more isolates. Resistance defined as in the original study.

^aR > 8 mg/L.

TABLE 5. STUDIES ON ANTIMICROBIAL-RESISTANT GRAM-POSITIVE COCCI IN VEGETABLES

Species	No. of isolates from vegetables	Specification of vegetables	Tested antimicrobials (method)	Region	Source
<i>Staphylococcus aureus</i>	53	Lettuce, perilla leaf, and sprouts	Amoxicillin-clavulanic acid , cephalothin , chloramphenicol, ciprofloxacin , clindamycin, erythromycin , gentamicin , imipenem , linezolid, nitrofurantoin, oxacillin , penicillin , quinupristin/dalfopristin , telithromycin, tetracycline , trimethoprim/sulfamethoxazole, vancomycin (<i>disk diffusion</i>)	Korea	Hong <i>et al.</i> (2015)
<i>Staphylococcus saprophyticus</i> , <i>Staphylococcus succinus</i> , <i>Staphylococcus xylosus</i>	49, 14, 18	Soybean products: meju and doenjang	Ampicillin , chloramphenicol, erythromycin , gentamicin, lincomycin , penicillin G , tetracycline , trimethoprim (<i>disk diffusion</i>)	Korea	Jeong <i>et al.</i> (2016)
<i>Enterococcus faecalis</i>	100	Fruit, root, bulbous vegetables, salads, and cereals	Amoxicillin-clavulanic acid, ampicillin, mezlocillin, imipenem , gentamicin high level, streptomycin high level , chloramphenicol , florfenicol, ciprofloxacin , enrofloxacin, moxifloxacin, vancomycin, teicoplanin, erythromycin , tylosin, <u>linezolid</u> , doxycycline , nitrofurantoin, rifampicin	Germany	Schwaiger <i>et al.</i> (2011a)
<i>Enterococcus faecium</i> , <i>E. faecalis</i>	26, 9	Herbs, spinach, and leafy greens	Ampicillin, bacitracin, chloramphenicol, ciprofloxacin , erythromycin , gentamicin, kanamycin , linezolid, nitrofurantoin , penicillin, quinupristin-dalfopristin , ^a streptomycin high level, teicoplanin, tetracycline , vancomycin (<i>disk diffusion</i>)	Canada	Allen <i>et al.</i> (2013)
<i>Enterococcus casseliflavus</i> , <i>E. faecalis</i> , <i>E. faecium</i> , <i>Enterococcus hirae</i> , <i>Enterococcus gallinarum</i> , <i>Enterococcus</i> spp.	40, 20, 18, 9, 5, 16	Ready-to-eat salads (split or mixed leaves, carrot, and corn)	Amoxicillin, chloramphenicol, ciprofloxacin, erythromycin , gentamicin, minocycline , nitrofurantoin, quinupristin-dalfopristin , ^b streptomycin, teicoplanin, tetracycline , vancomycin	Portugal	Campos <i>et al.</i> (2013)
<i>E. faecium</i> , <i>E. hirae</i> , <i>E. faecalis</i> , <i>E. casseliflavus</i>	34, 23, 4, 4	Vegetables, soil, and irrigation water	Ampicillin, chloramphenicol , ciprofloxacin , erythromycin , gentamicin high level , kanamycin high level , streptomycin high level , teicoplanin, tetracycline , trimethoprim-sulfamethoxazole, vancomycin ^c (<i>disk diffusion</i>)	Tunisia	Ben Said <i>et al.</i> (2016)
Each study <25	n.d.	n.d.	n.d.	n.d.	Abriouel <i>et al.</i> (2008); Gomes <i>et al.</i> (2008); Hassan <i>et al.</i> (2011); Burgos <i>et al.</i> (2014); Lavilla Lerma <i>et al.</i> (2014); Pesavento <i>et al.</i> (2014); Rodriguez-Palacios <i>et al.</i> (2014); Xu <i>et al.</i> (2014); Hao <i>et al.</i> (2015)

Font normal: resistance tested, but not found; underlined: resistance in one single isolate; bold face: resistance in two or more isolates. Resistance defined as in the original study.

^a*E. faecalis*; ^b*E. casseliflavus*, *E. faecalis*; ^c*E. casseliflavus*.

TABLE 6. COMPARATIVE PREVALENCE (%) OF ANTIMICROBIAL-RESISTANT *ESCHERICHIA COLI* IN MEAT AND VEGETABLES (NONSELECTIVE APPROACH; CALCULATED FROM PERCENT POSITIVE SAMPLES AND PERCENT RESISTANT ISOLATES)

Antibiotic	Retail chicken (n = 250)	Retail pork (n = 250)	Vegetables (farm+retail, n = 1001)
Amoxicillin–clavulanate	13.65 ^a	2.37 ^b	1.10 ^b
Ampicillin	34.58 ^a	5.90 ^b	1.70 ^c
Piperacillin	23.33 ^a	4.34 ^b	0.40 ^c
Cefaclor	9.30 ^a	2.76 ^b	1.90 ^b
Cefoxitin	2.40	0.78	1.30
Cefuroxime	1.95	0.39	0.80
Imipenem	1.65 ^a	0.00 ^b	0.00 ^b
Gentamicin	1.13 ^a	0.39 ^{a,b}	0.10 ^b
Streptomycin	23.63 ^a	7.12 ^b	0.30 ^c
Tobramycin	0.00	0.39	0.00
Chloramphenicol	3.60 ^a	1.17 ^{a,b}	0.50 ^b
Florfenicol	0.00	0.39	0.60
Ciprofloxacin	3.98 ^a	0.39 ^b	0.00 ^b
Enrofloxacin	4.50 ^a	0.39 ^b	0.00 ^b
Colistin	0.00	0.39	0.30
Doxycycline	30.53 ^a	9.46 ^b	0.50 ^c
Sulfamethoxazole—trimethoprim	31.35 ^a	5.51 ^b	0.20 ^c

Different superscripts in a row: values differ significantly in a chi-squared test (Fisher's exact test, if expected values are below 5 per cell).

is that of foodborne disease, so data are available for many pathogens. In contrast, dose–response data are completely missing for stable intestinal colonization or opportunistic extraintestinal disease. For colonization, hints are provided from studies using oronasal infection to colonize animals: Schoeni and Doyle (1994) isolated *E. coli* O157:H7 from cecal tissue of single chickens 3 months after experimental inoculation with as low as 2.6×10^1 colony-forming unit (cfu).

Exposure Assessment

Produce and human consumption of vegetables

Community members of the EU produced vegetables (including melons and strawberries) on a total area of 224,126,000 ha in 2016 (<http://ec.europa.eu>). In Germany, more than 8000 farms produced 3,672,660 tons of vegetables for human consumption (www.destatis.de); roots, salads, and other vegetables that are commonly consumed raw constituted approximately one-third of this produce. According to EFSA data (<https://dwh.efsa.europa.eu>) based on 60 studies (1997–2015) from 25 countries, the mean daily consumption of vegetables and vegetable products in adults ranged from 62 g per capita in Sweden to 382 g in Romania, with excessive consumption observed in individuals (99th percentile = 1047 g); these figures do not include plant products such as grains or legumes.

Prevalence of ARB in vegetables

A PubMed search combining the keywords “antimicrobial resistance” or “antibiotic resistance” with “vegetables” or “fresh produce” covered the period between January 2007 and April 2018. After applying the exclusion criteria as described above, 93 hits were condensed in Tables 1–7.

The studies included bacteria of more than 20 different genera (Tables 1–7), mainly *E. coli*, *S. enterica* subsp. *enterica*, *Listeria* spp., enterococci, pseudomonads, and *B. cereus*. To

provide basic information from the studies, resistances are marked as present or absent in Tables 1–5 (irrespective of whether they are acquired or intrinsic). Additionally, prevalences are included in Tables 6 and 7. In general, comparisons between studies should be avoided since different choices of breakpoints introduce significant bias when comparing resistance data according to the European EUCAST organization with the U.S. Clinical and Laboratory Standards Institute (CLSI). Moreover, many studies lack the information of whether they used appropriate reference strains in antimicrobial susceptibility testing.

For *Salmonella*, studies are hard to compare due to a high diversity in serotypes. A huge number of vegetables (more than 100,000 samples) were investigated in the United States from 2002 to 2012, resulting in isolation of 152 *Salmonella* strains, of which as few as 10 had detectable AMRs (Reddy *et al.*, 2016; Table 1). However, CLSI warned that *Salmonella* might be falsely reported as susceptible to several antibiotics (CLSI M100-S22E, table 2A in the CLSI document) due to differences between *in vitro* and *in vivo* susceptibility. Yoke-Kqueen *et al.* (2008) reported 56.7% of 134 isolates with a multiple antimicrobial resistance (MAR) index of more than 0.2. However, erythromycin was included in the MAR. Corrected for this fact, 44.3% of isolates from Malaysian vegetables were resistant to two or more antibiotics (mainly tetracycline+streptomycin—substances that are not the first- or second-line drugs in human therapy of salmonellosis). Relevant multiresistance (>5 resistances, including third-generation cephalosporins) was found in *Salmonella* Albany, *Salmonella* Brunei, and *Salmonella* Kralingen (Yoke-Kqueen *et al.*, 2008).

All identified studies dealing with AMR in vegetable-borne *B. cereus* (*n* isolates = 39–110, Table 1) referred to Korean fermented or traditional food (Lee *et al.*, 2012; Kim *et al.*, 2015; Yim *et al.*, 2015), thus statements are restrained to a limited variety of food and regions. In addition, resistance was assessed only by means of disk diffusion, and the source and

TABLE 7. PREVALENCE OF VEGETABLE-BORNE BACTERIA WITH PARTICULARLY RELEVANT ANTIMICROBIAL RESISTANCE (EXAMPLES)

Microorganism	Antimicrobial resistance	Prevalence of resistance: No. of resistant isolates/total No. of isolates (%)	Vegetable source	Region	References
<i>Salmonella enterica</i> subsp. <i>enterica</i> (serovar Albany, Brunei, Kralingen)	Third-generation cephalosporin	4/134 (3.0)	Indigenous vegetable food	Malaysia	Yoke-Kqueen <i>et al.</i> (2008)
<i>Bacillus cereus</i>	Vancomycin	1/87 (1.2)	Fermented soybeans	Korea	Yim <i>et al.</i> (2015)
		5/35 (14.3)	Sunsik	Korea	Lee <i>et al.</i> (2012)
<i>Escherichia coli</i>	ESBL genes	3/56 (5.4)	Lettuce	Portugal	Araújo <i>et al.</i> (2017)
	ESBL genes and phenotypes	One from 32 salad samples	Salad	Spain	Egea <i>et al.</i> (2011)
		Zero from 400 fruit or vegetable samples	Fruit and vegetables	United Kingdom	Randall <i>et al.</i> (2017)
	Carbapenem	12/180 (6.6)	Leek, radish, basil, spinach, lettuce, traditional salads, commercial salads	Iran	Shakerian <i>et al.</i> (2016)
<i>Enterobacteriaceae</i>	<i>mcr1</i>	1/?	Lettuce	Portugal	Jones-Dias <i>et al.</i> (2016)
		0/230	Vegetables	United States	Mavrici <i>et al.</i> (2017)
	ESBL phenotype	2/60 (3.3) from 42 samples	Imported vegetables	Switzerland	Zurfluh <i>et al.</i> (2016)
		44/97	Various vegetables and fruits	Algeria	Mesbah Zekar <i>et al.</i> (2017)
<i>Enterobacteriaceae</i> (<i>E. coli</i> , <i>Klebsiella pneumoniae</i> , <i>Enterobacter cloacae</i> , <i>Enterobacter aerogenes</i> , <i>Cronobacter sakazakii</i>)	ESBL phenotype+ESBL genes	25.4% of vegetable samples	Imported vegetables	Dominican Republic, India, Thailand, Vietnam (imported to Switzerland)	Zurfluh <i>et al.</i> (2015)
<i>Enterobacteriaceae</i> (<i>K. pneumoniae</i> , <i>Serratia marcescens</i>)	ESBL phenotype+ESBL genes	3/138 (2.2) from 9 samples	Iceberg lettuce	United States	Bhutani <i>et al.</i> (2015)
<i>Raoultella terrigena</i>	ESBL genes	2/2	Salads	Portugal	Campos <i>et al.</i> (2013)
<i>Pseudomonas</i> spp.	Colistin ^a , imipenem, meropenem ^a	8/401 (2.0), 26/401 (6.5), 1/401 (0.2) from 1001 samples	Fruit, root, bulbous vegetables, salads, and cereals	Germany	Schwaiger <i>et al.</i> (2011a)
<i>Staphylococcus aureus</i> (MSSA)	Linezolid	1/53	Leafy vegetables	Korea	Hong <i>et al.</i> (2015)
<i>Enterococcus faecalis</i> + <i>faecium</i>	Vancomycin	3/160 (1.9) from 540 samples	Vegetables+environment (not specified)	Korea	Kim <i>et al.</i> (2017a)
<i>E. faecalis</i> + <i>faecium</i>	Linezolid	1/100 (1.0), 1/59 (1.7) from 1001 samples	Fruit, root, bulbous vegetables, salads and cereals	Germany	Schwaiger <i>et al.</i> (2011a)

^aR > 8; CLSI breakpoint: R ≥ 8, for colistin recently adjusted to R ≥ 4.
?, unknown number; CLSI, Clinical and Laboratory Standards Institute; ESBL, extended-spectrum beta-lactamase; MSSA, methicillin-sensitive *Staphylococcus aureus*.

FIG. 1. Impact of antimicrobial resistance in different situations following vegetable-borne (a) infection, (b) intoxication, or (c) colonization (schematic illustration). *not for EHEC infection

validity of breakpoints is unclear, since these studies did not refer to CLSI or other acknowledged standards, but to the secondary literature. Apart from substances to which *B. cereus* should be considered as intrinsically resistant (e.g., ampicillin, oxacillin, penicillin, cefepime, rifampin), resistance rates were moderate, with complete susceptibility to gentamicin, imipenem, ciprofloxacin, tetracycline, and vancomycin in one study (Kim *et al.*, 2015). However, resistance to vancomycin—a first-line therapeutic in (extraintestinal) *B. cereus* infections—was reported in 1/87 isolates from fermented soybeans (Yim *et al.*, 2015) and 5/39 isolates from Sunsik (Lee *et al.*, 2012), a ready-to-eat food made from grains, fruits, and vegetables.

Significant data for *L. monocytogenes* ($n=144$, Table 1) from fresh and frozen vegetables were reported from a Polish study using E-test with CLSI-breakpoints for listeria and staphylococci (Korsak *et al.*, 2012): only one vegetable-borne isolate had detectable AMR at all (to tetracyclines). In Turkish ready-to-eat salad, listerial AMR was most pronounced to erythromycin and cephalothin (Gurler *et al.*, 2015). However, cephalosporin resistance is considered as an intrinsic feature of *L. monocytogenes* (Collins *et al.*, 2012), and erythromycin is not considered as a treatment of choice (Morvan *et al.*, 2010).

Campylobacter jejuni ($n=33$, Table 1) from Malaysian “salad style vegetables” had moderate resistance rates except for intrinsic resistances and erythromycin resistance (60.6%; Khalid *et al.*, 2015). The latter fact might be remarkable since macrolides have for long been used as a therapy of choice (Blaser *et al.*, 1979; Engberg *et al.*, 2001).

However, the study is generally limited by the fact that breakpoints were outdated (taken from a document from 2003, when the CLSI was still named NCCLS). In any case, fluoroquinolones—a first-line therapy of today—were highly effective in this study.

Resistance data for *E. coli* (60–239 isolates, Table 2) differed significantly between studies; this might be related to the very different locations (South Africa, India, United States, South America, Europe, Lebanon, Iran), the different nature of samples, and the different choice of antibiotics and breakpoints.

The majority of studies reported pronounced resistance to streptomycin and tetracyclines—a finding of limited significance, given the fact that clinical breakpoints cut within natural populations. However, studies also reported significant resistance to amoxicillin combined with clavulanic acid (>15%) (e.g., Araújo *et al.*, 2017) or resistance to carbapenems (Shakerian *et al.*, 2016) and third- or fourth-generation cephalosporin as well as presence of *bla*_{TEM1}-genes (Araújo *et al.*, 2017). Remarkably, Gómez-Aldapa *et al.* (2016) reported 100% resistance to amikacin and colistin in diarrheagenic *E. coli* pathotypes from Mexican cactus salads. The study referred to CLSI; however, CLSI does not provide a breakpoint for colistin in *E. coli*. In addition, colistin resistance was assessed by disk diffusion, a practice that is discouraged nowadays (www.eucast.org).

For pathogenic *E. coli*, one study assessed the prevalence of plasmid-borne colistin resistance genes *mcr1/mcr2* in STEC *E.*

coli from the United States; not a single positive isolate was found in 1000 strains from different sources, including about 230 isolates from vegetables (Mavrici *et al.*, 2017).

Due to their environmental sources, pseudomonads are frequently found in vegetables. Pseudomonads are not considered as major foodborne pathogens; however, six studies were identified, which reported resistance data of up to 401 *Pseudomonas* isolates from, mostly *Pseudomonas aeruginosa* (Table 3). Using microdilution and DIN58940-breakpoints where available, one study identified significantly higher resistance rates to aminoglycosides in *Pseudomonas* isolates from fruit vegetables compared with root vegetables or salads (Schwaiger *et al.*, 2011a). In this study, resistance also included antibiotics used in clinics, such as gentamicin, tobramycin, amikacin, ciprofloxacin, colistin, piperacillin, ceftazidime, imipenem, or meropenem; however, such resistances were rare (Schwaiger *et al.*, 2011a). Using CLSI breakpoints and disk diffusion, most of these resistances were also observed in Jamaican vegetable isolates, apart from resistance to carbapenems (Allydice-Francis and Brown, 2012).

Cronobacter spp. acts as a foodborne pathogen in immunocompromised patients (Healy *et al.*, 2010) and will then be antibiotically treated, for example, with ampicillin plus gentamicin or chloramphenicol (Lai, 2001). However, vegetable isolates were characterized by a high rate of pansusceptible strains and lacked resistance against clinically relevant antimicrobials (Vojkowska *et al.*, 2016; Table 4). For *Enterobacter* spp. ($n=264$), several AMRs were significantly lowered at retail, compared with farm level, pointing toward a need for sampling close to consumption for particular questions of risk assessment (Schwaiger *et al.*, 2011a; Table 4). Once more, resistance depended on the vegetable group: resistance to colistin was significantly more frequent in *Enterobacter cloacae* from fruit vegetables than from roots (Schwaiger *et al.*, 2011a).

None of the studies on Gram-positive cocci (Table 5) reported staphylococci to be resistant to vancomycin, a critically important antibiotic especially for infection caused by methicillin-resistant *S. aureus* (MRSA). Korean leafy vegetables were partly contaminated with multiresistant MRSA strains (Hong *et al.*, 2015). One methicillin-susceptible isolate was resistant to linezolid, a first-line drug for the treatment of MRSA-infected wounds (Gurusamy *et al.*, 2013). Single linezolid-resistant strains of *Enterococcus faecalis* (1/100) and *Enterococcus faecium* were isolated from German vegetables (Schwaiger *et al.*, 2011a). One linezolid-resistant *E. faecium* was detected in Canadian vegetables as well (Allen *et al.*, 2013). Detectable resistance in *E. faecalis* isolates ($n=20$) from Portuguese ready-to-eat salad was restrained to tetracyclines and erythromycin due to the presence of *tet*(M), *tet*(L), and *erm*(B) (Campos *et al.*, 2013). From Tunisia, Ben Said *et al.* (2016) reported additional resistance to high level concentrations of aminoglycosides and chloramphenicol. By multilocus sequence typing, Leavis *et al.* (2006) identified vegetable-borne ARB of a so-called high-risk enterococcal clonal complex. Importantly, vancomycin-teicoplanin-resistant enterococci were isolated from fresh produce or its environment in Korea (Kim *et al.*, 2017a).

Human exposure to vegetable-borne ARB

Assessing the exposure of humans to vegetable-borne ARB is not the same as assessing the prevalence of AMR in

vegetable-borne bacteria: Considerable exposure to ARB might arise in a situation when bacteria are rarely found, but frequently resistant, or vice versa in the situation when AMR is moderate, but the prevalence of bacterial contamination is very high. For *Salmonella* or *Listeria* spp., both prevalence of bacteria and prevalence of AMR within vegetable-borne bacteria are low (Reddy *et al.*, 2016). For other bacteria such as pseudomonads, prevalence of bacteria is high, but the prevalence of clinically significant resistance is low.

Vice versa, the prevalence of *E. coli* or *E. faecalis* is much lower in vegetables than in food of animal origin, but the prevalence of AMR is remarkably high for some antibiotics and, although rare, resistances to critically important isolates are present as well. For *E. coli*, we had the chance to comparatively calculate the prevalence of ARB (prevalence of bacteria $\times$ prevalence of resistance) in pork, poultry, and vegetables from two studies (Schwaiger *et al.*, 2011a, 2012 and unpublished data) using identical methods; pork and poultry were included for the sake of comparison. The studies resembled each other in identical (nonselective) bacterial isolation, identical susceptibility testing (microdilution, DIN 58940-81), and identical source of breakpoints. Results are shown in Table 6. In total, the prevalence of *E. coli* was lowest in vegetables (3.4% compared with 75% in chicken and 25% in pork). However, for several antibiotics, the prevalence of resistant *E. coli* was quite comparable between pork and vegetables. This was because a much lower prevalence of *E. coli* in vegetables was accompanied by a high prevalence of resistance: up to 55% of the vegetable-borne *E. coli* were resistant to beta-lactams.

For quantitative risk assessment, it would be crucial to know not just the prevalence but also the concentration of ARB in vegetables since risk characterization has to consider dose-related data. However, this information is more than rare. Some of the studies listed in Tables 1–7 quantified bacteria (e.g., Hassan *et al.*, 2011; Campos *et al.*, 2013; Holvoet *et al.*, 2013; Araújo *et al.*, 2017; Du Plessis *et al.*, 2017) but did not specifically refer to ARB. To assess these data, selective quantification for resistant bacteria would be needed, which is performed even rarer. Relevant data are provided by Ruimy *et al.* (2010), who reported densities of resistant bacteria as high as 10^4 cfu/g of product for Gram negatives grown in the presence of tetracycline, chloramphenicol, and nalidixic acid. Yang *et al.* (2016) quantified antibiotic-resistant endophytic bacteria in experimentally grown pak choi and found concentrations of up to 9×10^2 per g.

Risk Characterization

Risk characterization integrates the informations from hazard identification, hazard characterization, and exposure assessment into a quantitative estimate of risk (probability of adverse outcome or number of cases). Hazard identification related to resistant bacteria has to account for the fact whether this resistance is of therapeutic interest. This review revealed that clinically relevant antibiotics are underrepresented in current studies so that many pathogen-AMR combinations listed above do not imply any therapeutic consequence (hazard). However, ESBL production in Enterobacteriaceae or linezolid resistance in enterococci and staphylococci is surely of clinical interest.

Risk characterization would also include hazard characterization, which means dose–response relationships. Infectious

doses vary in the individual host and situation; for many environmental bacteria, such infectious doses have not been defined at all. In addition, the amount of opportunistic pathogens or commensals, which is necessary to produce stable colonization in the human intestine, is unknown for most bacteria, and dose–response studies are absent. For foodborne outbreaks in general, case numbers are also used as a substitute in hazard characterization if dose–response data are missing. However, no specific case numbers are available for infectious disease due to vegetable-borne ARB.

In the absence of known colonization doses or known frequencies of colonization events, at least qualitative data might help to further characterize the risk. Thus, attempts were made to identify genetically related antimicrobially resistant organisms in food and human intestines or human clinical samples in a distinct region and period (e.g., Abriouel *et al.*, 2008; Hannah *et al.*, 2009; Burgos *et al.*, 2014), but without showing significant association. Several studies tried to relate the presence of ARB in human feces to consumption behavior and found that vegetarians are at higher risk to carry ARB (Elder *et al.*, 1993; Sannes *et al.*, 2008). However, Sannes *et al.* (2008) also showed that this might be linked to a confounder related to the lifestyle of vegetarians: traveling abroad.

Exposure assessment identified vegetables as a relevant vehicle in terms of a high consumption frequency and in terms of frequent bacterial contamination in a ready-to-eat state. However, the exposure to clinically relevant pathogens or to commensals conferring clinically relevant resistance is low, compared with other food like meat.

Ultimately, taking together the individual steps of risk assessment, we have to state that the absence of dose–response data and the absence of published estimates on treatment failure due to AMR in foodborne infections do not allow quantitative risk characterization for vegetable-borne ARB yet.

Risk Prevention: Sources of Contamination

While sprout seeds from Egypt were considered as the most probable outbreak source for the multinational EHEC outbreak in 2011, the primary source of sprout contamination remained unknown. This is seen in most outbreaks (not only) related to vegetables, thus indirect clues are important for assessing infection chains. In rare cases, molecular typing of bacterial isolates from possible sources and contaminated vegetables is available to strengthen these indirect clues.

Irrigation water as a source of ARB

Worldwide, the FAO indicates a need for irrigation water as high as 1500 km³/year; of this irrigation water, 7.38% are used for vegetables and roots (www.fao.org, data from 2012). Groundwater constitutes 38.9% of this irrigation water, the rest is taken from sources such as surface water or used water, which are more easily prone to microbial contamination than groundwater. Microbial concentrations in irrigation water are generally much lower than they are in organic fertilizers; however, filter effects of soil might concentrate microorganisms at soil surfaces, and irrigation is more continuously applied to vegetables than is organic fertilizer. Blaak *et al.* (2015) found that Dutch surface water contains 2.2×10^2 multidrug-resistant *E. coli* (median) and up to 6×10^2 ESBL *E. coli* per microliter.

Several studies found that irrigation water and vegetables shared the same clones, as confirmed by repetitive-element polymerase chain reaction (rep-PCR: BOX-, REP-, and ERIC-PCR) (Jongman and Korsten, 2016; Araújo *et al.*, 2017). Njage and Buys (2015) identified irrigation water as a possible pathway of transmission for ESBL *E. coli*.

Soil and organic fertilizer as sources of ARB

Vegetables produced in or close to soil—such as carrots and leaf vegetables—are at special risk for contamination with soil-borne bacteria, either belonging to natural soil microbiota or introduced into soil by manure fertilization. While the application of manure in ready-to-eat crops is discouraged, for example, by the U.K. Food Standards Agency (www.food.gov.uk); it is not generally prohibited by European law. In Germany, however, the use of liquid manure for top dressing is interdicted by national law; any vegetables must not be planted earlier than 12 weeks after fields had been fertilized with liquid manure.

Several studies focused on the prevalence of ARB or AMR genes in vegetables at harvest and the soil from which these vegetables were harvested (Marti *et al.*, 2013, 2014; Wang *et al.*, 2015; He *et al.*, 2016; Lau *et al.*, 2017; Tien *et al.*, 2017); these studies found temporary—if any—impact of manuring compared with natural soil, which has a natural base level of ARG due to the presence of antimicrobial-producing microorganisms (Nesme and Simonet, 2015). Ruimy *et al.* (2010) showed that resistance scores were significantly higher in vegetables grown in or close to soil compared with fruits and vegetables grown above the soil. Rahube *et al.* (2014, 2016) reported that human biosolids are a relevant—but also temporary—source of ARG on vegetables at harvest, and Duan *et al.* (2017) found that ARG contents in soil and lettuce decreased after experimental application of biochar. Zhu *et al.* (2017) reported eightfold higher absolute copy numbers or ARG in manure-fertilized Chinese organic lettuce compared with conventionally produced lettuce; this difference was mainly due to different microbial communities in both types of samples. One recent study searched for ARGs in the viral DNA fraction (bacteriophages) from vegetables and soil and found that concentrations were highest for lettuce and soil, pointing toward soil as a relevant source of ARGs (Larrañaga *et al.*, 2018).

Antibiotic residues in manure or soil might be more critical than is presence of manure-borne ARB, due to higher persistence. Antibiotic residues were able to select antimicrobially resistant endophytic bacteria, which are not removed by washing or peeling (Zhang *et al.*, 2017; Esteban-Cuesta *et al.*, 2018).

Direct contamination by humans

To our knowledge, systematic studies on human contamination of vegetables with ARB during production or processing are missing, as is systematic source tracking by molecular microbiological methods.

In the 2011 EHEC outbreak, it was assumed that, in a regional suboutbreak, vegetable dishes were contaminated by an infected caterer (www.rki.de). Genotyping raised suspicion that (multiresistant) *S. aureus* from Korean leafy vegetables were of human origin (Hong *et al.*, 2015); the presence of new class 3-integrins in a pathogenic strain of *Klebsiella pneumoniae* was related to clinical sources by Jones-Dias

et al. (2016). As an additional hint, we found that fruit vegetables were generally more likely to be contaminated with ARB than other vegetables; touching by consumers is discussed as a possible explanation, although not proven yet (Schwaiger *et al.*, 2011a). A similar hypothesis was presented by Mesbah Zekar *et al.* (2017). Weak food hygiene standards on Nigerian farm were shown to correlate with higher bacterial loads in cut lettuce (Oyinlola *et al.*, 2017), but these standards included only some parameters of personal hygiene besides a couple of nonhuman hygiene factors, such as surface water irrigation and manure application.

Besides the mentioned routes of contamination, other plausible ways of contamination include a broad variety of sources, either for the primary products (e.g., *Salmonella* contamination by reptiles and amphibians, Gorski *et al.*, 2013) or during transport, processing, and distribution. Macrorestriction plus pulsed-field gel electrophoresis suggested that two different kinds of vegetables from two different countries were contaminated with a clonal *E. coli* strain due to packaging and distribution at the same factory. The role of processing is further illustrated by cut salads, which are generally considered as a risk food (www.fda.gov): here, processing is not only a possible source of cross-contamination but also a factor increasing water activity (thus decreasing microbial stability).

Outlook: Transfer of ARB to Humans by Food of Plant Origin—Current Lacks and Future Needs for Risk Assessment

A considerable share of studies used nonvalidated methods, reported intrinsic resistance, dealt with nonpathogenic bacteria, or provided data on AMRs/ARGs of low clinical relevance. Such limitations should be proactively addressed, if present. Commensal bacteria might be useful indicator organisms and serve as donors for AMR genes in the gut; in addition, the label pathogenic/nonpathogenic is conferred more carefully than in former since pathogenesis is more and more treated as a non-static interaction between strain-specific virulence and susceptibility of individual hosts. However, this fact gives even more emphasis to a need for prudent study design, overcoming the actual strategy of “count and collect” in favor of more functional approaches leading to dose–response data.

Important lacks of knowledge affect almost every part of risk assessment and management, starting from identifying sources of contamination and ending up to the black box of intestinal and infectious processes long after consumption. With regard to sources of contamination, search for plant-derived ARB should include processing and distribution steps since contamination close to consumption is more likely to result in human exposure, and human contamination might be more likely to introduce strains with AMR against human therapeutics. Further research should thus focus on clinically relevant antibiotics, should provide (or reject) evidence of transmission at clonal level, and should clarify and quantify the involvement of commensals in ARG transfer to pathogens as well as in extraintestinal opportunistic infections after transmission by food. Future studies addressing these gaps would be more than welcome.

Disclosure Statement

No competing financial interests exist.

References

- Abriouel H, Omar NB, Molinos AC, López RL, Grande MJ, Martínez-Viedma P, Ortega E, Cañamero MM, Galvez A. Comparative analysis of genetic diversity and incidence of virulence factors and antibiotic resistance among enterococcal populations from raw fruit and vegetable foods, water and soil, and clinical samples. *Int J Food Microbiol* 2008; 123:38–49.
- Allen KJ, Kovacevic J, Cancarevic A, Wood J, Xu J, Gill B, Allen JK, Mesak LR. Microbiological survey of imported produce available at retail across Canada. *Int J Food Microbiol* 2013;162:135–142.
- Allydice-Francis K, Brown PD. Diversity of antimicrobial resistance and virulence determinants in *Pseudomonas aeruginosa* associated with fresh vegetables. *Int J Microbiol* 2012; 2012:426241.
- Araújo S, A T Silva I, Tação M, Patinha C, Alves A, Henriques I. Characterization of antibiotic resistant and pathogenic *Escherichia coli* in irrigation water and vegetables in household farms. *Int J Food Microbiol* 2017;257:192–200.
- Ben Said L, Klibi N, Dziri R, Borgo F, Boudabous A, Ben Slama K, Torres C. Prevalence, antimicrobial resistance and genetic lineages of *Enterococcus* spp. from vegetable food, soil and irrigation water in farm environments in Tunisia. *J Sci Food Agric* 2016;96:1627–1633.
- Bergogne-Bérézin E. Treatment and prevention of antibiotic associated diarrhea. *Int J Antimicrob Agents* 2000;16:521–526.
- Bezanson GS, MacInnis R, Potter G, Hughes T. Presence and potential for horizontal transfer of antibiotic resistance in oxidase-positive bacteria populating raw salad vegetables. *Int J Food Microbiol* 2008;127:37–42.
- Bhutani N, Muraleedharan C, Talreja D, Rana SW, Walia S, Kumar A, Walia SK. Occurrence of multidrug resistant extended spectrum beta-lactamase-producing bacteria on iceberg lettuce retailed for human consumption. *Biomed Res Int* 2015;2015:547547.
- Blaak H, Lynch G, Italiaander R, Hamidjaja RA, Schets FM, de Roda Husman AM. Multidrug-resistant and extended spectrum beta-lactamase-producing *Escherichia coli* in Dutch surface water and wastewater. *PLoS One* 2015;10: e0127752.
- Blaser MJ, Berkowitz ID, LaForce FM, Cravens J, Reller LB, Wang WL. *Campylobacter* enteritis: Clinical and epidemiologic features. *Ann Intern Med* 1979;91:179.
- Borriello SP, Larson HE, Welch AR, Barclay F, Stringer MF, Bartholomew BA. Enterotoxigenic *Clostridium perfringens*: A possible cause of antibiotic-associated diarrhoea. *Lancet* 1984;1:305–307.
- Buchholz U, Bernard H, Werber D, Böhmer MM, Remschmidt C, Wilking H, Deleré Y, an der Heiden M, Adlhoch C, Dreesman J, Ehlers J, Ethelberg S, Faber M, Frank C, Fricke G, Greiner M, Höhle M, Ivarsson S, Jark U, Kirchner M, Koch J, Krause G, Luber P, Rosner B, Stark K, Kühne M. German outbreak of *Escherichia coli* O104:H4 associated with sprouts. *N Engl J Med* 2011;365:1763–1770.
- Burgos MJG, Aguayo MCL, Pulido RP, Gálvez A, López RL. Multilocus sequence typing and antimicrobial resistance in *Enterococcus faecium* isolates from fresh produce. *Antonie Van Leeuwenhoek* 2014;105:413–421.
- Butler AJ, Thomas MK, Pintar KDM. Expert elicitation as a means to attribute 28 enteric pathogens to foodborne, waterborne, animal contact, and person-to-person transmission routes in Canada. *Foodborne Pathog Dis* 2015;12:335–344.

- Campos J, Mourão J, Pestana N, Peixe L, Novais C, Antunes P. Microbiological quality of ready-to-eat salads: An underestimated vehicle of bacteria and clinically relevant antibiotic resistance genes. *Int J Food Microbiol* 2013;166:464–470.
- [CLSI] Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second Informational Supplement. CLSI document M100-S22. Wayne, PA: Clinical and Laboratory Standards Institute, 2012.
- Chen M, Wu Q, Zhang J, Wu S, Guo W. Prevalence, enumeration, and pheno- and genotypic characteristics of *Listeria monocytogenes* isolated from raw foods in South China. *Front Microbiol* 2015;6:1026.
- Collins B, Guinane CM, Cotter PD, Hill C, Ross RP. Assessing the contributions of the LiaS histidine kinase to the innate resistance of *Listeria monocytogenes* to nisin, cephalosporins, and disinfectants. *Appl Environ Microbiol* 2012;78:2923–2929.
- Danyluk MD, Schaffner DW. Quantitative assessment of the microbial risk of leafy greens from farm to consumption: Preliminary framework, data, and risk estimates. *J Food Prot* 2011;74:700–708.
- Darmon N, Darmon M, Maillot M, Drewnowski A. A nutrient density standard for vegetables and fruits: Nutrients per calorie and nutrients per unit cost. *J Am Diet Assoc* 2005;105:1881–1887.
- Depoorter P, Persoons D, Uyttendaele M, Butaye P, de Zutter L, Dierick K, Herman L, Imberechts H, van Huffel X, Dewulf J. Assessment of human exposure to 3rd generation cephalosporin resistant *E. coli* (CREC) through consumption of broiler meat in Belgium. *Int J Food Microbiol* 2012;159:30–38.
- Du Plessis EM, Govender S, Pillay B, Korsten L. Exploratory study into the microbiological quality of spinach and cabbage purchased from street vendors and retailers in Johannesburg, South Africa. *J Food Prot* 2017;80:1726–1733.
- Duan M, Li H, Gu J, Tuo X, Sun W, Qian X, Wang X. Effects of biochar on reducing the abundance of oxytetracycline, antibiotic resistance genes, and human pathogenic bacteria in soil and lettuce. *Environ Pollut* 2017;224:787–795.
- EFSA. Technical specifications on the harmonised monitoring and reporting of antimicrobial resistance in *Salmonella*, *Campylobacter* and indicator *Escherichia coli* and *Enterococcus* spp. bacteria transmitted through food. *EFSA J* 2012;10:311.
- EFSA. Risks for public health related to the presence of *Bacillus cereus* and other *Bacillus* spp. including *Bacillus thuringiensis* in foodstuffs. *EFSA J* 2016;14:122.
- Egea P, López-Cerero L, Navarro MD, Rodríguez-Baño J, Pascual A. Assessment of the presence of extended-spectrum beta-lactamase-producing *Escherichia coli* in eggshells and ready-to-eat products. *Eur J Clin Microbiol Infect Dis* 2011;30:1045–1047.
- Ehling-Schulz M, Svensson B, Guinebreteire M-H, Lindbäck T, Andersson M, Schulz A, Fricker M, Christiansson A, Granum PE, Märtlbauer E, Nguyen-The C, Salkinoja-Salonen M, Scherer S. Emetic toxin formation of *Bacillus cereus* is restricted to a single evolutionary lineage of closely related strains. *Microbiology* 2005;151(Pt 1):183–197.
- Elder HA, Roy I, Lehman S, Phillips RL, Kass EH. Human studies to measure the effect of antibiotic residues. *Vet Hum Toxicol* 1993;35(Suppl 1):31–36.
- Engberg J, Aarestrup FM, Taylor DE, Gerner-Smidt P, Nachamkin I. Quinolone and macrolide resistance in *Campylobacter jejuni* and *C. coli*: Resistance mechanisms and trends in human isolates. *Emerg Infect Dis* 2001;7:24–34.
- Esteban-Cuesta I, Drees N, Ulrich S, Stauch P, Sperner B, Schwaiger K, Gareis M, Gottschalk C. Endogenous microbial contamination of melons (*Cucumis melo*) from international trade: An underestimated risk for the consumer? *J Sci Food Agric* 2018;98:5074–5081.
- Estepa V, Rojo-Bezares B, Torres C, Sáenz Y. Genetic lineages and antimicrobial resistance in *Pseudomonas* spp. isolates recovered from food samples. *Foodborne Pathog Dis* 2015;12:486–491.
- Faour-Klingbeil D, Kuri V, Fadlallah S, Matar GM. Prevalence of antimicrobial-resistant *Escherichia coli* from raw vegetables in Lebanon. *J Infect Dev Ctries* 2016;10:354–362.
- Fei P, Jiang Y, Gong S, Li R, Jiang Y, Yuan X, Wang Z, Kang H, Ali MA. Occurrence, genotyping, and antibiotic susceptibility of *Cronobacter* spp. in drinking water and food samples from Northeast China. *J Food Prot* 2018;456–460.
- Franklin A, Acar J, Anthony F, Gupta R, Nicholls T, Tamura Y, Thompson S, Threlfall EJ, Vose D, van Vuuren M, White DG, Wegener HC, Costarrica ML. Antimicrobial resistance: Harmonisation of national antimicrobial resistance monitoring and surveillance programmes in animals and in animal-derived food. *Rev Sci Tech* 2001;20:859–870.
- Franz E, Tromp SO, Rijgersberg H, Van Der Fels-Klerx HJ. Quantitative microbial risk assessment for *Escherichia coli* O157:H7, *Salmonella*, and *Listeria monocytogenes* in leafy green vegetables consumed at salad bars. *J Food Prot* 2010;73:274–285.
- Goldwater PN, Bettelheim KA. Treatment of enterohemorrhagic *Escherichia coli* (EHEC) infection and hemolytic uremic syndrome (HUS). *BMC Med* 2012;10:12.
- Gomes BC, Esteves CT, Palazzo ICV, Darini ALC, Felis GE, Sechi LA, Franco BDGM, de Martinis ECP. Prevalence and characterization of *Enterococcus* spp. isolated from Brazilian foods. *Food Microbiol* 2008;25:668–675.
- Gómez-Aldapa CA, Cerna-Cortes JF, Rangel-Vargas E, Torres-Vitela MR, Villarruel-López A, Gutiérrez-Alcántara EJ, Castro-Rosas J. Presence of multidrug-resistant Shiga toxin-producing *Escherichia coli*, enteropathogenic *E. coli* and enterotoxigenic *E. coli*, on raw nopalitas (*Opuntia ficus-indica* L.) and in nopalitas salads from local retail markets in Mexico. *Foodborne Pathog Dis* 2016;13:269–274.
- Gorski L, Jay-Russell MT, Liang AS, Walker S, Bengson Y, Govoni J, Mandrell RE. Diversity of pulsed-field gel electrophoresis pulsotypes, serovars, and antibiotic resistance among *Salmonella* isolates from wild amphibians and reptiles in the California Central Coast. *Foodborne Pathog Dis* 2013;10:540–548.
- Grif K, Dierich MP, Karch H, Allerberger F. Strain-specific differences in the amount of Shiga toxin released from enterohemorrhagic *Escherichia coli* O157 following exposure to subinhibitory concentrations of antimicrobial agents. *Eur J Clin Microbiol Infect Dis* 1998;17:761–766.
- Guchi B, Ashenafi M. Microbial load, prevalence and antibiograms of *Salmonella* and *Shigella* in lettuce and green peppers. *Ethiop J Health Sci* 2010;20:41–48.
- Guerrant RL, van Gilder T, Steiner TS, Thielman NM, Slutsker L, Tauxe RV, Hennessy T, Griffin PM, DuPont H, Sack RB, Tarr P, Neill M, Nachamkin I, Reller LB, Osterholm MT, Bennis ML, Pickering LK. Practice guidelines for the management of infectious diarrhea. *Clin Infect Dis* 2001;32:331–351.
- Gurler Z, Pamuk S, Yildirim Y, Ertas N. The microbiological quality of ready-to-eat salads in Turkey: A focus on *Salmonella* spp. and *Listeria monocytogenes*. *Int J Food Microbiol* 2015;196:79–83.

- Gurusamy KS, Koti R, Toon CD, Wilson P, Davidson BR. Antibiotic therapy for the treatment of methicillin-resistant *Staphylococcus aureus* (MRSA) infections in surgical wounds. *Cochrane Database Syst Rev* 2013;(8):CD009726.
- Hamilton AJ, Stagnitti F, Premier R, Boland A-M, Hale G. Quantitative microbial risk assessment models for consumption of raw vegetables irrigated with reclaimed water. *Appl Environ Microbiol* 2006;72:3284–3290.
- Hannah EL, Johnson JR, Angulo F, Haddadin B, Williamson J, Samore MH. Molecular analysis of antimicrobial-susceptible and -resistant *Escherichia coli* from retail meats and human stool and clinical specimens in a rural community setting. *Foodborne Pathog Dis* 2009;6:285–295.
- Hao D, Xing X, Li G, Wang X, Zhang M, Zhang W, Xia X, Meng J. Prevalence, toxin gene profiles, and antimicrobial resistance of *Staphylococcus aureus* isolated from quick-frozen dumplings. *J Food Prot* 2015;78:218–223.
- Hassan SA, Altalhi AD, Gherbawy YA, El-Deeb BA. Bacterial load of fresh vegetables and their resistance to the currently used antibiotics in Saudi Arabia. *Foodborne Pathog Dis* 2011;8:1011–1018.
- He LY, Ying GG, Liu YS, Su HC, Chen J, Liu SS, Zhao JL. Discharge of swine wastes risks water quality and food safety: Antibiotics and antibiotic resistance genes from swine sources to the receiving environments. *Environ Int* 2016;92–93:210–219.
- Healy B, Cooney S, O'Brien S, Iversen C, Whyte P, Nally J, Callanan JJ, Fanning S. *Cronobacter* (*Enterobacter sakazakii*): An opportunistic foodborne pathogen. *Foodborne Pathog Dis* 2010;7:339–350.
- Hemmatinezhad B, Momtaz H, Rahimi E. *VacA*, *cagA*, *iceA* and *oipA* genotypes status and antimicrobial resistance properties of *Helicobacter pylori* isolated from various types of ready to eat foods. *Ann Clin Microbiol Antimicrob* 2016;15:2.
- Holvoet K, Sampers I, Callens B, Dewulf J, Uyttendaele M. Moderate prevalence of antimicrobial resistance in *Escherichia coli* isolates from lettuce, irrigation water, and soil. *Appl Environ Microbiol* 2013;79:6677–6683.
- Hong J, Kim Y, Kim J, Heu S, Kim S-R, Kim K-P, Roh E. Genetic diversity and antibiotic resistance patterns of *Staphylococcus aureus* isolated from leaf vegetables in Korea. *J Food Sci* 2015;80:M1526–M1531.
- Jeong D-W, Lee B, Her J-Y, Lee K-G, Lee J-H. Safety and technological characterization of coagulase-negative staphylococci isolates from traditional Korean fermented soybean foods for starter development. *Int J Food Microbiol* 2016;236:9–16.
- Jones-Dias D, Manageiro V, Ferreira E, Barreiro P, Vieira L, Moura IB, Caniça M. Architecture of class 1, 2, and 3 integrons from gram negative bacteria recovered among fruits and vegetables. *Front Microbiol* 2016;7:1400.
- Jongman M, Korsten L. Genetic diversity and antibiotic resistance of *Escherichia coli* isolates from different leafy green production systems. *J Food Prot* 2016;79:1846–1853.
- Jung Y, Matthews KR. Potential transfer of extended spectrum β -lactamase encoding gene, *blashv18* gene, between *Klebsiella pneumoniae* in raw foods. *Food Microbiol* 2016;60:39–48.
- Kadariya J, Smith TC, Thapaliya D. *Staphylococcus aureus* and staphylococcal food-borne disease: An ongoing challenge in public health. *Biomed Res Int* 2014;2014:827965.
- Khalid MI, Tang JYH, Baharuddin NH, Rahman NS, Rahimi NF, Radu S. Prevalence, antibiogram, and *cdt* genes of toxigenic *Campylobacter jejuni* in salad style vegetables (ulam) at farms and retail outlets in Terengganu. *J Food Prot* 2015;78:65–71.
- Kim C-W, Cho S-H, Kang S-H, Park Y-B, Yoon M-H, Lee J-B, No W-S, Kim J-B. Prevalence, genetic diversity, and antibiotic resistance of *Bacillus cereus* isolated from Korean fermented soybean products. *J Food Sci* 2015;80:M123–M128.
- Kim M-C, Cha M-H, Ryu J-G, Woo G-J. Characterization of vancomycin-resistant *Enterococcus faecalis* and *Enterococcus faecium* isolated from fresh produce and human fecal samples. *Foodborne Pathog Dis* 2017a;14:195–201.
- Kim S, Woo G-J. Prevalence and characterization of antimicrobial-resistant *Escherichia coli* isolated from conventional and organic vegetables. *Foodborne Pathog Dis* 2014;11:815–821.
- Kim YJ, Kim SH, Ahn J, Cho S, Kim D, Kim K, Lee H, Son H, Lee HJ, Yong D, Choi JY, Kim HR, Shin JH. Prevalence of *Clostridium perfringens* toxin in patients suspected of having antibiotic-associated diarrhea. *Anaerobe* 2017b;48:34–36.
- Korsak D, Borek A, Daniluk S, Grabowska A, Pappelbaum K. Antimicrobial susceptibilities of *Listeria monocytogenes* strains isolated from food and food processing environment in Poland. *Int J Food Microbiol* 2012;158:203–208.
- Lai KK. *Enterobacter sakazakii* infections among neonates, infants, children, and adults. Case reports and a review of the literature. *Medicine* 2001;80:113–122.
- Larrañaga O, Brown-Jaque M, Quirós P, Gómez-Gómez C, Blanch AR, Rodríguez-Rubio L, Muniesa M. Phage particles harboring antibiotic resistance genes in fresh-cut vegetables and agricultural soil. *Environ Int* 2018;115:133–141.
- Lau CH-F, Li B, Zhang T, Tien Y-C, Scott A, Murray R, Sabourin L, Lapen DR, Duenk P, Topp E. Impact of pre-application treatment on municipal sludge composition, soil dynamics of antibiotic resistance genes, and abundance of antibiotic-resistance genes on vegetables at harvest. *Sci Total Environ* 2017;587–588:214–222.
- Lavilla Lerma L, Benomar N, Valenzuela AS, Casado Muñoz Mdel C, Gálvez A, Abriouel H. Role of EfrAB efflux pump in biocide tolerance and antibiotic resistance of *Enterococcus faecalis* and *Enterococcus faecium* isolated from traditional fermented foods and the effect of EDTA as EfrAB inhibitor. *Food Microbiol* 2014;44:249–257.
- Leavis HL, Bonten MJM, Willems RJL. Identification of high-risk enterococcal clonal complexes: Global dispersion and antibiotic resistance. *Curr Opin Microbiol* 2006;9:454–460.
- Lee N, Sun JM, Kwon KY, Kim HJ, Koo M, Chun HS. Genetic diversity, antimicrobial resistance, and toxigenic profiles of *Bacillus cereus* strains isolated from Sunsik. *J Food Prot* 2012;75:225–230.
- Lee W-J, Kim H-B, Kim K-S. Isolation and characterization of spore-forming bacilli (SFB) from Shepherd's Purse (*Capsella bursa-pastoris*). *J Food Sci* 2016;81:M684–M691.
- Lima CM, Souza IEGL, Dos Santos Alves T, Leite CC, Evangelista-Barreto NS, de Castro Almeida RC. Antimicrobial resistance in diarrheagenic *Escherichia coli* from ready-to-eat foods. *J Food Sci Technol* 2017;54:3612–3619.
- Liu S, Kilonzo-Nthenge A. Prevalence of multidrug-resistant bacteria from U.S.-grown and imported fresh produce retailed in chain supermarkets and ethnic stores of Davidson County, Tennessee. *J Food Prot* 2017;80:506–514.
- Mąka Ł, Maćkiw E, Ścieżyńska H, Popowska M. Occurrence and antimicrobial resistance of *Salmonella* spp. isolated from food other than meat in Poland. *Ann Agric Environ Med* 2015;22:403–408.
- Marti R, Scott A, Tien Y-C, Murray R, Sabourin L, Zhang Y, Topp E. Impact of manure fertilization on the abundance of

- antibiotic-resistant bacteria and frequency of detection of antibiotic resistance genes in soil and on vegetables at harvest. *Appl Environ Microbiol* 2013;79:5701–5709.
- Marti R, Tien YC, Murray R, Scott A, Sabourin L, Topp E. Safely coupling livestock and crop production systems: How rapidly do antibiotic resistance genes dissipate in soil following a commercial application of swine or dairy manure? *Appl Environ Microbiol* 2014;80:3258–3265.
- Mavrici D, Yambao JC, Lee BG, Quiñones B, He X. Screening for the presence of *mcr-1/mcr-2* genes in Shiga toxin-producing *Escherichia coli* recovered from a major produce-production region in California. *PLoS One* 2017;12:e0187827.
- Mellmann A, Harmsen D, Cummings CA, Zentz EB, Leopold SR, Rico A, Prior K, Szczepanowski R, Ji Y, Zhang W, McLaughlin SF, Henkhaus JK, Leopold B, Bielaszewska M, Prager R, Brzoska PM, Moore RL, Guenther S, Rothberg JM, Karch H. Prospective genomic characterization of the German enterohemorrhagic *Escherichia coli* O104:H4 outbreak by rapid next generation sequencing technology. *PLoS One* 2011;6:e22751.
- Mesbah Zekar F, Granier SA, Marault M, Yaici L, Gassilloud B, Manceau C, Touati A, Millemann Y. From farms to markets: Gram-negative bacteria resistant to third-generation cephalosporins in fruits and vegetables in a region of North Africa. *Front Microbiol* 2017;8:1569.
- Miranda JM, Mondragón AC, Martínez B, Guarddon M, Rodríguez JA. Prevalence and antimicrobial resistance patterns of *Salmonella* from different raw foods in Mexico. *J Food Prot* 2009;72:966–971.
- Mokhtari W, Nsaibia S, Majouri D, Ben Hassen A, Gharbi A, Aouni M. Detection and characterization of *Shigella* species isolated from food and human stool samples in Nabeul, Tunisia, by molecular methods and culture techniques. *J Appl Microbiol* 2012;113:209–222.
- Morvan A, Moubareck C, Leclercq A, Hervé-Bazin M, Bremond S, Lecuit M, Courvalin P, Le Monnier A. Antimicrobial resistance of *Listeria monocytogenes* strains isolated from humans in France. *Antimicrob Agents Chemother* 2010;54:2728–2731.
- Nesme J, Simonet P. The soil resistome: A critical review on antibiotic resistance origins, ecology and dissemination potential in telluric bacteria. *Environ Microbiol* 2015;17:913–930.
- Niyomdechana N, Mungkorakaew N, Samosornsuk W. Serotypes and antimicrobial resistance of *Salmonella enterica* isolated from pork, chicken meat and lettuce, Bangkok and Central Thailand. *Southeast Asian J Trop Med Public Health* 2016;47:31–39.
- Njage PMK, Buys EM. Pathogenic and commensal *Escherichia coli* from irrigation water show potential in transmission of extended spectrum and AmpC β -lactamases determinants to isolates from lettuce. *Microb Biotechnol* 2015;8:462–473.
- Oyinlola LA, Obadina AO, Omemu AM, Oyewole OB. Prevention of microbial hazard on fresh-cut lettuce through adoption of food safety and hygienic practices by lettuce farmers. *Food Sci Nutr* 2017;5:67–75.
- Pang H, Lambertini E, Buchanan RL, Schaffner DW, Pradhan AK. Quantitative microbial risk assessment for *Escherichia coli* O157:H7 in fresh-cut lettuce. *J Food Prot* 2017;80:302–311.
- Pesavento G, Calónico C, Ducci B, Magnanini A, Lo Nostro A. Prevalence and antibiotic resistance of *Enterococcus* spp. isolated from retail cheese, ready-to-eat salads, ham, and raw meat. *Food Microbiol* 2014;41:1–7.
- Rahube TO, Marti R, Scott A, Tien YC, Murray R, Sabourin L, Duenk P, Lapen DR, Topp E. Persistence of antibiotic resistance and plasmid-associated genes in soil following application of sewage sludge and abundance on vegetables at harvest. *Can J Microbiol* 2016;62:600–607.
- Rahube TO, Marti R, Scott A, Tien YC, Murray R, Sabourin L, Zhang Y, Duenk P, Lapen DR, Topp E. Impact of fertilizing with raw or anaerobically digested sewage sludge on the abundance of antibiotic-resistant coliforms, antibiotic resistance genes, and pathogenic bacteria in soil and on vegetables at harvest. *Appl Environ Microbiol* 2014;80:6898–6907.
- Randall LP, Lodge MP, Elviss NC, Lemma FL, Hopkins KL, Teale CJ, Woodford N. Evaluation of meat, fruit and vegetables from retail stores in five United Kingdom regions as sources of extended-spectrum beta-lactamase (ESBL)-producing and carbapenem-resistant *Escherichia coli*. *Int J Food Microbiol* 2017;241:283–290.
- Rasheed MU, Thajuddin N, Ahamed P, Teklemariam Z, Jamil K. Antimicrobial drug resistance in strains of *Escherichia coli* isolated from food sources. *Rev Inst Med Trop Sao Paulo* 2014;56:341–346.
- Raufu I, Bortolaia V, Svendsen CA, Ameh JA, Ambali AG, Aarestrup FM, Hendriksen RS. The first attempt of an active integrated laboratory-based *Salmonella* surveillance programme in the north-eastern region of Nigeria. *J Appl Microbiol* 2013;115:1059–1067.
- Reddy SP, Wang H, Adams JK, Feng PCH. Prevalence and characteristics of *Salmonella* serotypes isolated from fresh produce marketed in the United States. *J Food Prot* 2016;79:6–16.
- Rodriguez-Palacios A, Ilic S, LeJeune JT. *Clostridium difficile* with moxifloxacin/clindamycin resistance in vegetables in Ohio, USA, and prevalence meta-analysis. *J Pathog* 2014;2014:158601.
- Rosenquist H, Smidt L, Andersen SR, Jensen GB, Wilcks A. Occurrence and significance of *Bacillus cereus* and *Bacillus thuringiensis* in ready-to-eat food. *FEMS Microbiol Lett* 2005;250:129–136.
- Ruimy R, Brisabois A, Bernede C, Skurnik D, Barnat S, Arlet G, Momcilovic S, Elbaz S, Moury F, Vibet M-A, Courvalin P, Guillemot D, Andremont A. Organic and conventional fruits and vegetables contain equivalent counts of gram-negative bacteria expressing resistance to antibacterial agents. *Environ Microbiol* 2010;12:608–615.
- Safdar A, Armstrong D. Antimicrobial activities against 84 *Listeria monocytogenes* isolates from patients with systemic listeriosis at a comprehensive cancer center (1955–1997). *J Clin Microbiol* 2003;41:483–485.
- Sannes MR, Belongia EA, Kieke B, Smith K, Kieke A, Vandermouse M, Bender J, Clabots C, Winokur P, Johnson JR. Predictors of antimicrobial-resistant *Escherichia coli* in the feces of vegetarians and newly hospitalized adults in Minnesota and Wisconsin. *J Infect Dis* 2008;197:430–434.
- Schoeni JL, Doyle MP. Variable colonization of chickens perorally inoculated with *Escherichia coli* O157:H7 and subsequent contamination of eggs. *Appl Environ Microbiol* 1994;60:2958–2962.
- Schwaiger K, Helmke K, Hölzel CS, Bauer J. Antibiotic resistance in bacteria isolated from vegetables with regards to the marketing stage (farm vs. supermarket). *Int J Food Microbiol* 2011a;148:191–196.
- Schwaiger K, Hölzel C, Bauer J. Detection of the macrolide-efflux protein A gene *mef(A)* in *Enterococcus faecalis*. *Microb Drug Resist* 2011b;17:429–432.
- Schwaiger K, Huther S, Hölzel C, Kämpf P, Bauer J. Prevalence of antibiotic-resistant Enterobacteriaceae isolated

- from chicken and pork meat purchased at the slaughterhouse and at retail in Bavaria, Germany. *Int J Food Microbiol* 2012; 154:206–211.
- Shakerian A, Rahimi E, Emad P. Vegetables and restaurant salads as a reservoir for Shiga toxinogenic *Escherichia coli*: Distribution of virulence factors, O-serogroups, and antibiotic resistance properties. *J Food Prot* 2016;79:1154–1160.
- Singh BR, Singh P, Agrawal S, Teotia U, Verma A, Sharma S, Chandra M, Babu N, Kant Agarwal R. Prevalence of multi-drug resistant *Salmonella* in coriander, mint, carrot, and radish in Bareilly and Kanpur, northern India. *Foodborne Pathog Dis* 2007;4:233–240.
- Skočková A, Karpíšková R, Kolářková I, Cupáková Š. Characteristics of *Escherichia coli* from raw vegetables at a retail market in the Czech Republic. *Int J Food Microbiol* 2013; 167:196–201.
- Smith KE, Wilker PR, Reiter PL, Hedican EB, Bender JB, Hedberg CW. Antibiotic treatment of *Escherichia coli* O157 infection and the risk of hemolytic uremic syndrome, Minnesota. *Pediatr Infect Dis J* 2012;31:37–41.
- Spiliopoulou A, Papachristou E, Foka A, Kolonitsiou F, Anastassiou ED, Goumenos DS, Spiliopoulou I. Relapsing *Bacillus cereus* peritonitis in a patient treated with continuous ambulatory peritoneal dialysis. *JMM Case Rep* 2014;1:e003400.
- Taban BM, Aytac SA, Akkoc N, Akcelik M. Characterization of antibiotic resistance in *Salmonella enterica* isolates determined from ready-to-eat (RTE) salad vegetables. *Braz J Microbiol* 2013;44:385–391.
- Tarr PI, Gordon CA, Chandler WL. Shiga-toxin-producing *Escherichia coli* and haemolytic uraemic syndrome. *Lancet* 2005;365:1073–1086.
- Tien Y-C, Li B, Zhang T, Scott A, Murray R, Sabourin L, Marti R, Topp E. Impact of dairy manure pre-application treatment on manure composition, soil dynamics of antibiotic resistance genes, and abundance of antibiotic-resistance genes on vegetables at harvest. *Sci Total Environ* 2017;581:582:32–39.
- Uzal FA, Freedman JC, Shrestha A, Theoret JR, Garcia J, Awad MM, Adams V, Moore RJ, Rood JJ, McClane BA. Towards an understanding of the role of *Clostridium perfringens* toxins in human and animal disease. *Future Microbiol* 2014;9:361–377.
- van de Sande-Bruinsma N, Leverstein van Hall MA, Janssen M, Nagtzaam N, Leenders S, Greeff SC de, Schneeberger PM. Impact of livestock-associated MRSA in a hospital setting. *Antimicrob Resist Infect Control* 2015;4:11.
- de Vasconcelos Byrne V, Hofer E, Vallim DC, de Castro Almeida RC. Occurrence and antimicrobial resistance patterns of *Listeria monocytogenes* isolated from vegetables. *Braz J Microbiol* 2016;47:438–443.
- Verma P, Saharan VV, Nimesh S, Singh AP. Phenotypic and virulence traits of *Escherichia coli* and *Salmonella* strains isolated from vegetables and fruits from India. *J Appl Microbiol* 2018;125:270–281.
- Vojtkovska H, Karpiskova R, Orieskova M, Drahovska H. Characterization of *Cronobacter* spp. isolated from food of plant origin and environmental samples collected from farms and from supermarkets in the Czech Republic. *Int J Food Microbiol* 2016;217:130–136.
- Wang F-H, Qiao M, Chen Z, Su J-Q, Zhu Y-G. Antibiotic resistance genes in manure-amended soil and vegetables at harvest. *J Hazard Mater* 2015;299:215–221.
- Wang L, Nakamura H, Kage-Nakadai E, Hara-Kudo Y, Nishikawa Y. Prevalence, antimicrobial resistance and multiple-locus variable-number tandem-repeat analysis profiles of diarrheagenic *Escherichia coli* isolated from different retail foods. *Int J Food Microbiol* 2017;249:44–52.
- Xu J, Shi C, Song M, Xu X, Yang P, Paoli G, Shi X. Phenotypic and genotypic antimicrobial resistance traits of foodborne *Staphylococcus aureus* isolates from Shanghai. *J Food Sci* 2014;79:M635–M642.
- Yahaghi E, Khamesipour F, Mashayekhi F, Safarpour Dehkordi F, Sakhaei MH, Masoudimanesh M, Khameneie MK. *Helicobacter pylori* in vegetables and salads: Genotyping and antimicrobial resistance properties. *Biomed Res Int* 2014; 2014:757941.
- Yan H, Neogi SB, Mo Z, Guan W, Shen Z, Zhang S, Li L, Yamasaki S, Shi L, Zhong N. Prevalence and characterization of antimicrobial resistance of foodborne *Listeria monocytogenes* isolates in Hebei province of Northern China, 2005–2007. *Int J Food Microbiol* 2010;144:310–316.
- Yang Q, Zhang H, Guo Y, Tian T. Influence of chicken manure fertilization on antibiotic-resistant bacteria in soil and the endophytic bacteria of Pakchoi. *Int J Environ Res Public Health* 2016;13:662.
- Yim J-H, Kim K-Y, Chon J-W, Kim D-H, Kim H-S, Choi D-S, Choi I-S, Seo K-H. Incidence, antibiotic susceptibility, and toxin profiles of *Bacillus cereus* sensu lato isolated from Korean fermented soybean products. *J Food Sci* 2015;80: M1266–M1270.
- Yoke-Kqueen C, Learn-Han L, Noorzaleha AS, Son R, Sabrina S, Jiun-Hong S, Chai-Hoon K. Characterization of multiple-antimicrobial-resistant *Salmonella enterica* subsp. *enterica* isolated from indigenous vegetables and poultry in Malaysia. *Lett Appl Microbiol* 2008;46:318–324.
- Yu T, Jiang X, Zhou Q, Wu J, Wu Z. Antimicrobial resistance, class 1 integrons, and horizontal transfer in *Salmonella* isolated from retail food in Henan, China. *J Infect Dev Ctries* 2014;8:705–711.
- Zhang H, Li X, Yang Q, Sun L, Yang X, Zhou M, Deng R, Bi L. Plant growth, antibiotic uptake, and prevalence of antibiotic resistance in an endophytic system of Pakchoi under antibiotic exposure. *Int J Environ Res Public Health* 2017;14:1336.
- Zhu B, Chen Q, Chen S, Zhu YG. Does organically produced lettuce harbor higher abundance of antibiotic resistance genes than conventionally produced? *Environ Int* 2017;98:152–159.
- Zurfluh K, Nüesch-Inderbinen M, Morach M, Zihler Berner A, Hächler H, Stephan R. Extended-spectrum- β -lactamase-producing Enterobacteriaceae isolated from vegetables imported from the Dominican Republic, India, Thailand, and Vietnam. *Appl Environ Microbiol* 2015;81:3115–3120.
- Zurfluh K, Poirel L, Nordmann P, Nüesch-Inderbinen M, Hächler H, Stephan R. Occurrence of the plasmid-borne *mcr-1* colistin resistance gene in extended-spectrum- β -lactamase-producing Enterobacteriaceae in river water and imported vegetable samples in Switzerland. *Antimicrob Agents Chemother* 2016;60:2594–2595.

Address correspondence to:

Christina Susanne Hölzel, PhD

Institute for Animal Breeding and Husbandry

Christian-Albrechts-University Kiel (CAU)

Hermann-Rodewald-Str. 6

24098 Kiel

Germany

E-mail: choelzel@tierzucht.uni-kiel.de