Global Inequality in Maternal Health Care Service Utilization: Implications for Sustainable Development Goals

Sanni Yaya and Bishwajit Ghose*

Abstract

Purpose: Globally, low-middle-income countries continue to account for almost all of the pregnancy-related mortalities that are largely preventable through adequate utilization of essential maternal health care services such as antenatal care (ANC) and skilled birth assistance (SBA). Promoting the use of ANC and SBA services are hindered by numerous policy- and capacity-related barriers along with widespread inequality in utilization of the existing services that further exacerbates the scenario. In an attempt to better understand the geography of inequality in service utilization, we conducted this brief descriptive study by using World Health Organization (WHO) data on ANC and SBA utilization among the member states.

Methods: This was a descriptive study based on open access data on ANC and SBA use between 2012 and 2015 available through the Global Health Observatory of WHO. Country-level data were collected for Asia (41 countries), Africa (35 countries), Europe (35 countries), North America (10 countries), Latin America and the Caribbean (25 countries), and Oceania (16 countries). Cross-country and continent comparisons were made using dot and bar charts.

Results: The overall prevalence of ANC and SBA use were, respectively, 78.17% and 88.33%. Considerable disparities were found in terms of ANC and SBA use across the continents, especially in Asia and Africa. Globally, the poorest performing countries included Afghanistan, Somalia, and South Sudan where more than three-quarters of the women remain deprived of ANC and SBA services during the period of 2012 and 2015. The greatest inequality in ANC use was observed in Africa (9.4% in Somalia and 99.9% in Libya), whereas that of SBA use was observed in Asia (17.8% in Afghanistan vs. 100% in Bahrain). Europe was the most equal of all regions in terms of both ANC (66.8% in Albania vs. 99.7% in Belarus) and SBA (94.4% in Denmark vs. 100% in Lithuania) use.

Conclusion: Although in the majority of countries more than three-quarters of the women receive ANC and SBA services, the extent of intraregional inequality remains overwhelming especially for Asia and Africa. Progress toward maternal health-related targets should be interpreted in terms of the disparities to ensure a more even and sustainable outcome at both national and global level.

Keywords: antenatal care; skilled-birth assistance; health inequality; sustainable development goals

Introduction

Equitable provision of maternal health care services such as antenatal care (ANC) and skilled birth assistance (SBA) comprise a critical component of the entire health system owing to its pivotal role in ensuring safe

motherhood and survival and health of newborns as well the overall well-being of families and communities.^{1–3} Maternal mortality is a leading cause of mortality among women of reproductive age with the prevalence being significantly pronounced in low-resource settings.

School of International Development and Global Studies, Faculty of Social Sciences, University of Ottawa, Ottawa, Canada.
Both authors' contributed equally to this study.

*Address correspondence to: Bishwajit Ghose, PhD, School of International Development and Global Studies, Faculty of Social Sciences, University of Ottawa, 120 University Private, Ottawa K1N 6N5 Canada Ottawa, Canada, E-mail: brammmaputram@gmail.com

© Sanni Yaya and Bishwajit Ghose 2019; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

During past few decades, especially since the 1990s (launching year of the millennium development goals [MDGs]), the global community has witnessed massive undertakings to reverse the situation and also seen unprecedented declines in maternal and child mortality rates.⁴ However, post-MDGs estimation reveals that the progress remains largely uneven across and within regions as high rates of pregnancy-related mortalities and non-/underutilization of essential maternal health care services continue to be disproportionately higher among the most marginalized communities.^{5–7} The widening socioeconomic gradient in maternal health care utilization is a growing concern for low-middle-income countries (LMICs) in pursuit of achieving the sustainable development goals (SDGs).^{7–9}

Current statistics suggest that in high-income countries very few women die from childbirth, and the key maternal health care services, for example, antenatal and professional child birth services are nearly universal.^{10,11} In contrast, LMICs account for ~99% of global maternal mortality¹² with the coverage of SBA being as low as 50% (in sub-Saharan Africa).¹³ The main challenges for promoting maternal health care utilization in resource-limited settings are surrounding those of accessibility and affordability barriers stemming from various infrastructural,¹⁴ skilled human resource,¹⁵ technological¹⁶ and financial¹⁷ issues at health systems level and poor health literacy,¹⁸ self-efficacy,¹⁹ and behavioral²⁰ factors at the individual level. Although much has been achieved in terms of reducing the structural barriers to utilization of care, much remains to be done. With the expiration of MDGs, further initiatives are being taken (e.g., SDGs) toward the goals of promoting maternal and child health through an enhanced focus on addressing inequality in utilization of obstetric care and professional delivery services.

Experience gained from the MDGs provides the basis for realizing that health care inequality is a multidimensional issue that undercuts all development efforts and delays progress.^{8,21–24} This is particularly the case for maternal health care utilization in the LMICs owing to women's inferior socioeconomic status^{25,26} and inadequate opportunities for empowerment.^{4,27} Therefore, addressing inequality in maternal health care, including SBA, is an overarching priority, particularly in the countries that failed to meet the MDGs. Health sectors in both HICs (high-income countries/countries with minimum per capita GNI of \$12,056, as per World Bank) and LMICs need to make strategic approaches to ensure that women's access to the core

maternal health care services is not conditional on their socioeconomic status and pay special attention to the marginalized population who are often unable to uptake services even when no fees are incurred.²⁸ The challenge is clearly of significant magnitude and would require huge input in financial and logistical terms accompanied by firm commitments for action. Addressing health inequality is also reliant on generating and provision of high-quality data to facilitate evidence-based actions and monitoring the progress at national and international levels. We undertook this study based on open-access data as an exemplary effort to show the relative inequality in ANC and SBA prevalence across and within the six continents. Through the descriptive analysis and interpretation in the context of SDGs, this article serves the purpose of informing the actors at local and international level regarding this increasingly relevant issue that deserves more intense research and policy attention.

Methods

Source of data

Data for this study were extracted from Global Health Observatory (GHO), which is popularly known as the gateway of World Health Organization (WHO) to population health data. GHO database is dedicated to providing most recent data on a wide range of health indicators for the WHO member states, including MDGs, infectious diseases, maternal and child nutrition, reproductive health, and health care utilization. The database has been growing in popularity given the user-friendly nature of the system and reliability of the data, which makes it a crucial resource of information for health researchers especially in the developing countries. Data are gathered from a multitude of sources such as national birth and death registration databases, demographic and health surveys, and research projects and publications.²⁹ For this study, we extracted data on two indicators relevant to maternal health care utilization, for example, ANC, and SBA. Years of data collection ranged from 2012 to 2015 and for some countries data were available on ANC only but not SBA, and vice versa.

Data analysis

This was a descriptive study and data analysis involved graphical presentation of the prevalence of utilizing ANC and SBA services. WHO defines ANC as the health services provided by skilled health care professionals to pregnant women and adolescent girls along

Countrywise percentages of ANC and SBA use were plotted side by side for easier comparison using dot and bar plots. Owing to space constraints and better interpretability, we grouped the countries into six regions: Asia, Africa, Europe, Latin America and the Caribbean, North America, and Oceania. Countries with higher and lower than global average of ANC and SBA use were presented as map charts. All charts were prepared with R studio.

Globally, the overall prevalence of ANC and SBA use were, respectively, 78.17% and 88.33%. The prevalence of ANC use among Asian countries was lowest in Afgha-

Among the countries in Latin America and the Caribbean, the lowest prevalence of ANC use was observed

FIG. 2. Prevalence of utilizing ANC and SBA services in Africa.

FIG. 3. Prevalence of utilizing ANC and SBA services in Europe.

for Suriname (66.8%), and that of SBA use in Guatemala (65.5%). ANC and SBA use were universal in Antigua and Trinidad (Fig. 4). In majority of the countries, use of ANC services was more prevalent than that of SBA use.

In Oceania, the prevalence of ANC was highest in Australia (95%) and lowest in Nauru (40.2%). In contrast, SBA use was lowest in Papua New Guinea (40%), and was universal in Palau, Niue, Micronesia, and Cook Islands (Fig. 5).

Figure 6 shows that among the North American nations, Haiti had the lowest prevalence of both ANC and SBA use, and was the country where the prevalence of ANC and SBA use differed most markedly (66.6% vs. 41.7%). ANC and SBA services were near universal in Canada.

Figure 7 indicates that Europe as a region had the highest prevalence of ANC and SBA use followed by Latin America and the Caribbean. Overall, the prevalence of ANC use was lowest in Asia and that of SBA was lowest in Africa. The overall prevalence of SBA use was higher than that of ANC in all continents except North America.

Countries with ANC and SBA coverage above and below global average are presented in Figures 8 and 9, respectively. The majority of the countries that rank below global average fall within the Asian and African region with a few exceptions in Latin America and Oceania.

Discussion

Several interesting findings emerged from the analysis that are worthy of notice. First, the prevalence of ANC use was comparatively lower than SBA. This pattern was observed for both high-middle-income countries and LMIC. At the continental level, however, only North America had a higher prevalence of ANC use than SBA. The difference between the prevalence of ANC and SBA use was most remarkable for Africa—57.85% ANC versus 87.3% SBA. Inadequate-/nonuse of ANC services reduces the opportunities for detecting and addressing the diverse complications that may arise for the mother (hypertension, diabetes, and miscarriage)^{32,33} as well as for fetal surveillance that is crucial for vital examinations such as intrauterine growth retardation.^{34,35} Furthermore, women who do not attend

FIG. 4. Prevalence of utilizing ANC and SBA services in Latin America and the Caribbean.

FIG. 5. Prevalence of utilizing ANC and SBA services in Oceania.

FIG. 6. Prevalence of utilizing ANC and SBA services in North America.

ANC are also less likely to use professional childbirth services and thus altogether bear an increased risk of poor pregnancy outcomes.

Second, the interregional disparity was also more pronounced for ANC use. For instance, among the regions with above average prevalence of ANC and SBA use, the prevalence of ANC ranged from 73.44% in Oceania to 90.68% in Europe. In contrast, that of SBA ranged from 96.31% in Europe and 91.54% in North America. Countries with suboptimal ANC prevalence should, therefore, be given special emphasis for policy and action to enable them to be on track for meeting the maternal and child health-related SDGs.

Third, inequality in the use of both ANC and SBA was most noticeable among countries in Africa and Asia. These are also the regions where the prevalence of both of the indicators was lower than the global average. In majority of the countries in Asia, ANC services are used by more than three-quarter of the women and used almost universally in several countries. In Africa, in contrast, although in majority of the countries more than three-quarter of the births are assisted by a skilled

professional and is near universal in some countries, only a handful of the countries had an ANC prevalence of 75% and higher, but was universal in none. In general, the prevalence of both ANC and SBA use was lower for African countries compared with their Asian countries of similar per capita GDP. For instance, Botswana, South Africa, and Namibia had a lower prevalence of ANC use compared with Asian economies having similar or lower per capita GDP such as Thailand, Sri Lanka, and Bhutan. This is an indication that coverage of ANC at national level is less sensitive to the measures of income in Africa than in Asia. The variation might be accounted for by differences in national macroeconomic objectives as well as health and disease control priorities in these two regions. For this analysis, we did not have data on intracountry disparities on ANC and SBA use; nonetheless, it is assumable that at national level the situation could be far worse among the disadvantaged communities and especially among poorest of the poor.

Last but not least, intercountry disparities were vivid in all the continents especially in Asia and Africa. With nearly four-fifth of the women receiving ANC and SBA services, the overall scenario of ANC and SBA use may seem appropriate at global scale. The implementation period of MDGs has brought about an unprecedented decline in global maternal and child mortality rates. Despite the remarkable progress, high maternal mortality rates continue to persist especially across Asia and Africa where a large proportion of the women do not avail essential maternal health care services such as ANC and SBA. Hence, progress achieved at global level may overshadow the widening inequalities within certain countries that are failing to keep up with the internationally set goals. Unequal progress toward the maternal health care-related MDGs suggests that overall success of the efforts would ultimately rely on eliminating the disparities at all levels. Ensuring progress in an equitable and sustainable manner is, therefore, a necessary imperative that the health policy makers need to prioritize, particularly in the countries that are lagging behind the targets regarding maternal health care.

One distinguishing feature of SDGs in relation to MDGs is its global scope aiming to enhance population health, environment, and development in all countries across the development spectrum. Not surprisingly, in both of the approaches maternal and child health-related goals occupy a pivotal position as improvement in these indicators is crucial to ensuring the overall

success of the agenda. Reducing the global maternal mortality ratio to <70/100,000 by 2030 and ensuring universal access to sexual and reproductive health care represent two hallmark objectives of SDGs. From the perspective of SDGs, the goals 3 (promoting health and well-being), 5 (gender equality and women's empowerment), and 10 (reducing inequality within and across countries) are directly relevant to maternal and health child health and health care, and failure to prioritize these areas can significantly halt overall success in the LMICs (countries with per capita GNI of \$996 to \$3895, as per World Bank). Evidence suggests that access to quality maternal health care has significant implications on quality of life and well-being among mothers^{36,37} which serve as the prerequisite of women's socioeconomic empowerment^{38,39} and promoting gender equality.^{40,41}

In this study, we took advantage of these publicly available data from WHO to explore the situation of inequality in ANC and SBA use across and within continents. Studies of this kind are particularly important as they can inform health professionals and policy makers and facilitate evidence-based interventions to combat inequality at local and global stage. To date, a few comparative analyses of maternal health care have been carried out; however, none has focused on the inequality within and across continents from the perspective of SDGs. Therefore, in this study we presented the WHO data in a more informative manner to better understand the geography of inequality in maternal health care utilization. One important limitation of the study was that we were unable to show the intracountry sociodemographic inequalities due to lack of data. The first step to addressing the inequalities would require identifying the at-risk population through conducting large-scale population surveys to facilitate informed decision making. Future studies should focus on exploring the social and political determinants that underlie the differential progress toward the use of maternal health care services.

Conclusion

Apart from the universal use of ANC and SBA in the high-income countries, overwhelming disparities exist in a large number of countries across Asia and Africa. To promote equality in achieving the maternal health and related targets, progress should be assessed in terms of the disparities at both national and global level. Maternal health care cuts across three core components of SDGs, including population health and well-being, gender equality, and reduction of inequal-

ity within and among countries. Increased international collaboration is necessary to help the low-performing countries to be on track by enhancing both the quality and coverage of maternal health care interventions.

Ethics Statement

Data used in this study were open access and represent country level statistics only; therefore, no institutional review board approval was necessary.

Author Disclosure Statement

No competing financial interests exist.

References

- McNamee P, Ternent L, Hussein J. Barriers in accessing maternal health-care: evidence from low- and middle-income countries. *Expert Rev Pharmacoecon Outcomes Res.* 2009;9:41–48.
- Onta S, Choulagai B, Shrestha B, et al. Perceptions of users and providers on barriers to utilizing skilled birth care in mid- and far-western Nepal: a qualitative study. *Glob Health Action.* 2014;7:24580.
- Choulagai B, Onta S, Subedi N, et al. Barriers to using skilled birth attendants' services in mid- and far-western Nepal: a cross-sectional study. *BMC Int Health Hum Rights.* 2013;13:49.
- Ghose B, Feng D, Tang S, et al. Women's decision-making autonomy and utilisation of maternal healthcare services: results from the Bangladesh Demographic and Health Survey. *BMJ Open.* 2017;7:e017142.
- Wirth M, Sacks E, Delamonica E, et al. "Delivering" on the MDGs?: equity and maternal health in Ghana, Ethiopia and Kenya. *East Afr J Public Health.* 2008;5:133–141.
- Lomazzi M, Borisch B, Laaser U. The Millennium Development Goals: experiences, achievements and what's next. *Glob Health Action.* 2014;7:23695.
- Cohen RL, Alfonso YN, Adam T, et al. Country progress towards the Millennium Development Goals: adjusting for socioeconomic factors reveals greater progress and new challenges. *Glob Health.* 2014;10:67.
- Bobo FT, Yesuf EA, Woldie M. Inequities in utilization of reproductive and maternal health services in Ethiopia. *Int J Equity Health.* 2017;16:105.
- Kachikis A, Moller A-B, Allen T, et al. Equity and intrapartum care by skilled birth attendant globally: protocol for a systematic review. *BMJ Open.* 2018;8:e019922.
- Shaw D, Guise JM, Shah N, et al. Drivers of maternity care in high-income countries: can health systems support woman-centred care? *Lancet Lond Engl.* 2016;388:2282–2295.
- Mariani G, Kasznia-Brown J, Paez D, et al. Improving women's health in low-income and middle-income countries. Part I: challenges and priorities. *Nucl Med Commun.* 2017;38:1019–1023.
- Das R, Biswas S. Eclampsia: the major cause of maternal mortality in Eastern India. *Ethiop J Health Sci.* 2015;25:111–116.
- WHO | Skilled attendants at birth. WHO. Available at www.who.int/gho/maternal_health/skilled_care/skilled_birth_attendance_text/en Accessed September 26, 2018.
- Essendi H, Johnson FA, Madise N, et al. Infrastructural challenges to better health in maternity facilities in rural Kenya: community and healthworker perceptions. *Reprod Health.* 2015;12:103.
- Ranson MK, Chopra M, Atkins S, et al. Priorities for research into human resources for health in low- and middle-income countries. *Bull World Health Organ.* 2010;88:435–443.
- Pagalday-Olivares P, Sjöqvist BA², Adjorodor-van de Beek J, et al. Exploring the feasibility of eHealth solutions to decrease delays in maternal healthcare in remote communities of Ghana. *BMC Med Inform Decis Mak.* 2017;17.
- Kalu-Umeh NN, Sambo MN, Idris SH, et al. Costs and patterns of financing maternal health care services in Rural Communities in Northern Nigeria: evidence for designing national fee exemption policy. *Int J MCH AIDS.* 2013;2:163–172.
- Thomas SD, Mobley SC, Hudgins JL, et al. Conditions and dynamics that impact maternal health literacy among high risk prenatal-interconceptional women. *Int J Environ Res Public Health.* 2018;15. pii:E1383

19. Yaya S, Bishwajit G, Ekholuenetale M. Factors associated with the utilization of institutional delivery services in Bangladesh. *PLoS ONE*. 2017;12:e0171573.
20. Yanagisawa S, Oum S, Wakai S. Determinants of skilled birth attendance in rural Cambodia. *Trop Med Int Health*. 2006;11:238–251.
21. Antai D. Inequalities in under-5 mortality in Nigeria: do ethnicity and socioeconomic position matter? *J Epidemiol*. 2011;21:13–20.
22. Haider MR, Rahman MM, Moinuddin M, et al. Impact of maternal and neonatal health initiatives on inequity in maternal health care utilization in Bangladesh. *PLoS ONE*. 2017;12:e0181408.
23. Yaya S, Bishwajit G, Shah V. Wealth, education and urban-rural inequality and maternal healthcare service usage in Malawi. *BMJ Glob Health*. 2016;1:e000085.
24. Yu W, Bain RE, Mansour S, et al. A cross-sectional ecological study of spatial scale and geographic inequality in access to drinking-water and sanitation. *Int J Equity Health*. 2014;13:113.
25. Makate M, Makate C. The evolution of socioeconomic status-related inequalities in maternal health care utilization: evidence from Zimbabwe, 1994–2011. *Glob Health Res Policy*. 2017;2:1.
26. Mezmur M, Navaneetham K, Letamo G, et al. Socioeconomic inequalities in the uptake of maternal healthcare services in Ethiopia. *BMC Health Serv Res*. 2017;17:367.
27. Sado L, Spaho A, Hotchkiss DR. The influence of women's empowerment on maternal health care utilization: evidence from Albania. *Soc Sci Med*. 2014;114:169–177.
28. Pieterse P, Lodge T. When free healthcare is not free. Corruption and mistrust in Sierra Leone's primary healthcare system immediately prior to the Ebola outbreak. *Int. Health*. 2015;7:400–404.
29. WHO | Global Health Observatory—the one-stop shop for health data. WHO. Available at www.who.int/features/2012/global_health_observatory/en Accessed September 28, 2018).
30. Information NC for B, Pike, USNL of M 8600 R, MD, B & USA, 20894. Introduction. World Health Organization, 2016.
31. WHO | Making pregnancy safer: the critical role of the skilled attendant. WHO Available at https://www.who.int/maternal_child_adolescent/documents/9241591692/en Accessed February 13, 2019.
32. Dave VR, Rana BM, Sonaliya KN, et al. Screening of gestational diabetes and hypertension among antenatal women in rural West India. *Cent Asian J Glob Health*. 2014;3:140.
33. Ali AD, Mehress AA-KO, Al-Adhroey AH, et al. Prevalence and risk factors of gestational diabetes mellitus in Yemen. *Int J Womens Health*. 2016;8:35–41.
34. Sovio U, White IR, Dacey A, et al. Screening for fetal growth restriction with universal third trimester ultrasonography in nulliparous women in the Pregnancy Outcome Prediction (POP) study: a prospective cohort study. *Lancet Lond Engl*. 2015;386:2089–2097.
35. Sharma D, Shastri S, Sharma P. Intrauterine growth restriction: antenatal and postnatal aspects. *Clin Med Insights Pediatr*. 2016;10:67–83.
36. Yeoh PL, Hornetz K, Ahmad Shauki NI, et al. Assessing the extent of adherence to the Recommended Antenatal Care Content in Malaysia: room for Improvement. *PLoS ONE*. 2015;10:e0135301.
37. Baston H. Midwifery basics. Antenatal care—monitoring maternal well-being. *Pract Midwife*. 2003;6:32–35.
38. Onarheim KH, Iversen JH, Bloom DE. Economic benefits of investing in women's health: a systematic review. *PLoS ONE*. 2016;11:e0150120.
39. Mandal M, Muralidharan A, Pappa S. A review of measures of women's empowerment and related gender constructs in family planning and maternal health program evaluations in low- and middle-income countries. *BMC Pregnancy Childbirth*. 2017;17:342.
40. Adjiwanou V, LeGrand T. Gender inequality and the use of maternal healthcare services in rural sub-Saharan Africa. *Health Place*. 2014;29:67–78.
41. Morgan R, Tetui M, Muhumuza Kananura R, et al. Gender dynamics affecting maternal health and health care access and use in Uganda. *Health Policy Plan*. 2017;32:v13–v21.

Cite this article as: Yaya S, Ghose B (2019) Global inequality in maternal health care service utilization: Implications for sustainable development goals, *Health Equity* 3:1, 145–154, DOI: 10.1089/heq.2018.0082.

Abbreviations Used

ANC = antenatal care
GHO = Global Health Observatory
LMIC = low-middle-income countries
MDG = millennium development goal
SBA = skilled birth assistance
SDG = sustainable development goals
WHO = World Health Organization

Publish in *Health Equity*

- Immediate, unrestricted online access
- Rigorous peer review
- Compliance with open access mandates
- Authors retain copyright
- Highly indexed
- Targeted email marketing

liebertpub.com/heq