ORIGINAL ARTICLE

Open Access

Physicians in Myanmar Provide Palliative Care Despite Limited Training and Low Confidence in Their Abilities

Emily Earl-Royal, MD, MPH,^{1,*} Michelle Feltes, MD,¹ Michael A. Gisondi, MD,² Loretta Matheson, PhD,¹ Maung Ohn "Tony" Htoo, MD,³ and Rebecca Walker, MD, MPH¹

Abstract

Background: Patients in low-income and middle-income countries (LMICs) have limited access to palliative care providers. In Myanmar, little is known about physician knowledge of or perceptions about palliative care. An assessment of physician practice and capacity to provide palliative care is needed.

Objective: Our objective was to identify physician practice patterns, knowledge gaps, and confidence in providing palliative and end-of-life care in Myanmar.

Design: This was a cross-sectional survey study.

Setting/Subjects: Participants were physicians practicing in Myanmar who attended the *Myanmar Emergency Medicine Updates Symposium* on November 10 to 11, 2018 in Yangon, Myanmar ($n = 89$).

Measurements: The survey used modified Likert scales to explore four aspects of palliative care practice and training: frequency of patient encounters, confidence in skills, previous training, and perceived importance of formal training.

Results: Study participants were young (median age 27 years old); 89% cared for terminally ill patients monthly, yet 94% reported less than two weeks of training in common palliative care domains. Lack of training significantly correlated with lack of confidence in providing care. Priorities for improving palliative care services in Myanmar include better provider training and medication access.

Conclusions: Despite limited training and low confidence in providing palliative care, physicians in Myanmar are treating patients with palliative needs on a monthly basis. Future palliative care education and advocacy in Myanmar and other LMICs could focus on physician training to improve end-of-life care, increase physician confidence, and reduce barriers to medication access.

Keywords: barriers; confidence; education; low-income and middle-income countries; Myanmar; training

Background

Palliative and end-of-life care are increasingly recognized as basic human rights, consistent with the rights to health care and dignity.^{1–5} However, there is unequal access to palliative care throughout the world, representing a significant health disparity for those with terminal illness. Many patients in low-income and middle-income countries (LMICs) have little or no ac-

cess to palliative care^{5–9} despite clear and compelling needs. For example, LMICs have an increasing prevalence of chronic noncommunicable disease, childhood cancers, and older population demographics, all of which require additional palliative care services.^{10–14}

Myanmar is one such LMIC without broad access to palliative care. Myanmar is located in Southeast Asia and was a British colony from the late 19th century

¹Department of Emergency Medicine, ²The Precision Education and Assessment Research Lab, Department of Emergency Medicine, Stanford University School of Medicine, Palo Alto, California, USA.

³Emergency Department, Parami General Hospital, Yangon, Myanmar.

*Address correspondence to: Emily Earl-Royal, MD, MPH, Department of Emergency Medicine, Stanford University School of Medicine, 900 Welch Road, Suite 350, Palo Alto, CA 94304, USA, E-mail: eearlroyal@gmail.com

until 1948; subsequently, the country was ruled by a military dictatorship until 2011, when it began instituting political and social reforms.¹⁵ As described in the Global Atlas of Palliative Care, Myanmar is a country with “isolated palliative care provision...[and has] palliative care activism that is patchy in scope and not well supported.”⁵ The 2015 Quality of Death Index, based on in-country interviews with palliative care experts, found that Myanmar ranked 76 out of 80 countries in overall quality of death, with low scores in multiple domains including community engagement, palliative and health care environment, human resources, and quality of care.¹² In addition, limited supplies of pain medications and health care funding do not reliably address the needs of their population.⁵

Little is known about Myanmar physicians’ training, confidence, perceived importance, and frequency of providing palliative care. However, an accurate understanding of physician knowledge, skills, and attitudes regarding end-of-life care is fundamental to building Myanmar’s capacity to meet its palliative care needs.

Objective

We examined whether physicians in Myanmar routinely provided palliative care and whether they felt adequately trained to do so. We hypothesized that Myanmar physicians would report high frequency of care, low rates of previous training, low confidence in their skills, and high perceived importance of formal training in palliative care.

Methods

Design

No prior investigation measured the gap between confidence and training needs in palliative care among Myanmar physicians; thus, there were no prior survey instruments on which to base this study. Accordingly, a survey tool was developed based on a literature review and expert opinion. In consultation with a health sciences librarian, literature review was conducted using Google Scholar™ (Mountain View, CA) with the search terms “data + Myanmar + end of life,” “Myanmar + cancer + palliative,” “palliative + Myanmar,” “end of life + Myanmar,” “needs assessment + palliative + Asia,” and “needs assessment + palliative + provider,” and using all databases in Web of Science (Clarivate Analytics, Philadelphia, PA) for “palliative or end of life + Myanmar + Burma or Burmese.”

Study and survey design included experts who are American physicians with training in palliative medi-

cine and medical education research study design, as well as both Myanmar and American physicians with experience in the practice of medicine and medical education in Myanmar. We created our survey instrument using a modified Delphi approach to optimize content and internal structure evidence. The survey was tested among five authors for item generation, survey functionality, matching of item content to the construct, optimal item phrasing, and overall quality control. The survey was screened and edited by multiple physicians practicing in Myanmar to ensure cultural and clinical relevance. The survey was then piloted within the study team before distribution. Pilot results were checked for consistency, providing some evidence of response process validity. The paper survey was then distributed in person to eligible participants after informed consent was obtained verbally. The survey was open for the two-day study period only. No individual identifying information was recorded.

Setting/subjects

This was a cross-sectional survey of physicians practicing in Myanmar. Our roster of eligible participants included attendees at *Myanmar Emergency Medicine Updates Symposium* on November 10 to 11, 2018, in Yangon, Myanmar. Participants at this conference represented a broad mix of unspecialized physicians and specialized physicians who were unlikely to have advanced training in areas of palliative care. Participants were limited to physicians in practice; trainees were excluded. This study was reviewed by Stanford University Institutional Review Board and given exempt status, Protocol No. 48211.

Measurements

Participants were asked to respond to questions about four aspects of palliative medicine using modified Likert scales: frequency of patient encounters, confidence in skills, previous training, and perceived importance of formal training. Participants were also asked to identify the two most important priorities for improving palliative medicine in Myanmar, selected from a list of options; this list included the option “Other” with an open-text response box (Supplementary Appendix SA1, “Palliative Medicine Physician Needs Assessment in Myanmar”).

Data analysis using descriptive statistics was performed in Microsoft Excel 2017 (Redmond, WA) and SAS Enterprise Guide version 8.2 (Cary, NC). Pearson correlations were used to determine significance.

Table 1. Participant Demographics

	N (%)
Total	89
Age, (years) median [IQR]	27 [24–33]
Female	48 (54)
Years in practice	
<5	45 (51)
5–10	28 (31)
11–20	11 (12)
21–30	3 (3)
>30	1 (1)
Type of hospital practice	
Public hospital only	25 (25)
Private hospital only	51 (57)
Both public and private	15 (17)
Medical training	
Unspecialized	59 (66)
Specialized	30 (34)
Emergency	13 (15)
Surgery	5 (6)
Family medicine	3 (3)
Internal medicine	3 (3)
Pediatrics	2 (2)
Administration	1 (1)
Anesthesiology	1 (1)
Infectious disease	1 (1)
Orthopedics	1 (1)
Palliative care training	5 (6)

Results

Eighty-nine participants completed the survey, over half of whom were female ($n=48$, 54%) (Table 1). Most respondents practiced medicine 10 or fewer years ($n=73$, 82%) and worked in private hospitals at

least part of the time ($n=66$, 74%). The majority of respondents were unspecialized practitioners ($n=59$, 66%).

Most of the study participants treated patients with palliative care needs at least monthly (Fig. 1). The most commonly treated patient symptoms included fatigue ($n=82$, 92%), shortness of breath ($n=81$, 91%), and terminal illness ($n=79$, 89%). Just over half of the study participants provide counseling to survivors ($n=49$, 55%). Few participants were very confident providing palliative care (Fig. 2). Over one-third of participants reported a lack of confidence providing care for patients with terminal illness ($n=29$, 34%) or patients at the end of life ($n=31$, 37%).

Most participants reported zero to two weeks of palliative care training in their careers to date (Fig. 3). Fewer than 1 in 10 participants reported more than two weeks of training in managing terminal illness, helping patients prepare for death, counseling survivors, and end-of-life care. Training and confidence were significantly correlated ($p<0.05$) in 14 of the 15 domains of palliative care explored in this study. Most participants ($n=71$, 81%) rated formal training as “important” or “very important” for all 15 domains (Fig. 4).

The top priorities for improving palliative care services in Myanmar were medication access and cost, health care access, and provider training (Fig. 5).

FIG. 1. Frequency of treating patients with palliative care.

FIG. 2. Level of confidence the participants feel in each area.

Discussion

This study of physician perspectives on palliative care in Myanmar found that most participants provided palliative services at least monthly, most reported little training or confidence in palliative care skills, and nearly all believed that palliative care training is impor-

tant. It was striking that half of all participants provided end-of-life care without any palliative care training. Top priorities for improving palliative care in Myanmar included barriers to medications and lack of provider training. Our findings highlight the need for improved medical education and health

FIG. 3. Level of training the participants have received in each area.

FIG. 4. Level of importance the participants assign to receiving training in each area.

advocacy in palliative care in Myanmar and similar LMICs, namely provider training and medication access and cost.¹⁶

Although previous studies examined patient-reported needs for palliative care in Southeast Asia and provider knowledge of and attitudes about geriatric palliative care in Vietnam, ours is the first study to report Myanmar physician perspectives on end-of-life and palliative care.^{17,18} Although Myanmar's current palliative care needs are already substantial, the demand for palliative services will only increase over time due to projected changes in demographics.^{10–14} The country's population currently skews very young: 27% of Myanmar's population is <15 years old and only 6% are of age >65 years.¹⁸ However, Myanmar's population >65

years old is expected to triple after 2050, which will drastically increase its need for elder and palliative care.¹⁹

We found strikingly low levels of provider confidence and training in key palliative care domains including goals of care, preparations for death, and bereavement counseling for survivors; these gaps extended to conditions of cancer, AIDS, and terminal delirium (Figs. 2 and 3). Within the limitations of our study, confidence in a specific domain is our best measure of a physician's ability to provide satisfactory care. Our data show that Myanmar physicians who reported previous training in certain areas of palliative care were statistically more likely to report confidence in their practice. That is to say, palliative care training correlates with confidence.

FIG. 5. Top priorities for improving palliative care in Myanmar.

Our finding is similar to a study of health care providers in Vietnam that found that “good knowledge” about palliative care was associated with a “positive attitude” about it.¹⁸ Based on our findings, we advocate for better training of Myanmar physicians in the most common domains of palliative and end-of-life care.

Although few physicians in Myanmar are trained in palliative care, local organizations are attempting to address knowledge gaps. The Asia-Pacific Hospice Palliative Care Network and the LIEN Foundation in Singapore are building educational programs in the region, which have trained 28 providers from Myanmar through a partnership with the Myanmar Medical Association.²⁰ “Life Asked Death,” a documentary about palliative care needs in Southeast Asia, is raising awareness about end-of-life care in Myanmar.²¹ Critical training in palliative care for Myanmar physicians still depends on foreign aid and the nonprofit sector, including the United States Agency for International Development, *Medicins Sans Frontieres*, and the U Hla Tung Hospice Foundation.^{22,23}

Three themes emerged from our study as priorities for future interventions in Myanmar: provider training, medication access, and medication cost. These themes echo previous studies describing barriers to palliative care in developing countries.^{19,22} Our findings regarding medication access correlate with a longstanding lack of pain medication in LMICs and restrictions on chronic use. Opioid access is unequal around the globe, with 15% of the world’s population accounting for 93% of opioid use.^{6,10,24–29} Ironically, Myanmar has a severe shortage of opioid analgesics despite producing much of the world’s opium supply.³⁰

Our study has several limitations. Our convenience sample may not be representative of physician perspectives throughout Myanmar, and our findings may not generalize to other LMICs. Participants in this study may be younger than average physician age in Myanmar. Although we enrolled participants at an emergency medicine conference, most of the physicians were unspecialized (66%). Similar to other LMICs, many of the “newer” medical specialties, such as emergency medicine and palliative care, are not well established in Myanmar. Our responses may have been affected by sponsorship bias, as some study authors also served as conference faculty. In addition, desirability bias is common among survey respondents. Finally, palliative care is a broad and complex discipline; some important domains related to teamwork, communication, grief, and cultural differences may not have been adequately captured in our survey.

Conclusion

In summary, Myanmar physicians in our study reported lack of training and minimal confidence across common palliative care domains. Future education and policy interventions should focus on provider training, medication access, and medication cost for Myanmar to better provide palliative and end-of-life care for its citizens.

Authorship Confirmation Statement

The authors affirm that they have each met criteria for authorship as defined by the International Committee of Medical Journal Editors.

Funding Information

No funding was received.

Author Disclosure Statement

No competing financial interests exist.

Supplementary Material

Supplementary Appendix SA1

References

1. Brennan F: Palliative care as an international human right. *J Pain Symptom Manage* 2007;33:494–499.
2. Gwyther L: Palliative care as a human right. *Palliative Care* 2014;6:116–121.
3. Gwyther L, Brennan F, Harding R: Advancing palliative care as a human right. *J Pain Symptom Manage* 2009;38:767–774.
4. Radbruch L, de Lima L, Lohmann D, et al.: The Prague charter: Urging governments to relieve suffering and ensure the right to palliative care. *Palliat Med* 2013;27:101–102.
5. Connor SR, Sepulveda Bermedo MC, Worldwide Palliative Care Alliance, World Health Organization: Global atlas of palliative care at the end of life. https://www.who.int/nmh/Global_Atlas_of_Palliative_Care.pdf. 2014. (Last accessed June 10, 2020).
6. Knaul FM, Farmer PE, Krakauer EL, et al.: Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: The Lancet Commission report. *Lancet* 2018;391:1391–1454.
7. Woodward M: A consensus plan for action to improve access to cancer care in the Association of Southeast Asian Nations (ASEAN) Region. *Asian Pac J Cancer Prev* 2014;15:8521–8526.
8. De Lima L: Palliative care and pain treatment in the global health agenda. *Pain* 2015;156 Suppl 1:S115–S118.
9. Odonkor CA, Osei-Bonsu E, Tetteh O, et al.: Minding the gaps in cancer pain management education: A multicenter study of clinical residents and fellows in a low- versus high-resource setting. *J Glob Oncol* 2016;2:387–396.
10. Basu A, Mittag-Leffler BN, Miller K: Palliative care in low- and medium-resource countries. *Cancer J* 2013;19:410–413.
11. Narain JP, Garg R, Fric A: Non-communicable diseases in the South-East Asia region: Burden, strategies and opportunities. *Natl Med J India* 2011; 24:8.
12. Lewington J, Namukwaya E, Limoges J, et al.: Provision of palliative care for life-limiting disease in a low income country national hospital setting: How much is needed? *BMJ Support Palliat Care* 2012;2:140–144.
13. Caruso Brown AE, Howard SC, Baker JN, et al.: Reported availability and gaps of pediatric palliative care in low- and middle-income countries: A systematic review of published data. *J Palliat Med* 2014;17:1369–1383.
14. Downing J, Powell RA, Marston J, et al.: Children’s palliative care in low- and middle-income countries. *Arch Dis Child* 2016;101:85–90.
15. East Asia/South east Asia:: Burma—The World Factbook-Central Intelligence Agency. <https://www.cia.gov/library/publications/the-world-factbook/geos/bm.html> (Last accessed July 26, 2019).

16. Pastrana T, Vallath N, Mastrojohn J, et al.: Disparities in the contribution of low- and middle-income countries to palliative care research. *J Pain Symptom Manage* 2010;39:54–68.
17. Molassiotis A, Yates P, Li Q, et al.: Mapping unmet supportive care needs, quality-of-life perceptions and current symptoms in cancer survivors across the Asia-Pacific region: Results from the International STEP Study. *Ann Oncol* 2017;28:2552–2558.
18. Vu HTT, Nguyen LH, Nguyen TX, et al.: Knowledge and attitude toward geriatric palliative care among health professionals in Vietnam. *Int J Environ Res Public Health* 2019 25;16.
19. Myanmar Population Data: Population Reference Bureau. 2018. www.worldpopdata.org/map (Last accessed June 10, 2020).
20. What does the project entail?. Asia Pacific hospice palliative care network: LIEN collaborative. <http://aphn.org/lien-collaborative-for-palliative-care/what-does-the-project-entail/> (Last accessed July 25, 2019).
21. Hill M: Life asked death|developing palliative care in Asia. Moonshine movies. 2016. <http://lifeaskeddeath.com/> (Last accessed July 29, 2019).
22. Wright M, Wood J, Lynch T, Clark D: Mapping levels of palliative care development: A global view. *J Pain Symptom Manage* 2008;35:469–485.
23. Wright M, Lynch T, Clark D: A review of donor organisations that support palliative care development in five world regions: A report prepared for the Open Society Institute. www.icpcn.org/wp-content/uploads/2013/07/A-review-of-donor-organisations-supporting-palliative-care.pdf. 2008. (Last accessed July 25, 2019).
24. Public health dimension of the world drug problem including in the context of the Special Session of the United Nations General Assembly on the World Drug Problem, to be held in 2016. World Health Organization, Executive Board. http://apps.who.int/gb/ebwha/pdf_files/EB138/B138_11-en.pdf. 2016. (Last accessed July 26, 2019).
25. Cleary J, Radbruch L, Torode J, Cherny NI: Formulary availability and regulatory barriers to accessibility of opioids for cancer pain in Asia: A report from the Global Opioid Policy Initiative (GOPI). *Ann Oncol* 2013; 24 Suppl 11:xi24–xi32.
26. Spruyt O: The status of palliative care in the Asia-Pacific region. *Asia Pac J* 2018;5:12–14.
27. Gladstone R: Much of world suffers not from abuse of painkillers, but absence of them. *The New York Times*. <https://www.nytimes.com/2016/05/18/world/much-of-world-suffers-not-from-abuse-of-painkillers-but-absence-of-them.html>. 2016. (Last accessed July 26, 2019).
28. Vijayan R, Afshan G, Bashir K, et al.: Tramadol: A valuable treatment for pain in Southeast Asian countries. *J Pain Res* 2018;11:2567–2575.
29. Gilson AM, Maurer MA, Lebaron VT, et al.: Multivariate analysis of countries' government and health-care system influences on opioid availability for cancer pain relief and palliative care: More than a function of human development. *Palliat Med* 2013;27:105–114.
30. Narcotic Drugs—Estimated World Requirements for 2019—Statistics for 2017. Vienna: United Nations. https://www.incb.org/documents/Narcotic-Drugs/Technical-Publications/2018/INCB-Narcotics_Drugs_Technical_Publication_2018.pdf. 2019. (Last accessed July 28, 2019).

Cite this article as: Earl-Royal E, Feltes M, Gisondi MA, Matheson L, Htoo MOT, and Walker R (2020) Physicians in Myanmar provide palliative care despite limited training and low confidence in their abilities, *Palliative Medicine Reports* 1:1, 314–320, DOI: 10.1089/pmr.2020.0090.

Abbreviation Used

LMICs = low-income and middle-income countries

Publish in *Palliative Medicine Reports*

- Immediate, unrestricted online access
- Rigorous peer review
- Compliance with open access mandates
- Authors retain copyright
- Highly indexed
- Targeted email marketing

liebertpub.com/pmr