Feline-Human Zoonosis Transmission in North Africa: A Systematic Review

Breck Peterson and Amber N. Barnes

Abstract

Throughout human history, domestic animal species have represented a unique zoonotic disease risk for the transmission of pathogens ranging from viral, bacterial, parasitic, and fungal. In North Africa, cats have a particularly long record and occupy a specialized niche within many communities. This systematic review was conducted to analyze the current and historical literature documenting the breadth and variety of zoonoses in North Africa, specifically relating to the domesticated feline. Multiple electronic databases were searched on January 16, 2019, for published reports on feline zoonoses in North Africa. A total of 76 studies met the inclusion criteria for a full assessment. Articles selected for the review ranged in publication dates from 1939 to 2019 and included a case study, cross-sectional surveys, genomic analyses, and a book chapter. The most commonly studied pathogen was *Toxoplasma gondii* ($n=17$) followed by a variety of helminths ($n=10$). Of the countries in the target region, most publications were of studies conducted in Egypt ($n=53$) followed by Tunisia ($n=12$), Algeria ($n=11$), Morocco ($n=5$), and Libya ($n=3$). The results of this review identify a variety of viral, bacterial, fungal, and parasitic zoonotic diseases associated with cats in North Africa, ranging from historically endemic diseases in both human and animal populations in the region, to emerging infections with recent confirmatory diagnoses. This review describes reported feline zoonoses in North Africa and provides recommendations for their prevention and control. In addition to vaccination campaigns for domesticated felines and postexposure prophylaxis for humans, prompt veterinary and medical care of exposure risks and subsequent infections are essential in limiting the zoonotic disease burden in North African communities of humans and cats.

Keywords: zoonotic disease, North Africa, One Health, veterinary, feline, global health

Introduction

AS LONG AS ANIMALS have been living alongside humans, domesticated species have represented a unique risk to humans in the transmission of pathogens ranging from viral, bacterial, parasitic, and fungal. Zoonoses are infectious diseases that spread between animals and humans, and account for nearly 60% of all infectious disease in humans (Salayer et al. 2017). Losses resulting from zoonotic disease span animal and human health, productivity and income for households and communities, and costs associated with human and veterinary treatment (Torgerson and Macpherson 2011). In low-resource countries, zoonotic disease accounts for almost half of all human deaths and make up a significant

proportion of disability-adjusted life-years within populations (Torgerson and Macpherson 2011, Salayer et al. 2017). Quantifying the global burden of zoonoses is difficult, particularly in North Africa. However, this region represents a substantial portion of the total zoonotic cases of helminthiasis, leishmaniasis, and leprosy (Hotez et al. 2012).

Research into zoonotic disease risks from companion animals have often focused on canines (Macpherson 2012). Yet the feline has a long history in North Africa and the Middle East, and occupies a unique niche in these communities. Complete domestication of the cat likely occurred in Egypt just 4000 years ago (Faure and Kitchener 2009). In addition to skeletal remains as archaeological evidence, depictions of cats through clay and stone figurines from Syria, Turkey, and

Department of Public Health, University of North Florida, Jacksonville, Florida, USA.

© Breck Peterson and Amber N. Barnes 2020; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are cited.

Israel, as well as hieroglyphs and decorated cat mummy votive offerings indicate a value of the tamed feline beyond simple rat-catchers and pelt sources (Faure and Kitchener 2009).

In the 1480s, a European traveler wrote about a shelter for street cats in Damascus, describing the phenomenon as something unique to Muslim societies (Alkhateeb Shehada 2012). The shelter had been commemorated in honor of an Islamic tradition, with popular oral versions of the story relating the birth of a litter of kittens on the Prophet Mohammed's robe, who chose to abandon his garment rather than disturb the mother cat (Alkhateeb Shehada 2012). Other histories relating the cat and the prophet are written in the canonical literature, such as the story of a woman being punished for mistreatment of a cat (Sahih al-Bukhari 3318), the disapproval of selling cats and dogs for profit (Sahih Muslim 3808), and of the prophet sanctioning an apostle to perform ablutions with water shared with a cat for drinking (Sunan Abu-Dawud 0075).

Cats are a popular pet globally and there are more domesticated felines than domesticated dogs (Vitale et al. 2019). In 2018, a quarter of all households in the United States owned at least one cat (American Veterinary Medical Association 2018). However, feline contact can put humans and other animals at risk for zoonotic disease transmission (Gerhold and Jessup 2013). For instance, cats are implicated in nearly 40% of all animal bites in the United States, with up to 80% of these bites becoming infected (Kravetz and Federman 2002). In addition to the risk of wound infection and complications such as abscess or septic arthritis, cat bites and scratches are also implicated in the spread of diseases such as rabies, cat-scratch disease (CSD), tularemia, and plague (Kravetz and Federman 2002).

In addition, feline zoonotic transmission may occur through contact with infected saliva or other excretions, contaminated vehicles like food, water, or fomites, and shared vectors or environmental exposures (Tuzio et al. 2005). These diseases and transmission routes are often associated with stray, feral, or community-owned cats, which may be less likely to receive preventive care or veterinary attention (Maia et al. 2014). The aim of this review is to identify, summarize, and describe the published literature linking felines to zoonoses in North Africa and illustrate opportunities for One Health prevention and control strategies that protect the health of humans and cats.

Materials and Methods

Search strategy and selection criteria

Electronic databases were searched on January 16, 2019, for publications on feline zoonoses in North Africa. Databases included the following: ABI/Inform Collection, Agricola, Agriculture Collection, Aquatic Sciences Collection, Biological Sciences, BioOne Complete, Biosis, CABI, Environment Complete, Environmental Science Collection, Environmental Studies and Policy, Google Scholar, Health and Medicine, MEDLINE, Nursing and Allied Health, PubMed, SAGE Journals Online, ScienceDirect, SpringerLink Journals, TOXLINE, Web of Science Core Collection, and Zoological Record. There were no date restrictions, but publications were limited to English or French due to language abilities of the reviewers. The countries of Algeria,

Egypt, Libya, Morocco, and Tunisia were selected based on the World Bank's classification of countries that comprise their Middle East and North Africa region (World Bank 2019).

Key words and search terms were developed from larger categories for the host (cat*, feline, pet*, companion animal*, kitten*, and kitty), routes of zoonotic exposure (transmission, respiratory, enteric, fecal-oral, gastrointestinal, airborne, direct contact, bite*, scratch*, wound*, saliva, contact, feces, exposure*, zoonoses, zoonosis, zoonotic, fecal, and urine), and location of interest (Algeria*, Egypt*, Libya*, Morocco*, Tunisia*, North Africa*, and MENA). These terms were used in the creation of search strings that included wildcards (*) and Boolean operators to try to capture all variations and pluralities, based upon the search principles of each database. Each database was examined for publications that contained at least one search term from each category in the title and/or abstract. Search strings included the use of OR between terms within the category and were combined using AND to make sure that the results included the correct host, a route of zoonotic exposure, and a location of interest such as the following:

1. Cat[tiab] OR cats[tiab] OR feline*[tiab] OR cats[MeSH] OR pet[tiab] OR Pets[MeSH] OR "companion animal"[tiab] OR kitten*[tiab] OR kitty[tiab]) AND
2. (Transmission[tiab] OR respiratory[tiab] OR enteric[tiab] OR "fecal oral"[tiab] OR gastrointestinal[tiab] OR airborne[tiab] OR "direct contact"[tiab] OR bite*[tiab] OR scratch*[tiab] OR wound*[tiab] OR saliva[tiab] OR contact*[tiab] OR feces[tiab] OR exposure*[tiab] OR zoonoses[MeSH] OR zoonotic[tiab] OR zoonosis[tiab] OR fecal[tiab] OR urine[tiab]) AND
3. (Algeria*[tiab] OR Egypt*[tiab] OR Libya*[tiab] OR Morocco*[tiab] OR Tunisia*[tiab] OR "North Africa"[tiab] OR "North African"[tiab] OR "North Africans"[tiab] OR "MENA"[tiab])

Publications were included if there was documentation of zoonoses or reverse zoonoses with feline participation in transmission within or in reference to one or more North African country. Only literature that described cases of feline-associated zoonotic transmission or in which the authors described a strong likelihood of transmission were included. Publications were excluded if the study or report referenced the zoonotic disease in only an animal or human population without discussion of confirmed zoonotic transmission or potential exposure risk from cat contact. Broad overviews and reviews were excluded if they did not specifically provide case reports or study details on feline-human zoonosis transmission.

Data screening

Guidelines from the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (<http://prisma-statement.org/PRISMAStatement/FlowDiagram>) were followed (Moher et al. 2015). The initial search results combined from all databases ($n=2820$) were reduced once duplicates were removed. The nonduplicate results ($n=1270$) were then screened for categories (host, route of zoonotic exposure, and location of interest) in the title and abstract of the document. When it was not clear from reading this section

of the publication alone, the authors erred on the side of caution and included it for the next round of screening. After exclusion from title and abstract review, 365 works remained for a full-text assessment. The authors first worked independently to read over each accessible document and then merged their findings to form a collective and unanimous decision on each article. Full-text works were excluded ($n=289$) based on being a language other than French or English, a duplicate copy that was not discovered using citation management software, inaccessibility, no reference to zoonoses, cats, or cat/human transmission, the study location being outside of North Africa, or the article being a broad overview without specific instances of feline zoonotic transmission. In the end, the search produced 76 studies for review (Fig. 1).

Results

Articles selected for the review ranged in publication dates from 1939 to 2019 and included a case study, cross-sectional surveys, genomic analyses, and a book chapter. The most commonly studied pathogen was *Toxoplasma gondii* ($n=17$) followed by various helminths ($n=10$). Of the countries in the target region, most publications were of studies conducted in Egypt ($n=53$) followed by Tunisia ($n=12$), Algeria ($n=11$), Morocco ($n=5$), and Libya ($n=3$) (Fig. 2). In human studies, the most common study population was patients presenting with clinical signs of infectious disease ($n=15$), followed by pregnant women ($n=11$). Only four studies specified children in their sample population, and only five

involved populations with prior animal contact or occupational exposure to animals. In animal studies, many species beyond cats were researched, including other companion animals, livestock, poultry, and wildlife.

Zoonotic diseases included in review

The results of this review identify a variety of viral, bacterial, fungal, and parasitic zoonotic diseases associated with cats in North Africa (Table 1). Articles in this review that discussed viral infections include studies on rabies, rotavirus, avian influenza, and hepatitis E. In regard to bacterial infections, the included studies examined *Bartonella* spp., *Brucella* spp., *Campylobacter* spp., *Coxiella burnetii*, pathogenic *Escherichia coli*, *Helicobacter pylori*, *Klebsiella* spp., *Leptospira* spp., *Mycobacterium avium* subsp. *paratuberculosis*, *Pasteurella* spp., *Rickettsia* spp., *Salmonella*, *Shigella* spp., *Staphylococcus* spp., and *Streptococcus* spp. Seven studies researched fungal infections, six within the umbrella term of “dermatophytes,” including *Trichophyton violaceum*, *T. rubrum*, *T. tonsurans*, *T. mentagrophytes*, *T. verrucosum*, *T. terrestre*, *Microsporum canis*, *M. gypseum*, *M. audouinii*, *M. praecox*, and *Epidermophyton floccosum*, and one on recently re-categorized microsporidia. Many of the included research articles studied parasitic infections in North African cats that pose a risk to humans, which were further categorized as helminths and unicellular parasites. Among the helminths studied, over 17 different genera, including cestodes, trematodes, and nematodes, are included in

FIG. 1. North African countries included in review with shading to correspond with published studies. Map created in ArcMap 10.6 (ESRI, Redlands, CA); no copyrighted material was used. ESRI, Environmental Systems Research Institute.

FIG. 2. Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram demonstrating selected studies for review.

Adapted from: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7)

this review. Additional research on unicellular parasites among the studies in this review included *Cryptosporidium* spp., *Entamoeba histolytica*, *Giardia* spp., *Leishmania*, and *Toxoplasma gondii*.

Discussion

North Africa is currently faced with many challenging zoonotic diseases such as echinococcosis in Tunisia, rickettsiosis in Algeria, campylobacteriosis in Egypt, rabies in Morocco, and brucellosis in Libya (Majorowski et al. 2005, Ahmed et al. 2010, Mouffok et al. 2011, Ducrotoy et al. 2015, Helmy et al. 2017). The identified pathogens in this review range from historically endemic diseases in both human and animal populations in the region, to emerging and reemerging infections of veterinary and public health importance. Thus, a review of documented feline-associated zoonoses in these countries provides a valuable tool for comparison and retrospective determination of disease status and risk in cats and humans in North Africa.

Viral infections

Historically, rabies in this region is well recorded and recognized. In 1863, rabies was reported in 2 cats, 5 humans, and 17 dogs in Algeria by the Veterinary Surgeon Delacroix (Matter et al. 2004). In 1917, 5% of reported animal bite victims in Algeria were of cat origin (Matter et al. 2004). In 1933, cats were responsible for 117 cases of human rabies exposure in Algeria, and from 1961 to 1980, in Tunisia, cats

were responsible for 8% of treatment sought for human rabies exposure (Matter et al. 2004). Yet, animal rabies vaccination records appear primarily only available for dogs (Matter et al. 2004). Despite this, records from 1970 to 1979 in Algeria show 122 of 333 tested cats (36.6%) were positive for rabies accounting for 10.7% of all positive animal cases during that time (Benelmouffok et al. 1982). This number increased the following year in 1980 when 48.3% of tested cats were positive for rabies (Benelmouffok et al. 1982). The researcher attributed this rise to the difficulty of capturing cats to vaccinate for rabies and that cat bites are typically serious injuries (Benelmouffok et al. 1982).

Between 1976 and 1982, in Morocco, 4.3% of animal rabies cases were in cats (Bakkali 1985). Cats were the origin of 640 rabies postexposure treatments out of nearly 120,000 patients seeking post-exposure prophylaxis (PEP) in Egypt in 2000. In the same year, over 70,000 stray cats were culled to reduce risk of rabies transmission to humans and other animals (Matter et al. 2004). However, recent data on the number of confirmed rabies cases in humans and animals, as well as the number of those vaccinated against or treated for exposure are lacking in this region compared to the overall global numbers. The lack of current data on rabies infection in the North African region, particularly with a breakdown by animal species, does not indicate a decrease in the public health threat surrounding this pathogen, but rather a potential gap in funding and/or research priority. Many global health and veterinary organizations focus on dog-mediated rabies infection and advocate for

TABLE 1. CHARACTERISTICS OF STUDIES ON FELINE ZOONOSES IN NORTH AFRICA

Author	Year ^a	Pathogen	Location	Animal population	Human population
Abdel-Moein and Samir	2011	<i>Staphylococcus aureus</i>	Egypt	Cats, cattle, buffalo, sheep, goat, camel	Dermatology patients with skin lesions and history of animal contact
Abdel-Raouf et al.	2014	<i>Helicobacter pylori</i>	Egypt	Pet and stray cats and dogs	Healthy and sick persons associated with each animal
Aboueisha and El-Mahallawy	2013	Dermatophytes	Egypt	—	Patients diagnosed with dermatophyte lesions
Ahmed et al.	2014	<i>Toxoplasma gondii</i>	Egypt	Sheep	Pregnant women
Al-Herrawy and Gad	2016	Microsporidia	Egypt	Dogs, cats, rabbits, cattle, buffaloes, sheep, goats, donkeys and pigs	—
Al-Kappany et al.	2010	<i>T. gondii</i>	Egypt	Stray cats	—
Al-Kappany et al.	2010	<i>T. gondii</i>	Egypt	Stray cats	—
Arafa et al.	1978	Helminths	Egypt	Stray cats	—
Arrouji et al.	1991	Rabies	Tunisia	Cats, dogs, cows, horses, sheep, and rodents	—
Azzag et al.	2012	<i>Bartonella</i> spp.	Algeria	Stray cats	—
Bakkali	1985	Rabies	Morocco	Cats, dogs, cows, horses, sheep, mules, pigs, donkeys, goats, camels	—
Bassiony et al.	2016	<i>T. gondii</i>	Egypt	—	Pregnant women
Fredj et al.	2013	<i>Rotavirus</i>	Tunisia	—	Case study of infected infant
Benelmouffok et al.	1982	Rabies	Algeria	Dogs, cats, cattle, rats, foxes, jackals, sheep, horses, donkeys, rabbits, guinea pigs, monkeys, pigs, camels	—
Benhamiche	1977	Rabies	Algeria	Dogs, cats, cows, foxes, jackals, rats, “other animals”	—
Benmezad et al.	2012	Dermatophytes	Algeria	—	Hospital outpatients presenting with skin lesions
Bessas et al.	2016	<i>Bartonella</i> spp.; <i>Coxiella</i>	Algeria	Stray dogs and cats	—
Boudebouch et al.	2011	<i>Bartonella</i> spp.; <i>Rickettsia</i> spp.	Morocco	Fleas collected from cats, dogs, sheep, and goats	—
Boumhil et al.	2010	Dermatophytes	Morocco	—	Hospital outpatients presenting with skin lesions
Byomi et al.	2018	<i>T. gondii</i>	Egypt	Sheep, cattle	Women 18–40 years of age
Childs et al.	1995	<i>Bartonella</i> spp.	Egypt	Cats	—
ElAshmawy and Ali	2016	Dermatophytes	Egypt	Pet cats, horses, and calves	—
El-Dakhly et al.	2017	Helminths	Egypt	Stray cats	—
El-Gamal et al.	2013	<i>T. gondii</i>	Egypt	—	Pregnant women
El-Gohary	1998	<i>Campylobacter</i> sp.	Egypt	Cattle, horse, sheep, dogs, goats, cats, turkey, pigeons	Hospital outpatients presenting with GI symptoms

(continued)

TABLE 1. (CONTINUED)

Author	Year ^a	Pathogen	Location	Animal population	Human population
El-Hohary and Abdel-Latif	1998	Cryptosporidia	Egypt	Dog, cat, cow, sheep, pigeon, chicken, goat	Hospital outpatients presenting with GI symptoms
El-Jakee	1991	<i>Campylobacter</i> sp.	Egypt	Dogs, cats	Infants
Elkassas et al.	2018	<i>H. pylori</i> ; <i>Escherichia coli</i>	Egypt	Dogs and cats	Healthy and GI symptomatic persons
El-Menyawe and Abdel Rahman	2007	<i>Toxocara</i>	Egypt	Stray and pet cats and dogs	—
Elmonir et al.	2017	<i>T. gondii</i>	Egypt	Stray cats	Pregnant women
El-Sayed et al.	2013	Avian influenza	Egypt	Cattle, buffalo, sheep, goats, horses, donkey, swine, sewage rats, stray dogs, and stray cats	Humans
El-Seify et al.	2017	Helminths	Egypt	Stray cats	—
El-Shabrawy and Imam	1978	Helminths	Egypt	Stray cats	—
El-Shqanqery et al.	2017	<i>T. gondii</i>	Egypt	—	Pregnant women
Fakhfakh et al.	2013	<i>T. gondii</i>	Tunisia	—	Pregnant women
Felt et al.	2011	<i>Leptospira</i> sp.	Egypt	Rats, dogs, sheep, buffaloes, cattle, donkeys, weasels, and cats	—
Gautret et al.	2010	Rabies	Algeria; Egypt; Libya; Morocco; Tunisia	—	International travelers seeking treatment for animal-related injuries in France
Gharsa et al.	2015	<i>S. aureus</i>	Tunisia	Cow, goat, dog, and cats	—
Ghoneim et al.	2013	Avian influenza	Egypt	Stray dogs and cats, donkeys, rats	—
Ghoneim et al.	1980	<i>S. aureus</i> ; <i>Streptococcus</i> spp.; <i>E. coli</i> ; <i>Salmonella</i> spp.; <i>Shigella</i> spp.; <i>Pasteurella</i> spp.; <i>Klebsiella</i> spp.	Egypt	Stray cats	—
Grami et al.	2013	<i>E. coli</i>	Tunisia	Chickens, pet dogs, and pet cats	—
Hafez Hassanain et al.	2018	<i>T. gondii</i>	Egypt	—	Pregnant women
Haggag et al.	2016	<i>H. pylori</i>	Egypt	Cattle, buffalo, cat, and dog	Healthy and GI symptomatic persons
Hasslinger et al.	1988	Helminths	Egypt	Stray cats	—
Ibrahim et al.	2017	<i>T. gondii</i>	Egypt	Sheep	Pregnant women
Ibrahim et al.	2006	<i>T. gondii</i>	Egypt	Cattle, sheep, camel	Pregnant women
Kaabia and Letaief	2009	Coxiella	Tunisia	—	—
Kaal et al.	2006	Fleas	Libya	Sheep, goats, cattle, camels, horses, rabbits, gazelle, rats, dogs, and cats	Farm workers

(continued)

TABLE 1. (CONTINUED)

Author	Year ^a	Pathogen	Location	Animal population	Human population
Khalafalla Khalifa	2011 1999	Helminths Giardia	Egypt Egypt	Stray cats Stray dogs and cats; rats and mice	— Children 1–10 years of age with diarrhea
Khalil	1977	Toxocara	Egypt	Stray dogs and cats	School children; hospital patients with clinical signs
Khayeche et al.	2014	<i>T. gondii</i>	Tunisia	Sheep	—
Kuntz and Chandler	1956	Helminths	Egypt	Pet cats, dogs, foxes, shrews, rats, and birds	—
Matter et al.	2004	Rabies	Algeria; Egypt; Libya; Morocco; Tunisia	—	—
Messerer et al.	2014	<i>T. gondii</i>	Tunisia	—	Pregnant women
Morsy and El Seoud Abou	1994	Leishmania	Algeria	Pet and stray cats	Primary care outpatients
Morsy et al.	1988	Leishmania	Egypt	Stray dogs and cats; rats and mice	—
Mouffok et al.	2008	Rickettsia	Algeria	—	Patients with clinical signs of rickettsioses
Moustafa Reda and Mostafa	1989 1995	<i>Campylobacter</i> sp.; <i>Salmonella</i> spp. <i>Bartonella</i> spp.	Egypt Egypt	Housed cats and stray dogs	Ophthalmic clinic patients
Rifaat et al.	1976	<i>T. gondii</i>	Egypt	Stray cats	—
Rifaat et al.	1981	<i>T. gondii</i> ; Entamoeba; Helminths	Egypt	Stray cats	—
Romia et al.	1988	<i>T. gondii</i>	Egypt	—	Governorate residents
Salem et al.	2019	Mycobacterium	Egypt	Cattle, camels, cats, rats	—
Salem et al.	2013	<i>E. coli</i>	Tunisia	Pet dogs and cats	Case study of woman with cat bite
Sbai et al.	2015	Pasteurella	Tunisia	—	—
Siam et al.	1973	<i>Staphylococcus</i> spp.; <i>Streptococcus</i> spp.; <i>Salmonella</i> spp.	Egypt	Pet cats, stray dogs	—
Stoszek et al.	2006	Hepatitis E	Egypt	—	Pregnant women
Waly et al.	2017	Dermatophytes	Egypt	Pet cats and dogs	—
Wareth et al.	2017	Brucella	Egypt	Cattle, farm dog, and farm cats	—
Younis et al.	2015	<i>E. coli</i>	Egypt	Pet dogs and cats presenting at clinics for GI symptoms	—
Yousfi et al.	1977	Dermatophytes	Egypt	Buffalo, camel, cattle, cat, dog, donkey, goat, horse, sheep, pig	—
Yousfi et al.	2016	<i>E. coli</i>	Algeria	Healthy pet cats and dogs	—
Yousfi et al.	2016	<i>E. coli</i>	Algeria	—	—
Zim	1939	Helminths	Egypt	Cats and dogs	—
Zouari et al.	2017	<i>Bartonella</i> spp.	Tunisia	Fleas infesting dogs, cats, goats, and sheep	—

^aYear of publication.
GI, gastrointestinal.

canine vaccination campaigns (World Health Organization 2019). Without data on the present state of rabies infection in North African countries, it is difficult to argue for the inclusion of felines in prevailing canine vaccination and public awareness and education campaigns.

A small number of studies were done on viral infections other than rabies, limiting direct comparisons due to inconsistencies in study methods, target populations, and outcomes measured. The two studies concerning avian influenza found disparate seroprevalence in cats ranging from 8% to 56.4% (El Sayed et al. 2013, Ghoneim et al. 2013). Stoszek et al. (2006) concluded that contact with cats was a significant risk factor associated with anti-hepatitis E virus antibody production in pregnant study participants. The single study of rotavirus in the region conducted a phylogenetic analysis on the strain found in an infected infant, which suggests a direct inter-species transmission from cat to human (Fredj et al. 2013). Viral infections represent a substantial portion of the circulating and novel infectious diseases impacting human and animal health. Viruses have the capability to quickly evolve, appear in new forms and with new properties, infect multiple species, and travel to unexpected parts of the world (Mack et al. 2015). Analysis of published literature illustrates the important role of cats as viral reservoirs and exposure risks for vulnerable populations of children, the elderly, pregnant women, and the immunocompromised.

Bacterial infections

The vector-borne bacterial disease agents researched in the review include *Bartonella*, *Coxiella burnetii*, and *Rickettsia*. These pathogens can be transmitted by feline fleas and ticks. The two direct studies of vector-borne pathogens (VBP) in Egyptian cats produced similar results, 17% and 19% seropositivity for *Bartonella* spp. (Childs et al. 1995, Azzag et al. 2012). However, they conflicted with an Algerian study that found just 1.87% of sampled cats were seropositive for VBP, including *Rickettsia felis*, *B. henselae*, and *B. clarridgea* (Bessas et al. 2016). Two additional studies, in Tunisia and Morocco, tested feline fleas for the presence of VBP and found a seropositivity ranging from 33.3% for *Bartonella* spp. to 77.8% for at least one VBP including *Rickettsia* spp. and *Bartonella* spp. (Boudebouch et al. 2011, Zouari et al. 2017). A human study confirmed 14.5% of cases that were examined for CSD were in fact *Bartonella* spp. Although the authors repeatedly remarked on cat contact as a risk factor, they did not provide any data from their questionnaire to corroborate the association (Reda and Mostafa 1995). Another study of Algerian patients in a teaching hospital presenting with clinical signs of rickettsioses (spotted fevers) confirmed only 2 of the 277 potential cases were true positives and both reported frequent and recent contact with cats (Mouffok et al. 2008). Similarly, the principal risk factor for pregnant women infected with *Coxiella burnetii* was self-reported contact with domestic animals. Despite the fact that cat contact was implicated, their association was not specified in the study results (Kaabia and Letaief 2009).

In the studies of enteric bacterial infection rates in cats, *Campylobacter* spp. was found in a range from 3.3% to 10%, *Escherichia coli* from 36.8% to 79.2%, *Helicobacter pylori* from 28% to 33.3%, *Klebsiella* spp. at 6.67%, *Leptospira* spp. at 50%, *Mycobacterium avium* subsp. *paratuberculosis* at 42.8%, *Salmonella* from 2% to 12%, *Shigella* spp. at 4%,

Staphylococcus spp. from 6% to 21.3%, *Streptococcus* spp. from 2% to 36%, and *Pasteurella* spp. at 4% (Siam et al. 1973, Ghoneim et al. 1980, El Gohary 1998, Moustafa 1989, El-Jakee et al. 1991, Abdel-Moein and Samir 2011, Felt et al. 2011, Grami et al. 2013, Sallem et al. 2013, Abdel-Raouf et al. 2014, Gharsa et al. 2015, Sbai et al. 2015, Younis et al. 2015, Haggag et al. 2016, Yousfi et al. 2016, Elkassas et al. 2018, Salem et al. 2019). However, not all of the strains that were found in cats are pathogenic to humans, and the different methodologies among the studies make direct comparisons of prevalence impractical. *Brucella* spp. was isolated in a cat in a single study from Egypt, which the authors postulate may have an understated impact on the reemergence and transmission of brucellosis on farms (Wareth et al. 2017).

There were limited data available on the association between cat contact in North Africa and subsequent bacterial infections in humans, as either the risk factor was mentioned, but not quantified or the prevalence in humans was not determined. One study on *Helicobacter pylori* did conclude that persons with cat contact had a higher probability of *H. pylori* infection than persons with contact with dogs or other animals (Abdel-Raouf et al. 2014). In addition, a case study of a woman with a cat bite was positively identified to have *Pasteurella* spp. in her infected wound, establishing a direct source of transmission (Sbai et al. 2015).

A substantial number of studies researched the existence of antibiotic resistance genes in samples isolated from their study populations, including cats. Resistant strains of *E. coli* were isolated in 7.2–36.8% of cat sampled in the relevant studies (Grami et al. 2013, Sallem et al. 2013, Younis et al. 2015, Yousfi et al. 2016). Another study found *S. aureus* in just 6% of cats, but noted that all isolates from cats showed penicillin resistance (Gharsa et al. 2015).

Research conducted outside of the region has found that antimicrobial resistance (AMR) is increasing within the pathogens infecting companion animals, including cats (Lloyd 2007). As demonstrated by the number of studies on AMR genes in zoonotic bacterial pathogens found infecting cats and humans in North Africa, the risk of AMR is expanding in the domestic animal population, just as in the human population (Lloyd 2007). With such risk identified, measures need to be taken to effectively prevent and control not only bacterial diseases but also the emergence of resistant genes as well.

Fungal infections

In 1977, Yousef et al. found that 18.7% of Egyptian cats were positive for at least one pathogenic fungus, while 40 years later, in 2017, Waly et al. found that 16.7% of healthy cats were positive, suggesting remarkable stability in dermatophyte prevalence. A single study contained definitive prevalence rate numbers relating both human and feline dermatophyte infections, with animal contact a risk factor in 56.7% of positive human cases, but did not differentiate cats from other animals (Boumhil et al. 2010). Several studies commented on the rise of zoophilic species diagnosis in humans, and an increased representation in the isolates from skin lesions taken from dermatological patients (Benmezad et al. 2012, Aboueisha and El-Mahallawy 2013).

An additional study of note, but not included in the results, is Farag et al.'s (2018) article on dermatophyte infections and associated risk factors in school children from Menoufia,

Egypt. Pet contact was one of the leading risk factors for dermatophyte infections, which had a 1.41% overall prevalence in the study population. However, the authors did not specify what animals were included in the category of “pets” and thus the data cannot be stipulated to definitively include cat contact as a risk factor for dermatophyte infection in this population and was disqualified for inclusion in the formal review results (Farag et al. 2018).

Microsporidia, now considered fungi and not protozoa, have a transmission much more akin to enteric parasites than to dermatophyte fungal infections (Hibbett et al. 2007). Microsporidia were found in 23.1% of fecal samples from cats sampled from Egypt, with further analysis identifying them to be *Enterocytozoon bieneusi* and *E. intestinalis* species, demonstrating a common co-infection (Al-Herrawy and Gad 2016).

Parasitic infections

Helminths. Helminths are parasitic worms that are primarily intestinal, although some may reside in blood or other tissue. Taxonomically diverse, helminths encompass flatworms like trematodes or flukes, cestodes or tapeworms, and nematodes like roundworms and hookworms. Helminths re-

searched for this study are transmitted to humans primarily through the soil, either through accidental ingestion of contaminated soil or by walking barefoot. While mass deworming campaigns are not uncommon in North Africa, the routine use of veterinary dewormers in the region is unknown. According to World Health Organization (WHO) data from 2016, just 11.2% of school-aged children requiring preventive chemotherapy for soil-transmitted helminths actually received it in the Eastern Mediterranean region, the WHO region, which encompasses North Africa (World Health Organization 2016). The prevalence of certain helminths in the feline population has remained surprisingly consistent according to historical data for this region. For example, Zim (1939) found a 10% prevalence of *Taenia* spp. in cats in 1938, while a prevalence of 9.6% was recorded in 2017 by El-Dakhly et al. (2017). It is important to note that experimental methods varied widely, as did the locations of the surveys (Table 2).

Unicellular parasites. A survey of *Cryptosporidium* spp. was performed in humans with gastroenteritis and animals they were in contact with, to establish the incidence of cryptosporidiosis in households and animals associated with

TABLE 2. ZOONOTIC FELINE HELMINTH INFECTIONS IN NORTH AFRICA

Type	Species	Location	Year ^a	Prevalence in cats (%)	Author
Trematodes	<i>Heterophyes</i> spp.	Beni-Suef, Egypt	2017	3.2	El-Dakhly et al.
		Alexandria, Egypt	2017	11.76	El-Seify et al.
		Giza and Cairo, Egypt	1978	1.4	Arafa et al.
		Egypt	1939	90	Zim
		Egypt	1978	45.5	El-Shabrawy and Imam
	<i>Alaria</i> spp.	Beni-Suef, Egypt	2017	1.6	El-Dakhly et al.
		Alexandria, Egypt	2017	1.76	El-Seify et al.
	<i>Haplorchis</i>	Beni-Suef, Egypt	2017	1.6	El-Dakhly et al.
	<i>Pygidioopsis</i>	Beni-Suef, Egypt	2017	1.6	El-Dakhly et al.
	<i>Proceroverum</i>	Beni-Suef, Egypt	2017	1.6	El-Dakhly et al.
	<i>Prohemistomum</i>	Beni-Suef, Egypt	2017	1.6	El-Dakhly et al.
	<i>Echinochasmus</i>	Beni-Suef, Egypt	2017	6.45	El-Dakhly et al.
	<i>Euparadistomum</i>	Beni-Suef, Egypt	2017	4.8	El-Dakhly et al.
	<i>Asocotyle</i>	Giza and Cairo, Egypt	1978	3.2	Arafa et al.
		Beni-Suef, Egypt	2017	1.6	El-Dakhly et al.
Cestodes	<i>Dipylidium caninum</i>	Giza and Cairo, Egypt	1978	1.4	Arafa et al.
		Beni-Suef, Egypt	2017	62.9	El-Dakhly et al.
		Alexandria, Egypt	2017	18.8	El-Seify et al.
		Giza and Cairo, Egypt	1978	3.2	Arafa et al.
		Egypt	1939	50	Zim
	<i>Hymenolepis</i> spp.	Cairo, Egypt	1978	45.5	El-Shabrawy and Imam
		Alexandria, Egypt	2017	3.52	El-Seify et al.
		Beni-Suef, Egypt	2017	9.6	El-Dakhly et al.
		Alexandria, Egypt	2017	1.17	El-Seify et al.
		Giza and Cairo, Egypt	1978	9.2	Arafa et al.
	<i>Taenia taeniaeformis</i>	Egypt	1939	10	Zim
		Cairo, Egypt	1978	30.3	El-Shabrawy and Imam
		Beni-Suef, Egypt	2017	8–22.5	El-Dakhly et al.
		Giza and Cairo, Egypt	1978	2.7	Arafa et al.
		Cairo, Egypt	1978	24.2	El-Shabrawy and Imam
	<i>Diplopylidium</i> spp.	Beni-Suef, Egypt	2017	14.5	El-Dakhly et al.
		Giza and Cairo, Egypt	1978	0.9	Arafa et al.
		Cairo, Egypt	1978	31.8	El-Shabrawy and Imam
		Beni-Suef, Egypt	2017	33.8	El-Dakhly et al.
		Alexandria, Egypt	2017	8.23	El-Seify et al.
Nematodes	<i>Toxascaris leonina</i>	Alexandria, Egypt	2017	8.23	El-Dakhly et al.
		Egypt	1939	0	Zim
	<i>Toxocara cati</i>	Giza and Cairo, Egypt	1978	0.5	Arafa et al.
	<i>Gnathostomata</i> sp.	Giza and Cairo, Egypt	1978	0.5	Arafa et al.
		Giza and Cairo, Egypt	1978	0.5	Arafa et al.

^aYear of publication.

TABLE 3. STUDIES ON FELINE ZOONOTIC *TOXOPLASMA GONDII* IN NORTH AFRICA

Author	Location	Population	Cat contact as a risk factor for human infection
Ahmed et al.	Egypt	Pregnant women	Yes
Al-Kappany et al. ^a	Egypt	Feral cats	N/A
Bassiony et al.	Egypt	Pregnant women	No
Byomi et al.	Egypt	Women	Yes
El-Gamal et al.	Egypt	Pregnant women	Yes
El-Shqanqery et al.	Egypt	Pregnant women	No
Elmonir et al.	Egypt	Pregnant women; stray cats	No
Fakhfakh et al.	Tunisia	Pregnant women	No
Hafez Hassanain et al.	Egypt	Pregnant women	Yes
Ibrahim et al. (2006)	Egypt	Pregnant women	Yes
Ibrahim et al. (2017)	Egypt	Pregnant women	Yes
Khayeche et al.	Tunisia	Sheep	Yes
Messerer et al.	Algeria	Pregnant women	Yes
Rifaat et al. (1976)	Egypt	Stray cats	N/A
Rifaat et al. (1981)	Egypt	Stray cats	N/A
Romia et al.	Egypt	Humans	Yes

^aWhile two distinct studies were published in 2010 by Al-Kappany et al., only one study presented data with seropositivity rates in cats. N/A, not available.

confirmed human cases. Overall, 6.25% of fecal samples from cats were positive for *Cryptosporidium* spp. oocysts (El-Hohary and Abdel-Latif 1998). While this does not exclusively imply an etiologic role between species, the high detection of oocysts in the soil establishes the relevance of animal feces as a risk factor for transmission.

Entamoeba histolytica antibodies were found in 14 of 44 stool specimens taken from stray cats, but only one had a titer of 1/128 or greater to indicate a positive diagnosis at the time of sampling (Rifaat et al. 1981). While *Giardia* spp. was found in only 1% of stray cats in a study conducted in Egypt in 2011, the prevalence of *Giardia* spp. in fecal samples of stray cats was 11% in Egypt in 1999 (Khalifa 1999, Khalafalla 2011). Within Egypt, the prevalence of *Giardia* spp. in school-aged children and other stray animals, including dogs, cats, rats, and mice, was also determined, but the association between positive cases across species was not explored (Khalifa 1999). Nonetheless, the author did assert that dogs and cats should be considered reservoirs of infection, while rodents may play a role in the spread of parasites (Khalifa 1999).

Leishmaniasis is a disease caused by subspecies of *Leishmania* pathogenic eukaryotes and causes two forms of the disease presentation (cutaneous and visceral) in humans (Morsy and el Seoud Abou 1994). The Middle East and North Africa have long been established as a region endemic to leishmaniasis (Morsy et al. 1988). The bacteria are transmitted through the vector species of phlebotomine flies (the sand fly). The sand fly may also bite other mammals, such as rodents, dogs, and cats, and thus has the potential to spread disease from animals to humans, and vice versa (Morsy and el Seoud Abou 1994). Although the main confirmed reservoir host of *Leishmania* species in North Africa is dogs, contemporary research has suggested the role cats have as more than just an accidental host. Evidence of *Leishmania* infecting cats dates back to 1912 in Algeria (Maia et al. 2008). More recently, confirmed feline cases have occurred worldwide, including the Middle East, Mediterranean, Europe, and South America (Maia and Campino 2011).

Morsy conducted a study in 1982 that instead focused on ectoparasites on rodents in Egypt, identifying positive reactions for *Leishmania* antibodies in 16.7% of rodents (Morsy

et al. 1982). Researchers have reported human cases of cutaneous leishmaniasis since 1984, and visceral dating back to 1904 (Deperet and Boinet 1884, Nicolle and Cathoire 1905). Morsy published another study 10 years later focusing on the natural infection of *Leishmania* in domestic felines in Egypt (Morsy and el Seoud Abou 1994). In a survey of three regions, 2 out of 60 stray cats were found positive for *Leishmania* spp. The authors posit on whether these cats are accidental or reservoir hosts as they, “more or less roam the entire villages” and, unlike rats, are not “unaccepted and hated” (Morsy and el Seoud Abou 1994, p. 202).

The protozoan parasite of *Toxoplasma gondii* was first discovered in North Africa, so it is not a surprise that this is the pathogen with the most published studies in this review (Rouatbi et al. 2019). A disproportionate amount of *T. gondii* studies was located in Egypt (14 out of 17), although this trend resembles that of the studies overall (Table 3). One study, not included in the review, argued that ownership or contact with domestic felines is uncommon in Moroccan society and contact with cats is not significantly associated with *T. gondii* seropositivity (Ouzennou et al. 2019). Another study researched *T. gondii* in rabbits and found that the presence of cats in rabbit raising areas was significantly associated (odds ratio = 3.39) with *T. gondii* infection in rabbits (Abou Elez et al. 2017). Human contact with cats was not recorded in the study by Abou Elez et al. (2017) and therefore any attempt to look for an association between cat presence and human infection was inconclusive. Thus, this study was not included in the review results either. However, studies on human *T. gondii* infection rates in North African represent a prevalence range from 1% to 82% depending upon the country and study population (Rouatbi et al. 2019). And although at least seven wild felid species in this region have the potential to transmit *T. gondii*, no published data currently exist on their role in toxoplasmosis zoonoses (Rouatbi et al. 2019).

Regional trends and recommendations

While the majority of research on this subject has been conducted in Egypt, the assumption that this nation has the

majority of pathogens or rates of infection remains unproven. The few studies conducted in each country of this region that allows for comparison of a specific pathogen's prevalence indicate that it is more likely that the countries lacking robust research or public health veterinary surveillance have equal, if not greater, burdens compared with Egypt. In addition, this research cannot encompass the risks posed by undocumented transmission or transmission by sources apart from the domestic feline, which may add an entirely new layer to the surveillance, control, and approach of zoonoses in North Africa. Many of the feline-associated zoonotic pathogens outlined in this review can also be transmitted through contaminated food, water, or soil (*i.e.*, *Brucella* spp., *Giardia* spp., and *Toxocara* spp.), by vectors (*i.e.*, *B. henselae*), person to person (*i.e.*, *Cryptosporidium* spp.), and by other animal hosts (*i.e.*, *Leishmania* spp.).

Although the zoonotic diseases of felines in North Africa are varied and numerous, efforts to prevent and control zoonoses and reverse zoonoses in these countries are ongoing. The first line of defense is a network of veterinary public health professionals. Unfortunately, there is a disparity between access to veterinary care in North African countries, as well as between North Africa and North America. According to the World Organization of Animal Health, in 2018, Egypt had ~0.64 veterinary professionals per 1000 people, Morocco 0.074/1000, and Tunisia 0.176/1000. For comparison, the United States is around 0.55/1000. No data were available for Algeria or Libya (Veterinarians and Veterinary Para-Professionals 2018).

The most consistent control approach for zoonoses remains vaccination of target domestic species to prevent disease spread to humans or other species (Monath 2013). In addition to vaccination, prompt veterinary and medical care of zoonoses are essential in limiting the disease burden in North African communities. This includes programs aimed at reducing the number of feral and stray populations, which can harbor many zoonotic pathogens.

Cats managed through trap-neuter-return programs and left free roaming had higher incidence of helminth infections and lower incidence of viral and protozoan infections than cats in shelters, sanctuaries, and foster care (Andersen et al. 2018). While the study by Andersen et al. (2018) was conducted in the United States, this difference in enteric pathogens between free-roaming and confined cats could be present in North Africa as well, where the majority of felines live in the community without clear ownership. In regions where dimensions of cat ownership vary and routine veterinary care is improbable, the gap between the expected and actual roles of human and animal health care providers has allowed for a vacuum in zoonotic disease education and prevention. Historically, veterinarians have been seen as the source of information about zoonoses and their prevention for humans as well as their pets. Most physicians in a U.S. survey reported patient education on infectious disease risk should be the role of veterinarians (Grant and Olsen 1999). However, messages surrounding safe animal contact and steps to take to protect people and animals from zoonotic exposure risks should come from both human and animal health care using a collaborative voice.

While this review examined published studies of feline-human zoonotic disease transmission within the region of North Africa, narrative accounts of feline-associated zoono-

tic pathogens in singular populations of humans or animals without specific mention to transmission or transmission risk were not analyzed. Although literature not meeting the study criteria were omitted to prevent the unintentional interjection of false risk associations into the analysis, several publications were excluded that could provide information on the larger question surrounding the burden of zoonotic pathogens found within human and/or feline inhabitants of North Africa. Additional research on the prevalence of these zoonotic pathogens of public health importance within the region overall is warranted.

Conclusions

Zoonotic diseases shared between humans and felines have been, and continue to be, a public health threat that should be addressed with education and intervention strategies, which take into consideration the welfare of both species. This review has outlined several of the zoonotic feline pathogen risks that have been studied in the countries of Algeria, Egypt, Libya, Morocco, and Tunisia, as well as their most significant findings. Yet, despite the published data over the past several decades, many questions still exist on the role of felines in the spread of disease within North Africa. More research in this area is needed to create effective and comprehensive interventions for the protection of residents of this region and their feline counterparts.

Acknowledgments

The authors would like to recognize the University of North Florida Department of Public Health and the resource librarians for their support.

Author Disclosure Statement

No conflicting financial interests exist.

Funding Information

No funding received for this study.

References

- Abdel-Moein KA, Samir A. Isolation of enterotoxigenic *Staphylococcus aureus* from pet dogs and cats: A public health implication. *Vector Borne Zoonotic Dis* 2011; 11:627–629.
- Abdel-Raouf M, Abdel-Gleel Y, Enab A. Study on the role of pet animals for *Helicobacter pylori* transmission. *J Am Sci* 2014; 10:20–28.
- Abou Elez RMM, Hassanen EAA, Tolba HMN, Elsohaby I. Seroprevalence and risk factors associated with *Toxoplasma gondii* infection in domestic rabbits and humans. *Vet Parasitol Reg Stud Rep* 2017; 8:133–137.
- Aboueisha AM, El-Mahallawy H. Public health significance of dermatophytes in Ismailia and Port Said provinces, Egypt. *Med Mycol J* 2013; 54:123–129.
- Ahmed HA, Shafik SM, Ali ME, Elghamry ST, et al. Molecular detection of *Toxoplasma gondii* DNA in milk and risk factors analysis of seroprevalence in pregnant women at Sharkia, Egypt. *Vet World* 2014; 7:594–600.
- Ahmed MO, Elmeshri SE, Abuzweda AR, Blauo M, et al. Seroprevalence of brucellosis in animals and human populations in the western mountains region in Libya, December 2006–January 2008. *Euro Surveill* 2010; 15:19625.

- Al-Herrawy AZ, Gad MA. Microsporidial spores in fecal samples of some domesticated animals living in Giza, Egypt. *Iran J Parasitol* 2016; 11:195–201.
- Al-Kappany YM, Rajendran C, Abu-Elwafa SA, Hilali M, et al. Genetic diversity of *Toxoplasma gondii* isolates in Egyptian feral cats reveals new genotypes. *J Parasitol* 2010; 96:1112–1114.
- Al-Kappany YM, Rajendran C, Ferreira LR, Kwok OC, et al. High prevalence of toxoplasmosis in cats from Egypt: Isolation of viable *Toxoplasma gondii*, tissue distribution, and isolate designation. *J Parasitol* 2010; 96:1115–1119.
- Alkhateeb Shehada H. *Mamluks and Animals: Veterinary Medicine in Medieval Islam*. Leiden, Netherlands: Brill, 2012.
- American Veterinary Medical Association. AVMA pet ownership and demographics sourcebook, 2017–2018 edition. 2018. Available at <https://ebusiness.avma.org/productcatalog/product.aspx?ID=1531>
- Andersen LA, Levy JK, McManus CM, McGorray SP, et al. Prevalence of enteropathogens in cats with and without diarrhea in four different management models for unowned cats in the southeast United States. *Vet J* 2018; 236:49–55.
- Arafa MS, Nasr NT, Khalifa R, Mahdi AH, et al. Cats as reservoir hosts of *Toxocara* and other parasites potentially transmissible to man in Egypt. *Acta Parasitol Pol* 1978; 25: 383–389.
- Arrouji A, Lasfar ZB, Makni K. Rabies diagnostic laboratory. Institut Pasteur de Tunis. Report. *Arch Inst Pasteur Tunis* 1991; 68:89–95.
- Azzag N, Haddad N, Durand B, Petit E, et al. Population structure of *Bartonella henselae* in Algerian urban stray cats. *PLoS One* 2012; 7:e43621.
- Bakkali MM. Epidémiologie et Prophylaxie de la Rage au Maroc. In: Kuwert E, Mérieux C, Koprowski H, Bögel K, eds. *Rabies in the Tropics*. Berlin, Heidelberg: Springer Berlin Heidelberg, 1985;371–386.
- Bassiony H, Soliman NK, El Tawab S, Eissa S, Eossa A. Sero-prevalence and risk factors associated with *Toxoplasma gondii* infection among pregnant women in Alexandria, Egypt. *J Reprod Contracept Obstet Gynecol* 2016; 5:4220–4227.
- Benelmouffok A, Benhassine M, Abrous M. Rabies in Algeria. Recrudescence and new epizootic aspect. *Comp Immunol Microbiol Infect Dis* 1982; 5:321–326.
- Benhamiche B. *État actuel de la rage en Algérie: Deductions Prophylactiques (Current Rabies Situation in Algeria. Deductions Relating to Prophylaxis)*. Thèse Méd Vet Alfort 1977:35–48.
- Benmezad A, Moulahem T, Beneyezar M, Djaballah M, et al. Tinea capitis in the university hospital of Constantine (Algeria). *J Mycol Med* 2012; 22:354–356.
- Bessas A, Leulmi H, Bitam I, Zaidi S, et al. Molecular evidence of vector-borne pathogens in dogs and cats and their ectoparasites in Algiers, Algeria. *Comp Immunol Microbiol Infect Dis* 2016; 45:23–28.
- Boudebouch N, Sarih M, Beaucournu JC, Amarouch H, et al. *Henselae* and *Rickettsia felis* in fleas from Morocco. *Ann Trop Med Parasitol* 2011; 105:493–498.
- Boumhil L, Hjira N, Naoui H, Zerrouk A, et al. Tinea capitis in the military hospital Mohammed V (Morocco). *J Mycol Med* 2010; 20:97–100.
- Byomi A, Zidan S, Salama A, Elsify A, et al. Public health implications of toxoplasmosis in animals and women in selected localities of Meoufia Governorate, Egypt. *Assiut Vet Med J* 2018; 64:12–130.
- Childs JE, Olson JG, Wolf A, Cohen N, et al. Prevalence of antibodies to *Rochalimaea* species (cat-scratch disease agent) in cats. *Vet Rec* 1995; 136:519.
- Deperet C, Boinet E. The boil of Gafsa in the camp of Sathonay. *Méd Pharma Militaires* 1884; 3:296–302.
- Ducrottoy MJ, Azami HY, El Berbri I, Bouslikhane M, et al. Integrated health messaging for multiple neglected zoonoses: Approaches, challenges and opportunities in Morocco. *Acta Trop* 2015; 152:17–25.
- El Menyawe SM, Abdel Rahman MA. A study on *Toxocara* species and its public health importance. *Vet Med J Giz* 2007; 55:219.
- ElAshmawy WR, Ali ME. Identification of different dermatophytes isolated from cattle, cats and horses suffered from skin lesions. *Alex J Vet Sci* 2016; 49:126–132.
- El-Dakhly KM, Aboshinaf AM, El-Nahass ES, Gharib AE. A preliminary study on the Helminth fauna in necropsied stray cats (*Felis catus*) in Beni-Suef, Egypt. *J Adv Vet Res* 2017; 7:87–92.
- El-Gamal R, Selim M, Mohamed S, Fathy G, et al. Comparison of PCR with ELISA in diagnosis of recent toxoplasmosis in pregnant women. *J Am Sci* 2013; 9:824–832.
- El-Gohary AH. Prospective studies on campylobacteriosis in human and animals in contact. *Assiut Vet Med J* 1998; 38: 192–208.
- El-Hohary AH, Abdel-Latif AM. Zoonotic importance of cryptosporidiosis among some animals at Gharbia province in Egypt. *Indian J Anim Sci* 1998; 68:305–307.
- El-Jakee J. *Campylobacter jejuni* associated with diarrhea of pet animals and infants. *Vet Med J Giz* 1991; 39:81–89.
- Elkassas FA, Khaliel SA, Shawky M. Characterization of *Helicobacter pylori* and *Escherichia coli* from stool of human and pet animals. *Alex J Vet Sci* 2018; 56:95–101.
- Elmonir W, Harfoush MA, El-Tras WF, Kotb SA. Toxoplasmosis in stray cats and pregnant women in Egypt: Association between socio-demographic variables and high-risk practices by pregnant women. *Life Sci J* 2017; 14:1–5.
- El-Sayed A, Prince A, Fawzy A, Abdou MI, et al. Sero-prevalence of avian influenza in animals and human in Egypt. *Pak J Biol Sci* 2013; 16:524–529.
- El-Seify MA, Aggour MG, Sultan K, Marey NM. Gastro-intestinal helminths of stray cats in Alexandria, Egypt: A fecal examination survey study. *Vet Parasitol Reg Stud Rep* 2017; 8:104–106.
- El-Shabrawy MN, Imam EA. Studies on cestodes of domestic cats in Egypt with particular reference to species belonging to genera *Diplopylidium* and *Joyeuxiella*. *J Egypt Vet Med Assoc* 1978; 38:19–27.
- El-Shqanqery HE, Ibrahim HM, Mohamed AH, El-Sharaawy AA. Seroprevalence of *Toxoplasma gondii* infection and associated risk factors among asymptomatic pregnant females in Egypt. *J Egypt Soc Parasitol* 2017; 47:93–100.
- Fakhfakh N, Kallel K, Ennigro S, Kaouech E, et al. [Risk factors for *Toxoplasma gondii* and immune status of pregnant women: Cause and effect?] *Tunis Med* 2013; 91:188–190 (Article in French).
- Farag AG, Hammam MA, Ibrahim RA, Mahfouz RZ, et al. Epidemiology of dermatophyte infections among school children in Menoufia Governorate, Egypt. *Mycoses* 2018; 61:321–325.
- Faure E, Kitchener AC. An archaeological and historical review of the relationships between felids and people. *Anthrozoos* 2009; 22:221–238.
- Felt SA, Wasfy MO, Wael F, Samir A, et al. Cross-species surveillance of *Leptospira* in domestic and peri-domestic

- animals in Mahalla city, Gharbeya Governorate, Egypt. *Am J Trop Med Hyg* 2011; 84:420–425.
- Fredj MB, Heylen E, Zeller M, Fodha I, et al. Feline origin of *Rotavirus* strain, Tunisia, 2008. *Emerg Infect Dis* 2013; 19:630.
- Gautret P, Adehossi E, Soula G, Soavi MJ, et al. Rabies exposure in international travelers: Do we miss the target? *Int J Infect Dis* 2010; 14:e243–e246.
- Gerhold RW, Jessup DA. Zoonotic diseases associated with free-roaming cats. *Zoonoses Public Health* 2013; 60:189–195.
- Gharsa H, Slama KB, Gómez-Sanz E, Lozano C, et al. Molecular characterization of *Staphylococcus aureus* from nasal samples of healthy farm animals and pets in Tunisia. *Vector Borne Zoonotic Dis* 2015; 15:109–115.
- Ghoneim NH, Abdel-Moein KA, Zaher HM. The potential role of animals in the epidemiology of avian influenza virus H5N1 and its public health implications. *Glob Vet* 2013; 11: 609–613.
- Ghoneim NH, Hamed O, Abdelkarim AM. The role of stray cats in harbouring pathogens of zoonotic importance. *J Egypt Vet Med Assoc* 1980; 40:69–75.
- Grami R, Mansour W, Dahmen S, Mehri W, et al. The bla CTX-M-1 IncII/ST3 plasmid is dominant in chickens and pets in Tunisia. *J Antimicrob Chemother* 2013; 68:2950–2952.
- Grant S, Olsen CW. Preventing zoonotic diseases in immunocompromised persons: The role of physicians and veterinarians. *Emerg Infect Dis* 1999; 5:159–163.
- Haggag YN, Samaha HA, Nossair MA, Al Aswally SA. Epidemiological studies on *Helicobacter pylori* in some animals and humans. *Alex J Vet Sci* 2016; 51:275–281.
- Hassanain NA, Shaapan RM, Hassanain MA. Associated antenatal health risk factors with incidence of toxoplasmosis in Egyptian pregnant women. *Pak J Biol Sci* 2018; 21:463–468.
- Hasslinger MA, Omar HM, Selim MK. Incidence of helminths in stray cats in Egypt and other Mediterranean countries [1988]. *Vet Med Rev* 2013; 59:76–81.
- Helmy YA, El-Adawy H, Abdelwhab EM. A comprehensive review of common bacterial, parasitic and viral zoonoses at the human-animal interface in Egypt. *Pathogens* 2017; 6:33.
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, et al. A higher level phylogenetic classification of the Fungi. *Mycol Res* 2007; 111:509–547.
- Hotez PJ, Savioli L, Fenwick A. Neglected tropical diseases of the Middle East and North Africa: Review of their prevalence, distribution, and opportunities for control. *PLoS Negl Trop Dis* 2012; 6:e1475.
- Ibrahim HM, Mohamed AH, El-Sharaawy AA, El-Shqanqery HE. Molecular and serological prevalence of *Toxoplasma gondii* in pregnant women and sheep in Egypt. *Asian Pac J Trop Biomed* 2017; 10:996–1001.
- Ibrahim MA, Ghoneim NH, Ellatef GK. Investigation of toxoplasmosis in pregnant women and slaughtered animals. *Vet Med J Giz* 2006; 54:215.
- Kaabia N, Letaief A. Q fever in Tunisia. *Pathol Biol* 2009; 57: 439–443.
- Kaal JF, Baker K, Torgerson PR. Epidemiology of flea infestation of ruminants in Libya. *Vet Parasitol* 2006; 141:313–318.
- Khalafalla RE. A survey study on gastrointestinal parasites of stray cats in northern region of Nile delta, Egypt. *PLoS One* 2011; 6:e20283.
- Khalifa NO. Prevalence of *Giardia* cysts in pets, rodents, with possible implications in infantile diarrhea in Kaliobia Governorate. *Vet Med J Giz* 1999; 47:59–65.
- Khalil HM. Toxocariasis in Egypt. *J Egypt Public Health Assoc* 1977; 52:330.
- Khayeche M, Mhadhbi M, Gharbi M, Nasfi I, et al. Detection of *Toxoplasma gondii* infection of sheep slaughtered in the Governorate of Sousse on the occasion of the Muslim sacrifice feast (Eid Al-Adha) and analysis of risk factors. *Bull Soc Pathol Exot* 2014; 107:60–63.
- Kravetz JD, Federman DG. Cat-associated zoonoses. *Arch Gen Intern Med* 2002; 162:1945–1952.
- Kuntz RE, Chandler AC. Studies on Egyptian trematodes with special reference to the heterophyids of mammals. I. Adult flukes, with descriptions of *Phagicola longicollis* n. sp., *Cynodiplostomum Namrui* n. sp., and a *Stephanoprora* from cats. *J Parasitol* 1956; 42:445–459.
- Lloyd DH. Reservoirs of antimicrobial resistance in pet animals. *Clin Infect Dis* 2007; 45(Suppl_2):S148–S152.
- Mack A, Choffness ER. *Emerging Viral Diseases: The One Health Connection: Workshop Summary*. Washington, DC: National Academies Press, 2015.
- Macpherson CN. *Dogs, Zoonoses and Public Health*. London, UK: CABI, 2012.
- Maia C, Campino L. Can domestic cats be considered reservoir hosts of zoonotic leishmaniasis? *Trends Parasitol* 2011; 27: 341–344.
- Maia C, Nunes M, Campino L. Importance of cats in zoonotic leishmaniasis in Portugal. *Vector Borne Zoonotic Dis* 2008; 8:555–560.
- Maia C, Ramos C, Coimbra M, Bastos F, et al. Bacterial and protozoal agents of feline vector-borne diseases in domestic and stray cats from southern Portugal. *Parasit Vectors* 2014; 7:115.
- Majorowski MM, Carabin H, Kilani M, Bensalah A. Echinococcosis in Tunisia: A cost analysis. *Trans R Soc Trop Med Hyg* 2005; 99:268–278.
- Matter H, Blancou J, Benelmouffok A, Hammami S, et al. Historical perspective of rabies in Europe and the Mediterranean Basin. OIE, Paris (France), Geneva (Switzerland), 2004:185–199.
- Messerer L, Bouzbid S, Gourbdji E, Mansouri R, et al. [Seroprevalence of toxoplasmosis in pregnant women in Annaba, Algeria]. *Rev Epidemiol Sante Publique* 2014; 62:160–165 (Article in French).
- Moher D, Shamseer L, Clarke M, Gherzi D, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015; 4:1.
- Monath TP. Vaccines against diseases transmitted from animals to humans: A one health paradigm. *Vaccine* 2013; 31:5321–5338.
- Morsy TA, el Seoud Abou SM. Natural infection in two pet cats in a house of a zoonotic cutaneous leishmaniasis patient in Imbaba area, Giza Governorate, Egypt. *J Egypt Soc Parasitol* 1994; 24:199–204.
- Morsy TA, Michael SA, Bassili WR, Saleh MS. Studies on rodents and their zoonotic parasites, particularly leishmania, in Ismailiya Governorate, A.R. Egypt. *J Egypt Soc Parasitol* 1982; 12:565–585.
- Morsy TA, Michael SA, Makhoul LM. *Leishmania* infection sought in non human hosts in Suez Governorate, Egypt. *J Egypt Soc Parasitol* 1988; 18:539–545.
- Mouffok N, Parola P, Raoult D. Murine typhus, Algeria. *Emerg Infect Dis* 2008; 14:676.
- Mouffok N, Socolovschi C, Benabdellah A, Renvoise A, et al. Diagnosis of rickettsioses from eschar swab samples, Algeria. *Emerg Infect Dis* 2011; 17:1968.
- Moustafa S. Prevalence of *Campylobacter* and *Salmonella* in cats and dogs [Egypt]. *Assiut Vet Med J* 1989; 22:47–53.
- Nicoll C, Cathoire M. Note on a case of the boil of Gafsa. *Caducee* 1905; 10:1–2.

- Ouzennou N, Boussaa S, Ben Alla S, Boumezzough A. Observational study to assess pregnant women's knowledge and behaviour related to toxoplasmosis in Essaouira province, Morocco. *Asian Pac J Trop Med* 2019; 12:87–90.
- Reda WW, Mostafa O. Epidemiological studies on ocular cat-scratch-disease. *Vet Med J Giz* 1995; 43:383–388.
- Rifaat MA, Arafa MS, Sadek MS, Nasr NT, et al. *Toxoplasma* infection of stray cats in Egypt. *J Trop Med Hyg* 1976; 79: 67–70.
- Rifaat MA, Morsy TA, Sadek MS, Mahmoud AM. Antibodies against some parasites in stray cats in Cairo. *J Egypt Soc Parasitol* 1981; 11:517–524.
- Romia SA, Makhoul LM, el-Malky SA, Handoussa AE, et al. Seroepidemiologic observations on human toxoplasmosis in Dakahlia Governorate. *J Egypt Soc Parasitol* 1988; 18:493.
- Rouatbi M, Amairia S, Amdouni Y, Boussaadoun MA, et al. *Toxoplasma gondii* infection and toxoplasmosis in North Africa: A review. *Parasite* 2019; 26:6.
- Salem M, El-Deeb W, Abdel-Moein K, El-Sayed A, et al. Detection of *Mycobacterium avium* subsp. *paratuberculosis* in an Egyptian mixed breeding farm and comparative molecular characterisation of isolates from cattle, camels, and cats a case report. *Bulg J Vet Med* 2019; 22:41–49.
- Sallem RB, Gharsa H, Slama KB, Rojo-Bezares B, et al. First detection of CTX-M-1, CMY-2, and QnrB19 resistance mechanisms in fecal *Escherichia coli* isolates from healthy pets in Tunisia. *Vector Borne Zoonotic Dis* 2013; 13:98–102.
- Salyer SJ, Silver R, Simone K, Barton Behraves C. Prioritizing zoonoses for global health capacity building-themes from one health zoonotic disease workshops in 7 countries, 2014–2016. *Emerg Infect Dis* 2017; 23:S55–S64.
- Sbai MA, Dabloun S, Benzarti S, Khechimi M, et al. Acute carpal tunnel syndrome of the hand following a cat bite. *Pan Afr Med J* 2015; 21:206.
- Siam MA, Karim AA, Hamed OM, Zakaria A. The possible role of stray dogs and cats in the epidemiology of some bacterial human pathogens in Egypt. *Zentralbl Veterinarmed B* 1973; 20:409–414.
- Stoszek SK, Abdel-Hamid M, Saleh DA, Kafrawy SE, et al. High prevalence of hepatitis E antibodies in pregnant Egyptian women. *Trans R Soc Trop Med Hyg* 2006; 100:95–101.
- Torgerson PR, Macpherson CNL. The socioeconomic burden of parasitic zoonoses: Global trends. *Vet Parasitol* 2011; 182: 79–95.
- Tuzio H, Edwards D, Elston T, Jarboe L, et al. Feline zoonoses guidelines from the American association of feline practitioners. *J Feline Med Surg* 2005; 7:243–274.
- Vitale KR, Behnke AC, Udell MA. Attachment bonds between domestic cats and humans. *Curr Biol* 2019; 29:R864–R865.
- Veterinarians and Veterinary Para-Professionals. WAHIS Interface from World Organisation for Animal Health (OIE). 2018. Available at https://oie.int/wahis_2/public/wahid.php/Countryinformation/Veterinarians
- Waly N, Kamel R, Ahmed L, Ismail M. Prevalence of dermatophytosis in dogs and cats in Egypt. In BSAVA Congress Proceedings 2017, Birmingham, UK, BSAVA Library 2017: 457.
- Wareth G, Melzer F, El-Diasty M, Schmooch G, et al. Isolation of *Brucella abortus* from a dog and a cat confirms their biological role in re-emergence and dissemination of bovine brucellosis on dairy farms. *Transbound Emerg Dis* 2017; 64:e27–e30.
- World Bank. Middle East and North Africa. 2019. Available at <https://worldbank.org/en/region/mena>
- World Health Organization. Summary of global update on preventative chemotherapy implementation in 2015. *Weekly Epidemiol Rec* 2016; 91:456–460.
- World Health Organization. United against rabies collaboration, first annual progress report: Global strategic plan to end human deaths from dog-mediated rabies by 2030. WHO, FAO, and OIE, Geneva, Switzerland. 2019. Available at <https://apps.who.int/iris/rest/bitstreams/1251282/retrieve>
- Younis K, Baddour M, Ibrahim MS. Detection of diarrheagenic *Escherichia coli* in pet animals and its antibiotic resistance in Alexandria Governorate. *Alex J Vet Sci* 2015; 45.
- Yousef Y, Atia M, Esmail A. Dermatophytes in animals, birds and man. 1. Animals as a potential reservoir of dermatophytes to man. *Assiut Vet Med J* 1977; 4:151.
- Yousfi M, Mairi A, Touati A, Hassissene L, et al. Extended spectrum β -lactamase and plasmid mediated quinolone resistance in *Escherichia coli* fecal isolates from healthy companion animals in Algeria. *J Infect Chemother* 2016; 22:431–435.
- Yousfi M, Touati A, Mairi A, Brasme L, et al. Emergence of carbapenemase-producing *Escherichia coli* isolated from companion animals in Algeria. *Microb Drug Resist* 2016; 22:342–346.
- Zim A. Helminths of dogs and cats in Egypt. *Ann Parasitol Hum Comp* 1939; 17:32–36.
- Zouari S, Khrouf F, M'ghirbi Y, Bouattour A. First molecular detection and characterization of zoonotic *Bartonella* species in fleas infesting domestic animals in Tunisia. *Parasit Vectors* 2017; 10:436.

Address correspondence to:
 Amber N. Barnes
 Department of Public Health
 University of North Florida
 Building 39, Room 3011
 1 UNF Drive
 Jacksonville, FL 32224
 USA

E-mail: amber.barnes@unf.edu