

2017 Guidelines of the American Thyroid Association for the Diagnosis and Management of Thyroid Disease During Pregnancy and the Postpartum

Erik K. Alexander,^{1,*} Elizabeth N. Pearce,^{2,*} Gregory A. Brent,³ Rosalind S. Brown,⁴ Herbert Chen,⁵ Chrysoula Dosiou,⁶ William A. Grobman,⁷ Peter Laurberg,^{8,†} John H. Lazarus,⁹ Susan J. Mandel,¹⁰ Robin P. Peeters,¹¹ and Scott Sullivan¹²

Background: Thyroid disease in pregnancy is a common clinical problem. Since the guidelines for the management of these disorders by the American Thyroid Association (ATA) were first published in 2011, significant clinical and scientific advances have occurred in the field. The aim of these guidelines is to inform clinicians, patients, researchers, and health policy makers on published evidence relating to the diagnosis and management of thyroid disease in women during pregnancy, preconception, and the postpartum period.

Methods: The specific clinical questions addressed in these guidelines were based on prior versions of the guidelines, stakeholder input, and input of task force members. Task force panel members were educated on knowledge synthesis methods, including electronic database searching, review and selection of relevant citations, and critical appraisal of selected studies. Published English language articles were eligible for inclusion. The American College of Physicians Guideline Grading System was used for critical appraisal of evidence and grading strength of recommendations. The guideline task force had complete editorial independence from the ATA. Competing interests of guideline task force members were regularly updated, managed, and communicated to the ATA and task force members.

Results: The revised guidelines for the management of thyroid disease in pregnancy include recommendations regarding the interpretation of thyroid function tests in pregnancy, iodine nutrition, thyroid autoantibodies and pregnancy complications, thyroid considerations in infertile women, hypothyroidism in pregnancy, thyrotoxicosis in pregnancy, thyroid nodules and cancer in pregnant women, fetal and neonatal considerations, thyroid disease and lactation, screening for thyroid dysfunction in pregnancy, and directions for future research.

Conclusions: We have developed evidence-based recommendations to inform clinical decision-making in the management of thyroid disease in pregnant and postpartum women. While all care must be individualized, such recommendations provide, in our opinion, optimal care paradigms for patients with these disorders.

Keywords: pregnancy, thyroid and pregnancy, thyroid function tests, postpartum thyroiditis

¹Division of Endocrinology, Diabetes and Hypertension, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts.

²Section of Endocrinology, Diabetes, and Nutrition, Boston University School of Medicine, Boston, Massachusetts.

³Department of Medicine, VA Greater Los Angeles Healthcare System and David Geffen School of Medicine at UCLA, Los Angeles, California.

⁴Division of Endocrinology, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts.

⁵Department of Surgery, University of Alabama at Birmingham, Birmingham, Alabama.

⁶Division of Endocrinology, Stanford University School of Medicine, Stanford, California.

⁷Department of Obstetrics and Gynecology, Northwestern University, Chicago, Illinois.

⁸Departments of Endocrinology & Clinical Medicine, Aalborg University Hospital, Aalborg, Denmark.

⁹Institute of Molecular Medicine, Cardiff University, Cardiff, United Kingdom.

¹⁰Division of Endocrinology, Diabetes and Metabolism, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania.

¹¹Department of Internal Medicine and Rotterdam Thyroid Center, Erasmus Medical Center, Rotterdam, The Netherlands.

¹²Department of Obstetrics and Gynecology, Medical University of South Carolina, Charleston, South Carolina.

*Co-chairpersons: Erik K. Alexander and Elizabeth N. Pearce. Excepting the co-chairpersons, the authors are listed in alphabetical order and were appointed by the ATA to independently formulate the content of this manuscript. None of the scientific or medical content of the manuscript was dictated by the ATA.

†Deceased.

DEDICATION

These guidelines are dedicated to the memory of Dr. Peter Laurberg, our friend and colleague, who died tragically during their preparation.

I. INTRODUCTION

PREGNANCY HAS A PROFOUND IMPACT on the thyroid gland and its function. During pregnancy, the thyroid gland increases in size by 10% in iodine replete countries but by 20% to 40% in areas of iodine deficiency. Production of the thyroid hormones, thyroxine (T₄), and triiodothyronine (T₃), increases by nearly 50%, in conjunction with a separate 50% increase in the daily iodine requirement. These physiological changes happen seamlessly in healthy women, but thyroid dysfunction can occur in many pregnant women because of pathologic processes. Furthermore, other thyroid illnesses such as nodular disease and thyroid cancer are occasionally detected during pregnancy and may require treatment. Together, the burden of thyroid disease affecting women, either before, during, or directly after pregnancy, is substantial.

For these reasons thyroid function is frequently assessed during the gestation period. However, accurate assessment of maternal (and fetal) thyroid function during pregnancy remains difficult, and interpretation of laboratory testing differs from the nonpregnant patient. Placental human chorionic gonadotropin (hCG) stimulates thyroid hormone secretion, often decreasing maternal thyrotropin (TSH) concentrations, especially in early pregnancy. But while such transiently suppressed maternal TSH concentrations are often observed and deemed safe, defining the upper reference limit for serum TSH in this population has remained controversial. Furthermore, up to 18% of all pregnant women are thyroid peroxidase antibody (TPOAb) or thyroglobulin antibody (TgAb) positive. Increasingly, data suggest that TPOAb positivity adversely modulates the impact of maternal thyroid status (especially hypothyroidism) on the pregnancy and the developing fetus. Thyroid antibody positivity separately increases the risk of thyroid dysfunction following delivery and during the postpartum period.

Studies have recently questioned the optimal treatment of hyperthyroidism during pregnancy. Clinical management of patients with Graves' disease (GD) is challenged by the understanding that maternal antibodies as well as antithyroid medication may differentially affect maternal and fetal thyroid function. Reports have also detailed the potential teratogenic effects of the antithyroid medications methimazole (MMI) and propylthiouracil (PTU). But while mild hyperthyroidism appears safe for the mother and fetus, moderate to severe hyperthyroidism can prove dangerous. Thus, when and how to treat affected mothers during pregnancy remains an important clinical question. Following delivery, mothers often choose to breastfeed. Separate questions surround the optimal approach to the treatment of hypo- and hyperthyroidism while lactating.

Given the prevalence and potential dangers detailed above, many have suggested universally evaluating thyroid function in all women either before or during pregnancy. Such a screening mandate, however, must take the cost, effectiveness, and practical nature of any such approach into account. To date, studies evaluating this question appear to demonstrate

mixed conclusions. Several ongoing investigations will shed further light on this difficult question.

Given the complexity surrounding thyroid physiology and thyroid illness during pregnancy and the postpartum period, how and when to evaluate for thyroid dysfunction and how and if to treat thyroid illness during this period remain challenging.

In 2011, the American Thyroid Association (ATA) first published guidelines on the diagnosis and management of thyroid disease during pregnancy and postpartum (1). There has been a substantial amount of new literature in this area since that publication.

It is in this context that the ATA charged a task force to develop revised clinical guidelines on the diagnosis and treatment of thyroid disease during pregnancy and the postpartum period. The task force consisted of both national and international experts in the field of thyroid disease and pregnancy and included representatives from the ATA, the European Thyroid Association (ETA), the American College of Obstetricians and Gynecologists, the Society for Maternal Fetal Medicine, and the Iodine Global Network. In addition to evidence-based updates of traditional content areas, the task force also sought to expand the prior document to address topics such as thyroid disease during lactation, the treatment of thyroid illness in infertile women and those undergoing assisted reproductive techniques (ART), as well as the approach to thyroid disease in the newborn.

The literature review for each section included an analysis of all primary studies in the area published since 1990 and selective review of the primary literature published prior to 1990 that was seminal in the field. In the past 25 years, there have been a number of recommendations and guideline statements relating to aspects of thyroid disease and pregnancy. In deriving the present guidelines, the task force conducted a new and comprehensive analysis of the primary literature and reformulated all clinical recommendations. In doing so, this document represents the best effort to create a useful, practical, and accurate guideline designed to help the practicing clinician, while also stimulating future research and discovery into this important and complex arena.

II. METHODS

ATA Thyroid Disease in Pregnancy guidelines were previously published in 2011 (1). Because of the rapid growth of the literature relating to this topic, plans for revising the guidelines within ~4–5 years of publication were made at the inception of the project. Task force chairs were appointed by the ATA President with approval of the Board. A task force of specialists with complementary expertise (adult and pediatric endocrinology, obstetrics, maternal-fetal medicine, endocrine surgery, iodine nutrition, and epidemiology) was appointed. In order to have broad specialty and geographic representation, as well as fresh perspectives, approximately one third of the task force is to be replaced for each iteration of the guidelines, as per ATA policy. In accordance with current ATA policies, the American College of Physicians Grading System was adopted for use in these guidelines (Tables 1 and 2) (2).

Prior to initiating the reviews, all task force members were provided written and verbal group advice on conducting electronic literature searches, critical appraisal of articles,

TABLE 1. RECOMMENDATIONS (FOR THERAPEUTIC INTERVENTIONS) BASED ON STRENGTH OF EVIDENCE

<i>Recommendation and evidence quality</i>	<i>Description of supporting evidence^a</i>	<i>Interpretation</i>
Strong recommendation		
High-quality evidence	RCT without important limitations or overwhelming evidence from observational studies	Can apply to most patients in most circumstances without reservation
Moderate-quality evidence	RCT with important limitations or strong evidence from observational studies	Can apply to most patients in most circumstances without reservation
Low-quality evidence	Observational studies/case studies	May change when higher-quality evidence becomes available
Weak recommendation		
High-quality evidence	RCT without important limitations or overwhelming evidence from observational studies	Best action may differ based on circumstances or patients' values
Moderate-quality evidence	RCT with important limitations or strong evidence from observational studies	Best action may differ based on circumstances or patients' values
Low-quality evidence	Observational studies/case studies	Other alternatives may be equally reasonable
Insufficient	Evidence is conflicting, of poor quality, or lacking	Insufficient evidence to recommend for or against

^aThis description of supporting evidence refers to therapy, therapeutic strategy, or prevention studies. The description of supporting evidence is different for diagnostic accuracy studies.

RCT, randomized controlled trial.

TABLE 2. RECOMMENDATIONS (FOR DIAGNOSTIC INTERVENTIONS) BASED ON STRENGTH OF EVIDENCE

<i>Recommendation and evidence quality</i>	<i>Methodologic quality of supporting evidence</i>	<i>Interpretation</i>
Strong recommendation		
High-quality evidence	Evidence from one or more well-designed nonrandomized diagnostic accuracy studies (i.e., observational—cross-sectional or cohort) or systematic reviews/meta-analyses of such observational studies (with no concern about internal validity or external generalizability of the results).	Implies the test can be offered to most patients in most applicable circumstances without reservation.
Moderate-quality evidence	Evidence from nonrandomized diagnostic accuracy studies (cross-sectional or cohort), with one or more possible limitations causing minor concern about internal validity or external generalizability of the results.	Implies the test can be offered to most patients in most applicable circumstances without reservation.
Low-quality evidence	Evidence from nonrandomized diagnostic accuracy studies with one or more important limitations causing serious concern about internal validity or external generalizability of the results.	Implies the test can be offered to most patients in most applicable circumstances, but the utilization of the test may change when higher-quality evidence becomes available.
Weak recommendation		
High-quality evidence	Evidence from one or more well-designed nonrandomized diagnostic accuracy studies (i.e., observational—cross-sectional or cohort) or systematic reviews/meta-analyses of such observational studies (with no concern about internal validity or external generalizability of the results).	The degree to which the diagnostic test is seriously considered may differ depending on circumstances or patients' or societal values.
Moderate-quality evidence	Evidence from nonrandomized diagnostic accuracy studies (cross-sectional or cohort), with one or more possible limitations causing minor concern about internal validity or external generalizability of the results.	The degree to which the diagnostic test is seriously considered may differ depending on individual patients' practice circumstances or patients' or societal values.
Low-quality evidence	Evidence from nonrandomized diagnostic accuracy studies with one or more important limitations causing serious concern about internal validity or external generalizability of the results.	Alternative options may be equally reasonable.
Insufficient	Evidence may be of such poor quality, conflicting, lacking (i.e., studies not done), or not externally generalizable to the target clinical population such that the estimate of the true effect of the test is uncertain and does not permit a reasonable conclusion to be made.	Insufficient evidence exists to recommend for or against routinely offering the diagnostic test.

and rationale for formulating strength of recommendations. Standardized data collection forms were used by all reviewers. For each question, a primary reviewer performed a literature search, appraised relevant literature, and generated recommendations, accompanying text, and a relevant bibliography. This information was then reviewed by both chairs, revised as needed, and presented for review by the entire panel. Feedback and suggestions for revisions from the Chairs and panel members were obtained via e-mail, regularly scheduled teleconferences, and face-to-face meetings. Once the manuscript was drafted, all suggestions for revisions were regularly reviewed by the entire panel in the form of a tracked changes draft manuscript and teleconferences. The draft document continued to be revised until no suggestions for further revisions were requested by any panel members. Thus, general consensus on acceptability of recommendations and manuscript text was achieved, with the fundamental understanding that not all recommendations may be feasible in all practice settings and adaptation of the guideline recommendations to individual care may be needed.

Formal stakeholder input in development of these guidelines was sought from ATA membership via an online survey distributed in October 2014. We also reviewed any letters, editorials, or reviews of the 2011 iteration of these guidelines (1) that were collected by the current chairs of the task force. Prepublication verbal feedback on some of the key guideline recommendations was received at a formal Satellite Symposium held in conjunction with the Endocrine Society meeting in Boston on March 31, 2016. The guidelines were then provided to the ATA membership for review and comments over a 2-week period. The guideline manuscript was next reviewed and approved by the ATA Board of Directors. Feedback and suggestions were formally discussed by the panel, and revisions were made to the manuscript prior to journal submission. The organization of management guideline recommendations is shown in Table 3.

TABLE 3. ORGANIZATION OF THE TASK FORCE'S RECOMMENDATIONS

<i>Location key</i>	<i>Description</i>	<i>Page</i>
I	Introduction	316
II	Methods	316
III	Thyroid Function Testing and Pregnancy	318
IV	Iodine Status and Nutrition	321
V	Thyroid Autoantibodies and Pregnancy Complications	324
VI	The Impact of Thyroid Illness upon Infertility and Assisted Reproduction	328
VII	Hypothyroidism and Pregnancy	331
VIII	Thyrotoxicosis in Pregnancy	343
IX	Thyroid Nodules and Thyroid Cancer During Pregnancy	350
X	Fetal and Neonatal Considerations	356
XI	Thyroid Disease and Lactation	359
XII	Postpartum Thyroiditis	361
XIII	Screening for Thyroid Dysfunction Before or During Pregnancy	363
XIV	Future Research Directions	367

The medical opinions expressed here are those of the authors, and the task force had complete editorial independence from the ATA in writing the guidelines. No funding was received by individual task force members from the ATA or industry for work on these guidelines. Competing interests of all task force members were reviewed at inception of the group, yearly, and upon completion of the guidelines and are included with this document.

These Guidelines were approved by the Board of the ATA.

III. THYROID FUNCTION TESTING AND PREGNANCY

Question 1: How do thyroid function tests change during pregnancy?

Normal pregnancy is associated with an increase in renal iodine excretion, an increase in thyroxine binding proteins, an increase in thyroid hormone production, and thyroid stimulatory effects of hCG. All of these factors influence thyroid function tests in the pregnant patient. The healthy thyroid adapts to these alterations through changes in thyroid hormone metabolism, iodine uptake, and the regulation of the hypothalamic-pituitary-thyroid axis (3,4). The thyroid function tests of healthy pregnant women, therefore, differ from those of healthy nonpregnant women. Furthermore, the reference ranges for the most widely applied tests, TSH and free thyroxine (FT4), may vary significantly in different populations.

Following conception, circulating thyroxine binding globulin (TBG) and total T4 (TT4) concentrations increase by week 7 of gestation and reach a peak by approximately week 16 of gestation (5). These concentrations then remain high until delivery. In the first trimester, maternal hCG directly stimulates the TSH receptor, increasing thyroid hormone production and resulting in a subsequent reduction in serum TSH concentration (4,6). Therefore, during pregnancy, women have lower serum TSH concentrations than before pregnancy, and a TSH below the nonpregnant lower limit of 0.4 mU/L is observed in as many as 15% of healthy women during the first trimester of pregnancy (7,8). In Japan, a suppressed TSH less than 0.6 mU/L is similarly frequently observed during the first trimester of pregnancy (9). The fraction of women with a suppressed TSH falls to about 10% in the second trimester, and 5% in the third trimester (4).

Measurement of FT4 concentration by automated immunoassays results in a significant and assay dependent reduction in the measured serum FT4 concentrations in the third trimester, even though direct measurement of free T4 by more precise methods does not show a similar degree of reduction (4,6,7). Automated immunoassays for FT4, which are employed in most clinical laboratories, are complicated in pregnant women by the increase in TBG and decrease in albumin concentrations (10,11). Other methods of direct measurement, such as measurement by equilibrium dialysis, ultrafiltration, or liquid chromatography/tandem mass spectrometry (LC/MS/MS) are less influenced by the pregnancy-associated changes in serum proteins but are significantly more expensive and less widely available. Thus, the automated immunoassays used for serum FT4 analysis are still widely used, but the important considerations discussed above must be noted. The use of population-based, trimester-specific reference ranges remains the best way to handle this issue.

Question 2: What is the normal reference range for serum TSH concentrations in each trimester of pregnancy?

A downward shift of the TSH reference range occurs during pregnancy, with a reduction in both the lower (decreased by about 0.1–0.2 mU/L) and the upper limit of maternal TSH (decreased by about 0.5–1.0 mU/L), relative to the typical nonpregnant TSH reference range. The largest decrease in serum TSH is observed during the first trimester because of elevated levels of serum hCG directly stimulating the TSH receptor and thereby increasing thyroid hormone production (Table 4). Thereafter, serum TSH and its reference range gradually rise in the second and third trimesters, but nonetheless remain lower than in nonpregnant women (12,13). Since hCG concentrations are higher in multiple pregnancies than in singleton pregnancies, the downward shift in the TSH reference interval is greater in twin pregnancies (11). In a study of 63 women with hCG concentrations >200,000 IU/L, TSH was suppressed (≤ 0.2 mU/L) in 67% of women and in 100% of women if hCG concentrations were >400,000 IU/L (14). Serum TSH reference range determinations should take into account iodine intake, TPO positivity, and according to some studies, body mass index (BMI).

Although the downward shift in TSH reference ranges is seen in essentially all populations, the extent of this reduction varies significantly between different racial and ethnic groups. Initial studies of pregnant women in the United States and Europe first led to recommendations for a TSH upper reference limit of 2.5 mU/L in the first trimester and 3.0 mU/L in the second and third trimesters (1,15). However, more recent studies in pregnant women in Asia, India, and the Netherlands, have demonstrated only a modest reduction in the upper reference limit (16–20). A study of 4800 pregnant women in China recently showed that the downward shift in the TSH reference range occurred at weeks 7–12, but the upper reference limit was only reduced from 5.31 to 4.34 mU/L (17). Separate data from a recent prospective intervention trial in the United States support this finding (21). Analysis of the TSH and FT4 “set-point” in pregnant women showed that reductions in FT4 were observed only when the serum TSH was greater than 4.8 mU/L. Similar studies of pregnant women in India and Korea show a modest reduction in the first-trimester upper TSH limit of 0.5–1.0 mU/L. In some cases, this was not statistically different from the nonpregnant state (18,20). Thus, the current evidence supports only a slight downward shift in the upper reference range of TSH occurring in the latter first trimester of pregnancy, typically not seen prior to week 7 (17).

A reduction in the lower TSH reference range is observed during pregnancy in almost all studies. In a small percentage of women, TSH can be undetectable (<0.01 mU/L), and yet still represent a normal pregnancy. In addressing the clinical importance of a reduced serum TSH during pregnancy, it is important to note that subclinical hyperthyroidism has not been associated with adverse pregnancy outcomes. Therefore, a maternal TSH concentration that is low but detectable is likely not clinically significant (22). TSH ranges have been shown to vary slightly depending on different methods of analysis, although this variation is not clinically significant (23). One approach to reducing this variability is to use the Multiple of Medians calculation to compare values between

assays. This calculation divides an individual value by the population median (24). The resulting value is not influenced by the differences between assays (24).

Significant geographic and ethnic diversity exist in TSH concentrations during pregnancy, as shown in Table 4. The task force recognizes the limited availability of trimester-specific reference ranges calculated for most ethnic and racial populations with adequate iodine intake who are free of thyroid autoantibodies. Nonetheless, to provide guidance to all patients and clinicians, the panel recommends use of the following trimester-specific ranges and cutoffs when local assessments are not available. In the first trimester, the lower reference range of TSH can be reduced by approximately 0.4 mU/L, while the upper reference range is reduced by approximately 0.5 mU/L. For the typical patient in early pregnancy, this corresponds to a TSH upper reference limit of 4.0 mU/L. This reference limit should generally be applied beginning with the late first trimester, weeks 7–12, with a gradual return towards the nonpregnant range in the second and third trimesters. For specific recommendations regarding the diagnosis and treatment of maternal hypothyroidism, see Section VII.

RECOMMENDATION 1

When possible, population-based trimester-specific reference ranges for serum TSH should be defined through assessment of local population data representative of a health care provider’s practice. Reference range determinations should only include pregnant women with no known thyroid disease, optimal iodine intake, and negative TPOAb status.

Strong recommendation, moderate-quality evidence.

Question 3: What is the optimal method to assess serum T4 concentration during pregnancy?

Unbound T4 represents only about 0.03% of serum TT4 content. Importantly, only FT4 is available for tissue uptake, with the remainder of T4 bound to serum proteins, primarily TBG. Serum TT4 concentrations are measured in the nanomolar range, while FT4 concentrations are measured in the picomolar range. In part because of this, measuring FT4 in the presence of high concentrations of bound T4 has proved to be challenging. This challenge is especially true in conditions where binding-protein levels are altered, such as pregnancy.

Equilibrium dialysis and ultrafiltration are used for physical separation of serum FT4 from bound T4 prior to analysis of the dialysate or ultrafiltrate. While they are theoretically not influenced by changes in binding proteins and heterophilic antibodies, assays based on classical equilibrium dialysis or ultrafiltration are laborious, time-consuming, expensive, and not widely available.

As noted above, FT4 measurement performed by indirect analog immunoassays is used by the majority of clinical laboratories, largely because of its ability to be quickly performed on automated platforms. Unfortunately, this approach is prone to inaccuracy in the setting of pregnancy because of disruption of the original equilibrium—a process dependent upon dilution, temperature, buffer composition, affinity, and the concentration of the T4 antibody reagent and the T4-binding capacity within the serum sample (25). High protein concentrations in serum samples tend to result in higher FT4

TABLE 4. REFERENCE RANGES FOR THYROTROPIN AND FREE THYROXINE DURING EARLY PREGNANCY WORLDWIDE

Author, country (reference) (analyzing method)	N	Gestation (week)	TSH, mU/L		FT4, pmol/L (ng/dL)		Population characteristics			
			Median	2.5th– 97.5th	Median	2.5th– 97.5th	(Median, 2.5th–97.5th)	Iodine insufficiency	Mean BMI	Ethnicities
Bestwick <i>et al.</i> , Italy (24) (AutoDELFIA)	5505	<16	1.07	0.04–3.19	9.3	7.4–12.2	(0.73, 0.58–0.95)	Moderate-mild ^a	NR	NR
Bestwick <i>et al.</i> , UK (24) (Advia Centaur)	16,334	<16	1.11	0.06–3.50	13.9	10.9–17.9	(1.08, 0.85–1.40)	Moderate-mild ^a	NR	NR
Bocos-Terraz <i>et al.</i> , Spain (264) (Architect)	481	<14	0.94	0.41–2.63	13.9	10.8–17.8	(1.08, 0.84–1.38)	Mild	NR	Caucasian (93%)
Gilbert <i>et al.</i> , Australia (271) ^b (Architect)	1817	9–13	0.74	0.02–2.15	13.5	10.4–17.8	(1.05, 0.81–1.39)	Borderline	NR	Australian
Lambert-Messerlian <i>et al.</i> , USA (270) ^c (Immolute 2000)	8351	T1	1.00	0.12–3.37	14.2	10.4–17.8	(1.10, 0.81–1.38)	Mild	NR	Caucasian (67%) and Hispanic (23%) ^d
La'ulu <i>et al.</i> , USA (139,265) ^e	8415	T2	1.19	0.35–3.35	13.0	9.3–16.2	(1.01, 0.72–1.26)		NR	Hispanic (37%), Caucasian (29%), African American (27%), Asian (8%)
	2172	10–13	0.94	0.02–2.69	14.7	11.4–18.6	(1.15, 0.89–1.45)	Mild	NR	
	2683	14–20	1.14	0.15–3.11	12.0	9.3–15.2	(0.94, 0.73–1.19)		NR	
Li <i>et al.</i> , China (17) (Cobas Elesys 601)	640	7–12	1.47	0.10–4.34	15.8	12.3–20.9	(1.23, 0.96–1.63)	Proven sufficient ^f	NR	Chinese (presumed)
Männistö <i>et al.</i> , Finland (266) (Architect i2000)	4333	T1	1.11	0.08–3.54	15.3	11.7–22.8	(1.12, 0.86–1.58)	Sufficient	22.4	Finnish (presumed)
	747	T2	1.37	0.11–4.24	14.6	11.2–23.4	(1.13, 0.87–1.82)			
Medici <i>et al.</i> , the Netherlands (267) (Vitros ECI)	5186	8–18	1.30	0.03–4.04	14.7	10.4–22.0	(1.15, 0.81–1.72)	Proven sufficient ^f	24.5	Dutch (52%), Surinamese/Antillean (12%), Turkish (8%), Moroccan (6%)
Pearce <i>et al.</i> , USA (142) (Advia Centaur)	585	<14	1.1	0.04–3.60	2.1 ^h	1.5–2.9 ^g	—	Borderline	NR	White (77%) and African American (10%)
Quinn <i>et al.</i> , Russia (272) (Abbott AxSYM)	380	T1	1.66	0.09–4.67	—	—	—	Moderate	NR	Russian (presumed)
	549	T2	2.00	0.20–4.68	—	—	—		NR	
Springer <i>et al.</i> , Czech Republic (268) ^h (ADVIA Centaur)	4337	9–11	1.21	0.06–3.67	—	—	—	Mild	NR	Caucasian (99%)
Stricker <i>et al.</i> , Switzerland (262) (Architect i2000SR)	575	6–12	0.95	0.07–2.82	13.9	10.5–18.5	(1.08, 0.82–1.44)	Sufficient	NR	Swiss (presumed)
	528	T2	1.02	0.20–2.79	12.2	9.5–15.7	(0.95, 0.74–1.22)			
Vaidya <i>et al.</i> , UK (Modular E 170) (274)	1089	<12	1.08	0.14–3.19	14.6	10.7–19.4	(1.12, 0.83–1.59)	Mild-moderate	NR	Caucasian (91) and South Asian (4)

Studies were selected according to the following criteria: $N \geq 500$, exclusion of thyroid peroxidase antibody (TPOAb)-positive women and availability of data from the manuscript or via personal communication. Iodine status was estimated based on references from article, WHO iodine status reports or from the Vitamin and Mineral Nutrition Information System (VMINIS).

^aWeight reported (Bestwick *et al.* median weight 59 kg in Italian and 67 kg in UK population).

^bReported FT4 level is a mean.

^cLimits are 5th and 98th percentiles for TSH and 2nd and 95th percentiles for FT4.

^dBased on reports of the total FASTER population.

^eFT4 determined in normal-range TSH only.

^fBased on iodine measurements in study population.

^gFree T4 index (reference range 1.0–4.0).

^hHigh hCG levels excluded.

TSH, thyrotropin; FT4, free thyroxine; NR, not reported; T1, first trimester; T2, second trimester.

values, whereas low protein concentrations are likely to yield lower FT4 values. In order to decrease nonspecific binding and neutralize the effect of nonesterified fatty acids on serum FT4, albumin is added in some assays. Albumin itself, however, binds T4 and when added in sufficient amounts, may disrupt the equilibrium. Nevertheless, the currently used FT4 immunoassays perform reasonably well in many circumstances, and most often accurately report both low FT4 levels in the setting of thyroid hormone deficiency and high FT4 levels in the setting of thyroid hormone excess (26,27).

Sera of pregnant women are characterized by higher concentrations of TBG and nonesterified fatty acids and by lower concentrations of albumin relative to the sera of nonpregnant women. In part because of this, many current FT4 analog immunoassays fail dilutional assessment (25,28). Because FT4 reference intervals in pregnancy vary widely between methods, interpretation of FT4 values requires method-specific as well as trimester-specific ranges (10,11,27). Whereas it is customary for manufacturers to suggest that laboratories establish their own reference range for such a test, this suggestion is frequently impractical for FT4 assessment because it is especially difficult to recruit subjects with specific conditions such as pregnancy from which to independently establish method- and trimester-specific reference ranges. Therefore, laboratories customarily adopt the pregnancy ranges provided by the test manufacturers. Typically, the characteristics of these reference pregnant cohorts are not disclosed, and differences in iodine intake and ethnicity may compromise the ability to generalize the manufacturer ranges across different populations. This problem adds to the complexity of accurate measurement of serum FT4 in the pregnant individual.

Current uncertainty around FT4 estimates in pregnancy has led some to question the wisdom of relying on any FT4 immunoassays during pregnancy (29,30). In contrast, measurement of TT4 and the calculated FT4 index do show the expected inverse relationship with serum TSH (29). This finding suggests that TT4 measurements may be superior to immunoassay measurement of FT4 measurements in pregnant women. However, reference values should take the 50% increase in TBG witnessed during pregnancy into account by calculating the FT4 index using a serum thyroid hormone uptake test (such as the thyroid hormone binding ratio). Changes in total serum T4 concentration through pregnancy among euthyroid women have been previously reported (5). Changes are predictable, with an increase in TT4 concentration from weeks 7–16 of gestation, ultimately reaching ~50% above the prepregnancy level. This level is then sustained through pregnancy. Therefore, a clinically acceptable upper range determination can be calculated by shifting the nonpregnant limit 50% higher. However, this limit can only be used after week 16 of pregnancy. If a T4 measurement is required before that time (i.e., weeks 7–16 of pregnancy), a calculation can be made for the upper reference range based on increasing the nonpregnant upper reference limit by 5% per week, beginning with week 7. For example, at 11 weeks of gestation (4 weeks beyond week 7), the upper reference range for T4 is increased by 20% (4 weeks \times 5%/week) (5).

As described above, free thyroid hormones can also be measured in the dialysate or ultrafiltrate using online solid phase extraction (LC/MS/MS). However, this approach is time con-

suming, costly, and often impractical. Using direct equilibrium dialysis and LC/MS/MS, the 95% FT4 reference intervals decrease gradually with advancing gestational age: from 1.08–1.82 ng/dL in week 14 to 0.86–1.53 ng/dL in week 20 (31). Serum FT4 by LC/MS/MS correlates very well with serum FT4 measured by classical equilibrium dialysis, but correlation with results from the FT4 immunoassay are less satisfactory (8). The use of isotope dilution-LC/MS/MS for measuring T4 in the dialysate from equilibrium dialysis of serum is helpful to obtain a gold-standard reference measurement procedure for serum FT4 (32). This assay technology, unfortunately, is currently not widely available due to high instrument and operating costs.

■ RECOMMENDATION 2

The accuracy of serum FT4 measurement by the indirect analog immunoassays is influenced by pregnancy and also varies significantly by manufacturer. If measured in pregnant women, assay method-specific and trimester-specific pregnancy reference ranges should be applied.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 3

In lieu of measuring FT4, TT4 measurement (with a pregnancy-adjusted reference range) is a highly reliable means of estimating hormone concentration during the last part of pregnancy. Accurate estimation of the FT4 concentrations can also be done by calculating a FT4 index.

Strong recommendation, moderate-quality evidence.

IV. IODINE STATUS AND NUTRITION

Because of increased thyroid hormone production, increased renal iodine excretion, and fetal iodine requirements, dietary iodine requirements are higher in pregnancy than they are for nonpregnant adults (33). Women with adequate iodine intake before and during pregnancy have adequate intrathyroidal iodine stores and have no difficulty adapting to the increased demand for thyroid hormone during gestation. In these women, total-body iodine levels remain stable throughout pregnancy (34). However, in areas of even mild to moderate iodine deficiency, total-body iodine stores, as reflected by urinary iodine values, decline gradually from the first to the third trimester of pregnancy (35). Iodine, required for infant nutrition, is secreted into breast milk. Therefore, lactating women also have increased dietary iodine requirements (36).

Spot urinary iodine values are used most frequently for determination of iodine status in populations. Because there is substantial diurnal and day-to-day variation in urinary iodine excretion, urinary iodine concentrations (UICs) cannot be used to identify particular individuals with iodine deficiency (37,38). Therefore, iodine levels are a population rather than individual marker and outside unusual settings urinary iodide testing is not beneficial for individual use.

■ RECOMMENDATION 4

Median UICs can be used to assess the iodine status of populations, but single spot or 24-hour UICs are not a valid marker for the iodine nutritional status of individual patients.

Strong recommendation, high-quality evidence.

Question 4: What is the impact of severe iodine deficiency on the mother, fetus, and child?

Maternal dietary iodine deficiency results in impaired maternal and fetal thyroid hormone synthesis. Low thyroid hormone values stimulate increased pituitary TSH production, and the increased TSH stimulates thyroid growth, resulting in maternal and fetal goiter (39). In areas of severe iodine deficiency, thyroid nodules can be present in as many as 30% of pregnant women (40). Severe iodine deficiency in pregnant women has been associated with increased rates of pregnancy loss, stillbirth, and increased perinatal and infant mortality (41).

Normal levels of thyroid hormone are essential for neuronal migration, myelination, and other structural changes of the fetal brain. Because thyroid hormones are needed throughout pregnancy, iodine deficiency affects both maternal and fetal thyroid hormone production, and insufficient iodine intake can lead to detrimental effects. Specifically, maternal and fetal iodine deficiency in pregnancy have adverse effects on the cognitive function of offspring (42–45). Children whose mothers were severely iodine deficient during pregnancy may exhibit cretinism, characterized by profound intellectual impairment, deaf-mutism, and motor rigidity. Iodine deficiency is the leading cause of preventable intellectual deficits worldwide (42).

Universal salt iodization is the most cost-effective way of delivering iodine and improving maternal and infant health (46).

Question 5: What is the impact of mild to moderate iodine deficiency on the mother, fetus, and child?

Groups of pregnant women whose median UICs are 50–150 $\mu\text{g/L}$ are defined as mildly to moderately iodine deficient. Women with mild to moderate iodine deficiency during pregnancy are at increased risk for the development of goiter (39) and thyroid disorders (47). Low maternal UIC in pregnancy has been associated with reduced placental weight and neonatal head circumference (48). However, in areas with adequate dietary iodine intake, variations in maternal UICs have a limited influence on physical developmental outcomes (49). Mild to moderate maternal iodine deficiency has also been associated with attention deficit and hyperactivity disorders in children (50) as well as impaired cognitive outcomes (51–53). In an iodine-deficient area iodized salt intake before pregnancy did improve maternal thyroid function; no difference in child neurodevelopment was noted (54), but improvement has been noted in other studies (55).

Question 6: What is the iodine status of pregnant and breastfeeding women in the United States?

Surveillance of urinary iodine values of the U.S. population has been carried out at intervals since 1971. Following a precipitous drop in urinary iodine values between 1971 and 1994, U.S. dietary iodine intake has stabilized (56–61). The U.S. population overall remains iodine sufficient. However, U.S. women of reproductive age are the most likely group to have low urinary iodine values. According to the World Health Organization (WHO) guidelines, median urinary iodine values for pregnant women between 149 and 249 $\mu\text{g/L}$

are consistent with optimal iodine intake (42). In the 2005–2010 National Health and Nutrition Examination Survey (NHANES) surveys, the median UIC for U.S. pregnant women was 129 $\mu\text{g/L}$, consistent with mild iodine deficiency (62). Current data regarding iodine sufficiency among lactating U.S. women are very limited. It is possible that a subset of pregnant and lactating U.S. women may have mildly to moderately inadequate dietary iodine intake resulting in insufficient amounts of iodine in the breast milk to meet infants' dietary requirements, but studies have been small and inconsistent (63–65).

Question 7: What is the iodine status of pregnant and breastfeeding women worldwide?

Since 1990, the number of households worldwide using iodized salt has risen from less than 20% to more than 70% (66). Despite these advances, however, 30 countries remain iodine deficient, and iodine deficiency remains the leading cause of preventable intellectual deficits worldwide (44). Recent reports highlight the value of iodized salt in correcting iodine deficiency states in India (67,68), although remaining iodine deficiency was noted in one report (69). Only 6% of pregnant women had low UICs in rural Bangladesh (70), whereas 80% had low urine iodine concentrations in Pakistan (71). In Shanghai, China, iodine deficiency was noted in a subset of pregnant women (72), whereas in Shenyang city slight iodine excess was noted, with a consequent increase in subclinical hypothyroidism (73). Iodine status in Korea is more than adequate (median UIC 427 $\mu\text{g/L}$) (74). Iodine status in Japan is also more than adequate (median UIC 328 $\mu\text{g/L}$) (9). On the African continent, iodine status in pregnancy was inadequate in Niger (75) and was also poor in Ethiopia (76). Iodine nutrition in Iran was sufficient (77). In Brazil, median UIC was 138 $\mu\text{g/L}$, possibly due to reduced concentration of iodine in salt (78). In Europe many countries, including Belgium, the Czech Republic, Denmark, France, Latvia, Norway, Spain, and the United Kingdom, have recorded significant iodine deficiency in their pregnant populations (47,79–85).

Question 8: Does iodine supplementation in pregnancy and lactation improve outcomes in severe iodine deficiency?

In areas of severe iodine deficiency, iodine supplementation of mothers prior to conception or in early pregnancy results in children with improved cognitive performance relative to children of mothers given a placebo (86–88). The prevalence of cretinism and other severe neurological abnormalities is significantly reduced (89). Maternal iodine supplementation in severely iodine-deficient areas also decreases rates of stillbirth and neonatal and infant mortality (90,91). Oral administration of iodized oil can increase birth weight in addition to correcting iodine deficiency (92).

A recent randomized clinical trial demonstrated that in moderately to severely iodine-deficient areas without universal salt iodization, lactating women who receive one dose of 400 mg oral iodized oil after delivery can provide adequate iodine to their infants through breast milk for at least 6 months. Direct infant iodine supplementation was less effective at improving infant iodine status (93).

Question 9: Does iodine supplementation in pregnancy and lactation improve outcomes in mildly to moderately iodine-deficient women?

Eight controlled trials of iodine supplementation in mildly to moderately iodine-deficient pregnant European women have been published (94–101), although doses and timing of iodine supplementation varied and only two trials examined effects on offspring development. Iodine supplementation of moderately deficient pregnant women appears to consistently decrease maternal and neonatal thyroid volumes and thyroglobulin (Tg) levels. Effects on maternal thyroid function have been mixed, with significant maternal TSH decreases with supplementation described in four (88,90,91,95) of the eight published trials, and increases in maternal T4 or FT4 noted in just two (90,95). A reduction in cord TSH also indicates improvement in gestational iodine status (102).

In two nonrandomized studies, neurodevelopmental outcomes were improved in children from mildly to moderately iodine-deficient areas whose mothers received iodine supplementation early in pregnancy (87,95). Another study failed to show neuropsychological improvement in 16-month-old children of mothers who received supplementation (103). The timing of supplementation is likely to be critical because the beneficial effects of iodine on offspring development appeared to be lost if supplementation is started after 10–20 weeks gestation. If iodine supplementation is started before pregnancy in iodine-deficient women better maternal thyroid function can be observed but, depending on dose and the timing of initiation, supplementation may not fully correct iodine deficiency in an already iodine-deficient population (104). A meta-analysis concluded that iodine supplementation improves some maternal thyroid indices and may benefit aspects of cognitive function in school age children, even in marginally iodine-deficient areas (105). Another review highlighted the lack of high-quality evidence in relation to these outcomes and suggested that randomized controlled trial (RCTs) may not be feasible where iodine supplementation is common (106).

Recently, there has been controversy regarding whether it is ethical to perform randomized clinical trials of iodine supplementation in pregnancy in regions that are mildly to moderately iodine deficient (107–109). Caution in accepting the necessity of supplementation has been expressed, especially in areas where iodized salt is already in use (110).

No trials to date have specifically examined the effects of iodine supplementation in lactation in mildly to moderately iodine-deficient regions.

Question 10: What is the recommended daily iodine intake in women planning pregnancy, women who are pregnant, and women who are breastfeeding?

Iodine is an essential nutrient required for thyroid hormone production and is primarily derived from the diet and from vitamin/mineral preparations. The U.S. Institute of Medicine recommended dietary allowances to be used as goals for individual total daily iodine intake (dietary and supplement) are 150 $\mu\text{g}/\text{d}$ for women planning a pregnancy, 220 $\mu\text{g}/\text{d}$ for pregnant women, and 290 $\mu\text{g}/\text{d}$ for women who are breastfeeding (111). The WHO recommends 250 $\mu\text{g}/\text{d}$ for pregnant and lactating women (39). This level is supported by a study of more than 7000 pregnant Chinese women in whom it was

found that subclinical hypothyroidism and hypothyroxinemia were least common at the urinary iodine range of 150–249 $\mu\text{g}/\text{L}$, but risk for both these abnormalities rose when the UIC was lower or higher than this range (73).

Dietary iodine sources vary regionally. Sources of iodine in the United States diet have been difficult to identify, in part because there are a wide variety of potential sources and food iodine content is not listed on packaging. Iodized salt remains the mainstay of iodine deficiency disorder elimination efforts worldwide. However, salt iodization has never been mandated in the United States and only approximately 50% of salt sold for household use in the United States is iodized (112). In the United States, dairy foods are another important source of dietary iodine because of iodine in cattle feed and the use of iodophor disinfectants by the dairy industry (113–115). Commercially baked breads have been another major source of iodine in the United States because of the use of iodate bread conditioners (115). However, the use of iodate bread conditioners has decreased over the past several decades. Other sources of iodine in the United States diet are seafood, eggs, meat, and poultry (116). Foods of marine origin generally have high concentrations of iodine because marine animals concentrate iodine from seawater (96–98), although the amount of iodine in different fish and shellfish species is quite variable.

In the United States, the dietary iodine intake of individuals cannot be reliably ascertained either by patient history or by any laboratory measure. Because of concerns that a subset of pregnant U.S. women may be mildly to moderately iodine deficient and an inability to identify individual women who may be at risk, the ATA has previously recommended 150 μg daily iodine supplementation for all North American women who are pregnant or breastfeeding (117). More recently the Endocrine Society, Teratology Society, and American Academy of Pediatrics have also advocated iodine supplementation for pregnant and lactating U.S. women (15,118,119). The goal is supplementation of dietary iodine intake, rather than its replacement. Special attention may need to be paid to those with dietary restrictions (e.g., lactose intolerant, gluten intolerant, low-carbohydrate, or vegan) because those individuals may have additional needs for supplementation (120).

Unfortunately, recommendations regarding iodine supplementation in the United States have not been widely adopted. In the NHANES 2001–2006 dataset, only 20% of pregnant women and 15% of lactating women reported ingesting iodine-containing supplements (121). When assessed in 2009, of the 223 types of prenatal multivitamins available in the United States, only 51% contained any iodine (122). However, this may recently have changed; in 2015 the Council for Responsible Nutrition, the U.S. supplement industry trade group, recommended that their members include 150 μg iodine in all prenatal multivitamin preparations. Iodine in U.S. prenatal multivitamins is typically derived either from potassium iodide or from kelp. The iodine content in prenatal multivitamin brands containing kelp may be inconsistent because of variability in kelp iodine content (123). Women consuming levothyroxine (LT4) regularly do not require supplemental iodine because the substrate is no longer needed for hormone formation.

The task force recommendations for iodine supplementation in the setting of lactation and breastfeeding are provided in Section XI.

■ RECOMMENDATION 5

All pregnant women should ingest approximately 250 μg iodine daily. To achieve a total of 250 μg iodine ingestion daily, strategies may need to be varied based on country of origin.

Strong recommendation, high-quality evidence.

■ RECOMMENDATION 6

In most regions, including the United States, women who are planning pregnancy or currently pregnant, should supplement their diet with a daily oral supplement that contains 150 μg of iodine in the form of potassium iodide. This is optimally started 3 months in advance of planned pregnancy.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 7

In low-resource countries and regions where neither salt iodization nor daily iodine supplements are feasible, a single annual dose of ~ 400 mg of iodized oil for pregnant women and women of childbearing age can be used as a temporary measure to protect vulnerable populations. This should not be employed as a long-term strategy or in regions where other options are available.

Weak recommendation, moderate-quality evidence.

■ RECOMMENDATION 8

There is no need to initiate iodine supplementation in pregnant women who are being treated for hyperthyroidism or who are taking LT4.

Weak recommendation, low-quality evidence.

Question 11: What is the safe upper limit for iodine consumption in pregnant and breastfeeding women?

Most people are tolerant of chronic excess dietary iodine intake due to a homeostatic mechanism known as the Wolff–Chaikoff effect (124,125). In response to a large iodine load, there is a transient inhibition of thyroid hormone synthesis. Following several days of continued exposure to high iodine levels, escape from the acute Wolff–Chaikoff effect is mediated by a decrease in the active transport of iodine into the thyroid gland, and thyroid hormone production resumes at normal levels (126).

Some individuals do not appropriately escape from the acute Wolff–Chaikoff effect, making them susceptible to hypothyroidism in the setting of high iodine intake. The fetus may be particularly susceptible, since the ability to escape from the acute Wolff–Chaikoff effect does not fully mature until about week 36 of gestation (127,128).

Tolerable upper intake levels for iodine have been established to determine the highest level of daily nutrient intake that is likely to be tolerated biologically and to pose no risk of adverse health effects for almost all individuals in the general population. The upper intake levels are based on total intake of a nutrient from food, water, and supplements and apply to chronic daily use. The U.S. Institute of Medicine has defined the tolerable upper limit for daily iodine intake as 1100 $\mu\text{g}/\text{d}$ in all adults, including pregnant women (96) and the WHO has stated that daily iodine intake

>500 μg may be excessive in pregnancy. Recent population data support the WHO threshold (73). The exception may be communities that have historically and consistently consumed greater than 500 μg daily without experiencing adverse effects (e.g., Japan).

Medications may be a source of excessive iodine intake for some individuals. Amiodarone, an antiarrhythmic agent (129), contains 75 mg iodine per 200 mg tablet. Iodinated intravenous radiographic contrast agents contain up to 380 mg/mL. Some topical antiseptics contain iodine, although systemic absorption is generally not clinically significant in adults except in patients with severe burns (130). Iodine-containing anti-asthmatic medications and expectorants are occasionally used. In addition, some dietary supplements such as kelp and some iodine preparations may contain very large amounts of iodine (several thousand times higher than the daily upper limit) and should not be taken. Ingestion of iodine and kelp supplements containing in excess of 500 $\mu\text{g}/\text{d}$ is not recommended in pregnancy or lactation (131).

Concern exists that some populations may be exposed to excess iodine, possibly resulting in a high prevalence of thyroid dysfunction (132,133), an increased rate of hyperthyrotrophinemia (134), and an increased rate of hyperthyroid newborns (135). In addition, iodine-induced hypothyroidism has been reported in infants exposed to excess iodine from radiocontrast agents (136). It should be recognized that even low-dose iodine supplementation may trigger thyroid autoimmunity in a small proportion of women (73). The exception may be communities that have historically and consistently consumed greater than 500 μg daily without experiencing adverse effects. In one study, approximately one third of Japanese pregnant women demonstrated higher UIC beyond 500 $\mu\text{g}/\text{Cr}$, but there were no problems in pregnancy progress and fetal development (9).

■ RECOMMENDATION 9

Excessive doses of iodine exposure during pregnancy should be avoided, except in preparation for the surgical treatment of GD. Clinicians should carefully weigh the risks and benefits when ordering medications or diagnostic tests that will result in high iodine exposure.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 10

Sustained iodine intake from diet and dietary supplements exceeding 500 μg daily should be avoided during pregnancy due to concerns about the potential for fetal thyroid dysfunction.

Strong recommendation, moderate-quality evidence.

V. THYROID AUTO-ANTIBODIES AND PREGNANCY COMPLICATIONS

Question 12: What is the prevalence of thyroid autoantibodies in pregnant women?

Anti-TPO or anti-Tg thyroid autoantibodies are present in 2% to 17% of unselected pregnant women (47,137–144). The prevalence of antibodies varies with ethnicity. In U.S. populations, thyroid antibodies are most frequent in Caucasian and Asian women and least frequent in African Americans

(139,145). Dietary iodine intake may also be associated with anti-thyroid Ab positivity during pregnancy. Shi and colleagues (73) recently demonstrated a U-shaped relationship between UICs and antibody positivity among pregnant women.

A recent study from Belgium in women seeking fertility treatment showed that both TPOAb and TgAb were present in 8% of women, while 5% demonstrated isolated Tg antibodies and 4% demonstrated isolated TPOAb concentrations (146). Those women with isolated TgAb positivity had a significantly higher serum TSH than women without thyroid autoimmunity. While the task force acknowledges that testing for thyroid autoimmunity using only TPOAb would likely miss a small proportion of women with isolated Tg antibodies, we note that the vast majority of studies investigating thyroid autoimmunity and clinical outcomes used only TPOAb measurements. For this reason, the task force recommends assessment of TPOAb when testing for the presence of thyroid autoimmunity.

Question 13: What is the natural history of antithyroid antibodies in pregnant women?

Question 14: How should euthyroid women who are thyroid antibody (Ab)-positive be monitored during pregnancy?

In women with thyroid autoimmunity, hypothyroidism may occur because of the stress of pregnancy because the ability of the thyroid to augment hormone production is compromised. In 1994, Glinoe *et al.* (147) performed a prospective study of 87 euthyroid (TSH ≤ 4 mU/L) women who were positive for TPOAb, TgAb, or both. Twenty percent of women in the study developed a serum TSH >4 mU/L during gestation despite normal prepregnancy TSH values. Antithyroid Ab titers were highest in the first trimester, although they decreased by about 60% over the course of gestation. Twelve years later, in a prospective study, Negro *et al.* (28) demonstrated similar results. The authors found that in TPOAb-positive euthyroid women, TSH levels increased as gestation progressed, from a mean of 1.7 mU/L (12th week) to 3.5 mU/L (term), with 19% of women having a supranormal TSH value at delivery. Because the risk of TSH elevation is increased in this population, increased surveillance of euthyroid thyroid Ab-positive women should occur.

TPO antibodies are able to cross the placenta. At the time of delivery, cord blood TPOAb levels strongly correlate with third-trimester maternal TPOAb concentrations (148). However, maternal passage of either TPOAb or TgAb is not associated with fetal thyroid dysfunction.

■ **RECOMMENDATION 11**

Euthyroid pregnant women who are TPOAb or TgAb positive should have measurement of serum TSH concentration performed at time of pregnancy confirmation and every 4 weeks through midpregnancy.

Strong recommendation, high-quality evidence.

Question 15: Should euthyroid women with thyroid autoimmunity be treated with selenium?

Some studies evaluating nonpregnant women have shown that selenium can diminish TPOAb concentrations (149–152).

However, this reduction has not been observed in all studies (153). Negro *et al.* (154) noted that euthyroid TPOAb-positive pregnant women randomized to treatment with 200 $\mu\text{g/d}$ selenium not only had a significant decrease in the frequency of postpartum thyroid dysfunction ($p < 0.01$) but also had lower TPOAb concentrations during pregnancy compared to those in the untreated group. Importantly, this trial did not measure urinary iodine, a potential confounder because iodine status may influence the thyroidal effects of selenium. However, in another recent randomized clinical trial (155) performed in mildly iodine-deficient British pregnant women, treatment with 60 μg of selenium daily did not affect TPO concentrations or TPOAb positivity. Thus, conflicting data regarding selenium supplementation make any generalized recommendation unreliable, especially to regions with different intakes of iodine, selenium, or both. In addition, patients treated with selenium could be at higher risk for developing type 2 diabetes mellitus (156). For these reasons, the risk-to-benefit comparison does not presently support routine selenium supplementation of TPOAb-positive women during pregnancy.

■ **RECOMMENDATION 12**

Selenium supplementation is not recommended for the treatment of TPOAb-positive women during pregnancy.

Weak recommendation, moderate-quality evidence.

Question 16: Is there an association between thyroid antibodies and sporadic spontaneous pregnancy loss in euthyroid women?

Spontaneous pregnancy loss (miscarriage), occurs in 17%–31% of all gestations (157,158). A spontaneous pregnancy loss is usually defined as one occurring at less than 20 weeks of gestation. The individual risk varies according to clinical factors including maternal age, family history, environmental exposures, and medical comorbidities (159). Pregnancy losses are a significant emotional burden to patients and may also result in bleeding, infection, pain, and need for surgical intervention.

Endocrine disorders have been previously recognized as risk factors for spontaneous pregnancy loss. Patients with poorly controlled diabetes mellitus may have up to a 50% risk of loss (160). Thyroid dysfunction has similarly been associated with increased pregnancy loss (161). Stagnaro-Green and colleagues (162) first demonstrated an association between pregnancy loss and thyroid antibodies in a prospective observational study. Patients who were positive for thyroid antibodies (TPOAb, TgAb, or both) demonstrated a 2-fold increase in the risk for pregnancy loss (17% vs. 8.4%, $p = 0.01$). Since that time, numerous other studies have examined the association between maternal antithyroid Ab status and pregnancy loss risk, showing similar findings. In a recent meta-analysis of eight case-control studies, the pooled OR for pregnancy loss in women with thyroid autoimmunity versus women without antithyroid antibodies was 2.55 [95% CI 1.42–4.57]. Meta-analysis of 14 cohort studies showed a similar increased OR of 2.31 [95% CI 1.90–2.82] (163). However, Ab-positive women were noted to be slightly older and to have slightly higher serum TSH values. In support of these data, a separate meta-analysis similarly found an increase in loss rate among thyroid Ab-positive women (OR

3.90 for cohort studies [95% CI 2.48–6.12]; OR 1.80 for case control studies [95% CI 1.25–2.60]) (164).

Although a clear association has been demonstrated between thyroid antibodies and spontaneous pregnancy loss, it does not prove causality and the underlying mechanisms for such an association remain unclear. Three research groups have demonstrated one possible mechanism through increased fetal resorption in active immunization murine models (165–167). Several other mechanistic hypotheses have been proposed, including Ab-mediated mild thyroid hypofunction, cross-reactivity of antithyroid antibodies with hCG receptors on the zona pellucida, the presence of concurrent non-organ-specific autoimmunity, and increased levels of endometrial cytokines in women with thyroid autoimmunity (168).

Question 17: Is there an association between thyroid antibodies and recurrent spontaneous pregnancy loss in euthyroid women?

Recurrent pregnancy loss is defined as either two consecutive spontaneous losses or three or more spontaneous losses, and it may occur in up to 1% of all women (169). Several causes have been reported, including parental chromosomal anomalies, immunologic derangements, uterine pathology, and endocrine dysfunction (170).

In a case-control study, Iravani and colleagues (171) reported that patients with primary recurrent pregnancy losses (three or more) had a higher incidence of TgAb and/or TPOAb positivity (OR 2.24 [95% CI 1.5–3.3]). Kutteh *et al.* (172) reported similar findings, with an increased rate of positivity for TgAb, TPOAb, or both in 700 women with recurrent pregnancy loss compared to 200 healthy controls (22.5% vs. 14.5%, $p=0.01$). On the other hand, in a prospective observational study, Esplin and colleagues (173) demonstrated no difference in positivity for TgAb, TPOAb, or both between patients with recurrent pregnancy loss and healthy controls. Pratt and colleagues (174) reported a higher rate of subsequent pregnancy loss in patients with recurrent losses and thyroid Ab positivity. In a larger study with a similar population, Rushworth and colleagues (175) reported no significant difference in live birth rates between women with recurrent losses who were TgAb or thyroid microsomal Ab positive and those who were not. In a case-control study, Lata *et al.* (176) found that the prevalence of TPO antibodies was higher in women with recurrent pregnancy loss than in healthy pregnant controls without a history of recurrent loss (31% vs. 18%, $p=0.031$). A meta-analysis of eight studies that included 460 Ab-positive patients and 1923 controls noted a significant association between thyroid Ab positivity and recurrent pregnancy loss (OR 2.3 [95% CI 1.5–3.5]) (177).

Thus, the data for an association between thyroid antibodies and recurrent pregnancy loss are less robust than for sporadic loss. This finding may be because recurrent pregnancy loss has many potential causes, and endocrine dysfunction may only account for 15%–20% of all such cases (170). Many of the studies described above did not control for other potential causes of recurrent losses. One intriguing study reported an apparent interaction of antiphospholipid antibodies and antithyroid antibodies in the risk for recurrent pregnancy loss (178). In support of this, Kim and

colleagues (179) reported that women with recurrent pregnancy loss who were antithyroid Ab positive also demonstrated higher levels of anticardiolipin Ab and other non-organ-specific antibodies.

Question 18: Does treatment with LT4 or intravenous immunoglobulin therapy decrease the risk for pregnancy loss in euthyroid women with thyroid autoimmunity?

Negro and colleagues (28) reported a prospective, randomized interventional trial of LT4 in euthyroid patients who were TPOAb positive. The researchers reported a significantly decreased rate of pregnancy loss in the LT4-treated group (3.5% vs. 13.8%, $p<0.05$). A limitation of the study is that the mean estimated gestational age at starting LT4 was 10 weeks, and all but one of the eight losses in the untreated group had occurred before 11 weeks. This finding raises uncertainty as to the impact of the intervention upon the endpoint.

However, in a separate nonrandomized, retrospective study, Lepoutre *et al.* (180) analyzed data from 65 TPOAb-positive pregnant women with serum TSH values of 1–3.5 mU/L at the first antenatal visit. Thirty-four of these women were treated with 50 µg LT4 daily starting at a mean 10 weeks gestation, while the others were not treated. None of the LT4-treated women miscarried, but 5 of 31 untreated women (16%) experienced pregnancy loss. Although limited in nature, these data support the findings of Negro *et al.* (28) and begin to suggest a potential benefit to this treatment approach. The underlying mechanism is, however, uncertain.

Three small nonrandomized case series have been published on the use of intravenous immunoglobulin (IVIG) therapy for the prevention of recurrent pregnancy loss in women with antithyroid antibodies (181–183). The live birth rates ranged from 80% to 95%, and the one study with a control group (consisting of women who refused IVIG therapy) reported a highly significant improvement in live births in the IVIG-treated cohort (95% vs. 0% $p=0.001$) (182). Comparison of a LT4 intervention to an IVIG intervention in one study demonstrated a higher rate of term delivery in the LT4-treated group (183). However, all three studies had serious design flaws including small sample sizes, heterogeneous patient populations, lack of or limited randomization, and differences in the timing of treatment initiation. Nonetheless, intervention trials with LT4 (and less so IVIG) in euthyroid TPOAb-positive women with recurrent abortion appear to show a possible decrease in miscarriage rates. Further randomized trials are needed to better understand the effectiveness of both LT4 and IVIG intervention. At present, however, the cost, complexity, and side effect profile associated with IVIG infusion must be noted and make its use undesirable given the questionable benefit. In contrast, LT4 administration in low dosage (25–50 µg/d) is safe. Therefore, its use among patients with recurrent pregnancy loss may be reasonably considered in the setting of early gestation, especially when no other known cause of prior pregnancy loss has been identified.

The task force makes note that two randomized clinical trials are currently ongoing. The Thyroid AntiBodies and LEvoThyroxine study (TABLET) trial in the United Kingdom is randomizing euthyroid TPOAb-positive women with a

history of infertility or recurrent losses to either LT4 or placebo to assess effects on live birth rates. Separately, the T4Lifetrial in the Netherlands is examining the effects of LT4 treatment in euthyroid pregnant women with a history of recurrent loss. The primary outcome of this trial is the live birth rate.

■ RECOMMENDATION 13

Intravenous immunoglobulin treatment of euthyroid women with a history of recurrent pregnancy loss is not recommended.

Weak recommendation, low-quality evidence.

■ RECOMMENDATION 14

Insufficient evidence exists to conclusively determine whether LT4 therapy decreases pregnancy loss risk in TPOAb-positive euthyroid women who are newly pregnant. However, administration of LT4 to TPOAb-positive euthyroid pregnant women with a prior history of loss may be considered given its potential benefits in comparison with its minimal risk. In such cases, 25–50 µg of LT4 is a typical starting dose.

Weak recommendation, low-quality evidence.

Question 19: Is there an association between thyroid autoantibody positivity and premature delivery?

Preterm deliveries are defined as those occurring before 37 weeks gestation. In the United States, 11.4% of all births are preterm (184). Preterm birth remains one of the most prevalent and morbid perinatal complications. It is the leading cause of neonatal death and is associated with increased risks for acute respiratory, gastrointestinal, immunologic, central nervous system (CNS), hearing, and vision problems, as well as longer-term motor, cognitive, visual, hearing, behavioral, and growth impairment (185). A decade ago, the annual cost of preterm delivery to the U.S. health care system was estimated at \$26 billion (185). Preterm birth has remained difficult to predict, prevent, and treat primarily because there are multiple potential causes and pathways that end in premature labor (186). Examples include infection, trauma, cervical insufficiency, premature rupture of membranes, and maternal medical conditions.

The relationship between thyroid autoantibodies and preterm delivery has been investigated with mixed results. Gli- noer *et al.* (187) reported in a prospective cohort that euthyroid women positive for either TPOAb or TgAb had a significantly increased incidence of preterm birth (16% vs. 8%, $p < 0.005$). Ghafoor *et al.* (188) evaluated 1500 euthyroid women and found an increase in preterm delivery in TPOAb-positive women compared to women who were TPOAb negative (26.8% vs. 8.0%, $p < 0.01$). In contrast, Iijima *et al.* (189) did not find an increased risk for preterm birth in women positive for seven different autoantibodies and thyroid antibodies. This study had an unusually low rate of preterm birth in both study and control groups (3% vs. 3.1%). Had- dow *et al.* (190) reported a significant increase in preterm premature rupture of the membranes, but not in preterm delivery, among women who were TPOAb and/or TgAb positive in the first trimester.

Importantly, three recent large prospective cohort studies failed to find significant associations between antithyroid antibody positivity and risk for premature delivery. However, in each of the studies, prematurity rates were nonsignificantly higher in TPOAb-positive women than in women who were antibody negative (19,190,191). By contrast, Karakosta *et al.* (192) found that euthyroid women (TSH < 2.5 mIU/L) with positivity for TPOAb, TgAb, or both in early pregnancy were at increased risk for spontaneous preterm delivery (RR 1.7 [95% CI 1.1–2.8]). In a small cohort of 395 pregnant women, Kumru and colleagues (193) noted that spontaneous preterm delivery (< 37 weeks) was more frequent in TPOAb-positive euthyroid (TSH < 2.6 mU/L) women (OR 2.5 [95% CI 1.06–5.89]). Negro and colleagues found that untreated TPOAb-positive women had a substantially higher risk for preterm delivery than women without TPOAb (RR 12.18 [95% CI 7.93–18.7]) (28). A 2011 meta-analysis of seven studies, including about 23,000 participants, concluded that there was an association between thyroid autoimmunity and preterm delivery (OR = 1.6 [95% CI 1.44–1.94]) (194). In support, a meta-analysis of five cohort studies including 12,566 women similarly concluded that a positive association existed between the presence of thyroid Ab and preterm birth (OR 2.907 [95% CI 1.17–3.68]) (164). Finally, a third meta-analysis of 11 prospective cohorts including 35,467 participants determined that the relative risk for delivery at less than 37 weeks for women with positivity for TgAb, TPOAb, or both was 1.41 [95% CI 1.08–1.84] (195). Interestingly, in subgroup analyses, TPOAb but not TgAb positivity was associated with prematurity. Together, these data suggest that thyroid autoantibody positivity is associated with increased risk for preterm delivery.

Question 20: Does LT4 treatment of euthyroid women who are thyroid autoantibody positive reduce risk for premature delivery?

In contrast to association studies, interventional studies of LT4 therapy for the prevention of preterm delivery are sparse. Negro *et al.* (28) reported an increased risk of preterm delivery among euthyroid TPOAb-positive women compared to euthyroid TPOAb-negative women in the only prospective interventional trial to date (22.4% vs. 8.2%, $p < 0.01$). The TPOAb-positive subjects were randomized to either treatment with LT4 or no treatment, with the LT4 dose based on TSH level. The treated group had a significantly lower rate of preterm delivery than did the untreated group (7% vs. 22.4%, $p < 0.05$). This finding has not been replicated. The ongoing TABLET study in the United Kingdom, a randomized clinical trial examining effects of LT4 treatment of euthyroid women with a history of infertility or recurrent pregnancy loss, will examine preterm birth as a secondary outcome. Therefore, at present, there are insufficient data from which to draw any conclusion regarding the utility of LT4 administration for the purpose of reducing preterm delivery.

■ RECOMMENDATION 15

Insufficient evidence exists to recommend for or against treating euthyroid pregnant women who are thyroid autoantibody positive with LT4 to prevent preterm delivery.

No recommendation, insufficient evidence.

Question 21: Is thyroid autoimmunity in euthyroid pregnant women associated with adverse obstetric or child outcomes other than pregnancy loss and premature birth?

Antithyroid antibodies have been associated with perinatal death in some (191) but not all (140,196) studies. In two pregnancy cohorts an increased risk for placental abruption was observed in women with thyroid autoimmunity and without clinical hypothyroidism (140,190). Maternal thyroid autoimmunity has also been linked to postpartum depression (197,198) and to neonatal respiratory distress syndrome (196).

Several studies have examined associations between maternal thyroid autoimmunity and child development. In a nested case-control study, Li *et al.* (199) reported lower motor and intellectual development at age 25–30 months in the offspring of euthyroid women who were TPOAb positive compared to children of TPOAb-negative controls. Williams *et al.* (200) followed 97 full-term mothers and their children, assessing child neurocognition at age 5.5 years. Lower perceptual performance and motor scores were described in children of TgAb-positive mothers, and lower perceptual performance scores noted in children with TgAb-positive cord blood. However, no neurodevelopmental outcomes were associated with maternal or infant TPOAb status. Wasserman and colleagues (141) described higher rates of sensorineural hearing loss (22.7% vs. 4.3%, $p=0.004$) at age 8 in children whose mothers were TPOAb positive in pregnancy compared to children of TPOAb-negative women. Although maternal thyroid function was not assessed in this study, the association remained significant after controlling for a known history of maternal hypothyroidism. In the same cohort, cognitive assessments were performed in children at ages 4 and 7 years (201). Maternal TPOAb positivity was associated with lower child intelligence quotient (IQ) at age 4, but effects were attenuated by age 7. The authors speculated that the lower IQ scores at age 4 might be mediated by sensorineural hearing loss. Ghassabian and colleagues (202) assessed a cohort of 3139 mother–child pairs in which child cognitive function was assessed at age 2.5 years and behavior assessed at age 3. Maternal TPOAb status did not predict child neurocognition, but TPOAb positivity was associated with externalizing problems in children (OR 1.64 [95% CI 1.17–2.29]), especially attention deficit/hyperactivity problems (OR 1.77 [95% CI 1.15–2.72]). This result remained significant after adjustment for maternal TSH. Finally, in a nested case–control study, Brown *et al.* (203) found that women with children on the autism spectrum were more likely to have had positive TPOAb during pregnancy than mothers of controls.

VI. THE IMPACT OF THYROID ILLNESS UPON INFERTILITY AND ASSISTED REPRODUCTION

Infertility is defined as the failure to achieve a clinical pregnancy after 12 or more months of regular unprotected sexual intercourse (204). Infertility affects 7.4% of U.S. women aged 15–44 years (205). Infertility is due to female factors in about 35% of cases, due to male factors in 30% of cases, and due to both female and male factors in 20% of cases. In approximately 15% of cases the cause of infertility is unknown (206).

Question 22: Is overt thyroid dysfunction associated with infertility in women?

Irregular menses may occur in women with overt hyperthyroidism. Krassas and colleagues (207) reported that menstrual irregularities were present in 22% of hyperthyroid patients as compared to 8% in age- and weight-matched euthyroid controls. However, in a cross-sectional study, the prevalence of hyperthyroidism (both subclinical and overt) was similar in infertile women compared to fertile controls (208).

The risk of infertility in women with overt hypothyroidism is less well studied. In a study of 171 hypothyroid women with TSH concentrations >15 mU/L, 68% reported having irregular menses, far higher than the 12% rate of menstrual irregularities reported by euthyroid controls (209). In one cross-sectional study among 129 infertile women, 5% had serum TSH levels >4.5 mU/L (210). In a separate cross-sectional study among women age 18–50 years, the prevalence of infertility was 52.3% in women with GD and 47% among women with Hashimoto's thyroiditis; however, thyroid function was not reported in either group (211). Thus, despite imperfect data, the majority of evidence appears to support an association between overt thyroid dysfunction and an increased risk of infertility. Thyroid dysfunction is also reversible, and treatment is generally safe and may exert a positive effect on fertility. Therefore, it is reasonable to treat overt thyroid dysfunction in infertile women, with the goal of normalizing thyroid function.

Question 23: Is subclinical hypothyroidism associated with infertility in women?

Different definitions of subclinical hypothyroidism have been used in different studies examining this question, and results have been inconsistent. In a cross-sectional study of 704 women with infertility, the prevalence of TSH elevations was 2.3%, similar to background rates in the general population (212). In a prospective study, Poppe and colleagues (208) did not find increased rates of subclinical hypothyroidism among infertile women but did report slightly higher median serum TSH levels (1.3 vs. 1.1 mU/L) in the infertile women compared to controls. However, in a retrospective study, higher rates of subclinical hypothyroidism (13.9% vs. 3.9%) were reported in infertile women as compared to fertile controls (213). Among women with baseline TSH values 2.5–5 mU/L, thyrotropin-releasing hormone (TRH) stimulation testing caused TSH increases to >30 mU/L in 46% of women with female-factor infertility versus 7% of women whose infertility was due to male factors (214). Among women presenting with infertility, TSH levels were highest among women with ovulatory dysfunction and unknown causes of infertility and lower among those women with tubal infertility and whose infertility was due to male factors (215). In an uncontrolled study, 94 infertile women with hypothyroidism (TSH >4.2 mU/L) were treated with LT4 (25–150 μ g/d) and 72 of them spontaneously conceived within 1 year (216). In a retrospective study, Yoshioka *et al.* (217) reported that 84.1% of infertile women with subclinical hypothyroidism (TSH >3.0 mU/L) successfully conceived and their infertility duration was shortened after LT4 therapy, suggesting that LT4 therapy may enhance fertility in patients with subclinical hypothyroidism. Importantly, whether or not LT4 treatment increases the likelihood of conception in subclinically hypothyroid women not undergoing ART has not been studied in controlled trials. Thus, insufficient data exist for

recommending for or against routine LT4 therapy in subclinically hypothyroid, thyroid autoantibody–negative infertile women who are attempting conception but not undergoing ART.

Question 24: Is thyroid autoimmunity linked to infertility in women?

Limited evidence suggests that women with female-factor infertility are more likely to be TPOAb positive than age-matched women who are not infertile, even if euthyroid (208). In addition, the prevalence of antithyroid antibodies may be higher in women with polycystic ovarian syndrome (PCOS) than in age-matched controls (218). Antithyroid antibodies are detectable in the ovarian follicles of women with thyroid autoimmunity and correlate with serum antibody levels (219), although whether such antibodies interfere with the fertilization potential of maturing oocytes is unknown. Among infertile women with PCOS, the presence of antithyroid antibodies has been associated with a decreased likelihood of developing ovarian follicles in response to treatment with clomiphene citrate (220).

■ **RECOMMENDATION 16**

Evaluation of serum TSH concentration is recommended for all women seeking care for infertility.

Weak recommendation, moderate-quality evidence.

■ **RECOMMENDATION 17**

LT4 treatment is recommended for infertile women with overt hypothyroidism who desire pregnancy.

Strong recommendation, moderate-quality evidence.

■ **RECOMMENDATION 18**

Insufficient evidence exist to determine if LT4 therapy improves fertility in subclinically hypothyroid, thyroid autoantibody–negative women who are attempting natural conception (not undergoing ART). However, administration of LT4 may be considered in this setting given its ability to prevent progression to more significant hypothyroidism once pregnancy is achieved. Furthermore, low dose LT4 therapy (25–50 µg/d) carries minimal risk.

Weak recommendation, low-quality evidence.

■ **RECOMMENDATION 19**

Insufficient evidence exists to determine if LT4 therapy improves fertility in nonpregnant, thyroid autoantibody–positive euthyroid women who are attempting natural conception (not undergoing ART). Therefore, no recommendation can be made for LT4 therapy in this setting.

No recommendation, insufficient evidence.

Question 25: Is maternal subclinical hypothyroidism associated with worse ART outcomes?

Question 26: Does treatment of subclinically hypothyroid women improve ART outcomes?

In a retrospective cohort study among women undergoing intrauterine insemination (IUI), the use of LT4 to treat subclinical hypothyroidism (TSH >2.5 mU/L) was associated

with higher pregnancy rates, and the use of LT4 to treat overt hypothyroidism with lower pregnancy rates, despite the fact that all LT4-treated patients had a serum TSH <2.5 at the time of IUI (221). In a separate prospective cohort study of 1477 women undergoing IUI, preconception serum TSH in the range of 0.4–4.99 mU/L was not associated with IUI parameters, pregnancy rates, or live birth rates per cycle (222).

The majority of evidence suggests that IVF outcomes do not differ between women with serum TSH <2.5 mU/L and those with very mild TSH elevations, defined as a TSH between 2.5 and 5 mU/L. A retrospective cohort of 195 IVF cycles that resulted in deliveries found that gestational age at birth was lower and mean birth weight was lower among women with preconception TSH >2.5 mU/L (223). In another retrospective cohort of 164 women undergoing IVF, clinical pregnancy rates were higher (22% vs. 9%, $p=0.045$) in women with TSH ≤2.5 versus those with TSH >2.5 mU/L (224). However, in a retrospective cohort of 1055 women undergoing their first IVF cycle, in age-adjusted analyses there was no difference in pregnancy, pregnancy loss, or delivery rates among women with serum TSH <2.5 mU/L versus those with TSH <4.5 mU/L (225). Another recent retrospective study examined IVF outcomes in 627 women without known thyroid disease and concluded that the presence of prepregnancy subclinical hypothyroidism (serum TSH >4.5 mU/L) did not affect pregnancy rates, live birth rates, or miscarriage rates (226). A case–control study among women undergoing first IVF cycles found no difference in embryo quality among women with serum TSH 0.45–2.5 mU/L versus 2.5–4.5 mU/L (227). Another retrospective cohort similarly reported no associations between serum TSH in the range of 0.5–4.5 mU/L and IVF outcomes (228). Finally, in a retrospective study which aimed to control for embryo quality by limiting the cohort to 1599 euploid blastocyst transfer cycles, variations in serum TSH ≤2.5 mU/L were not associated with differences in implantation, pregnancy loss, or live birth rates (229).

However, data generally demonstrate that treatment of more significant elevations in TSH concentrations (although still classified as subclinical hypothyroidism) appears beneficial. A randomized clinical trial was conducted in women aged 20–40 years with subclinical hypothyroidism (serum TSH >4.5 mU/L with normal FT4) who were undergoing IVF (230). A total of 64 women were randomized to treatment with LT4 (50 µg/d starting at the time of initiation of ovarian stimulation, and titrated to maintain TSH <2.5 mU/L in the first trimester in those women who did achieve pregnancy) versus placebo. The treated women had higher rates of clinical pregnancy, lower rates of miscarriage, and higher delivery rates. Another trial randomized 64 infertile women with subclinical hypothyroidism (TSH >4.2 mU/L with normal FT4) to treatment with 50 µg/d LT4 versus placebo and similarly reported higher pregnancy rates, lower pregnancy loss rates, and higher live birth rates in the treated women than in the control group (231). A recent meta-analysis pooled results of these trials with a third study (232) examining the effects of LT4 treatment for TPOAb-positive euthyroid women undergoing ART and concluded that although LT4 treatment did not have any effect on clinical pregnancy rates (pooled relative risk 1.75 [95% CI 0.90–3.38]), it did result in a higher delivery rate (pooled relative risk 2.76 [95% CI 1.20–6.44]) (233).

Together, these data suggest that subclinical hypothyroidism likely affects ART in a dose-related fashion, such

that impact worsens as TSH concentrations rise. It is also well known that TSH concentrations are variable and may differ by several milli-units per liter from week to week, despite no change in treatment (234). It therefore seems prudent to recommend treatment of subclinically hypothyroid women seeking pregnancy with ART for any TSH elevation >2.5 mU/L.

■ RECOMMENDATION 20

Subclinically hypothyroid women undergoing IVF or intracytoplasmic sperm injection (ICSI) should be treated with LT4. The goal of treatment is to achieve a TSH concentration <2.5 mU/L.

Strong recommendation, moderate-quality evidence.

Question 27: Is treated maternal hypothyroidism associated with worse ART outcomes compared to healthy controls?

In a retrospective cohort study, among 240 women aged ≤ 37 years undergoing first IVF retrieval cycles, 21 euthyroid (serum TSH 0.35 – 4.0 mU/L) women with treated hypothyroidism had lower pregnancy rates than the women without thyroid disease (235). However, a subsequent case–control study including 137 LT4-treated hypothyroid women (mean serum TSH 1.6 ± 0.7 mU/L) and 274 age-matched euthyroid controls determined that the presence of adequately treated hypothyroidism was not associated with inferior rates of pregnancy, implantation, or live birth following IVF/ICSI (236). Thus, no conclusion can be drawn from these data.

Question 28: Is maternal antithyroid Ab positivity associated with poorer outcomes following ART?

Studies examining ART outcomes in thyroid autoantibody-positive and -negative women have enrolled heterogeneous populations with differing underlying etiologies for infertility. Some, but not all studies have been limited to women who were euthyroid. Different ART/IVF protocols have also been employed in different places and over time. A meta-analysis of prospective cohort studies suggests that pregnancy rates following IVF do not differ between Ab-positive and Ab-negative women, but as discussed previously (see Section V), that risk of pregnancy loss is higher in women with thyroid autoantibodies positivity (237). However, a recent retrospective study examined IVF outcomes in 627 women without known thyroid disease and concluded that the presence of prepregnancy thyroid autoimmunity did not affect pregnancy rates, live birth rates, or pregnancy loss rates (226). Similarly, two recent retrospective cohorts reported no differences in pregnancy, pregnancy loss, or live birth rates in thyroid Ab-positive versus thyroid Ab-negative euthyroid women undergoing IVF with ICSI (238,239). Separately, a prospective cohort study found no differences in outcomes of IVF with ICSI in TPOAb-positive and/or TgAb-positive versus euthyroid women with Ab negativity (240). By contrast, a recent retrospective IVF cohort found that fertilization, implantation, and pregnancy rates were lower in 90 antithyroid Ab-positive compared to 676 Ab-negative women, but thyroid function was not reported in either group (241).

Question 29: Does treatment of antithyroid Ab-positive euthyroid women improve ART outcomes?

Based on a single small randomized clinical trial and one retrospective cohort, LT4 treatment for thyroid Ab-positive women without thyroid dysfunction undergoing IVF does not appear to improve outcomes (232,242). Two small trials suggest the potential for improved pregnancy rates in thyroid Ab-positive infertile women who are treated with glucocorticoids prior to ART (243,244). In the trial by Litwicka *et al.* (243), 60 euthyroid (TSH <2.5 mU/L) TPOAb- and/or TgAb-positive women undergoing IVF were randomized to prednisolone 5 mg/d starting from the day of oocyte retrieval and continuing through the first trimester. There were higher overall pregnancy (60% vs. 30%, $p=0.02$) and live birth (46.6% vs. 20%, $p=0.05$) rates in the treated versus untreated women. Turi and colleagues (244) randomized 48 TPOAb-positive infertile women to prednisone versus placebo for 4 weeks before IUI. The pregnancy rate was 33.3% in treated women compared to 8.4% in the placebo group ($p=0.03$). Pregnancy loss rates were not significantly different between the two groups. Although these small trials appear promising, the risks of corticosteroid use in early pregnancy are not well understood (245).

■ RECOMMENDATION 21

Insufficient evidence exists to determine whether LT4 therapy improves the success of pregnancy following ART in TPOAb-positive euthyroid women. However, administration of LT4 to TPOAb-positive euthyroid women undergoing ART may be considered given its potential benefits in comparison to its minimal risk. In such cases, 25–50 μ g of LT4 is a typical starting dose.

Weak recommendation, low-quality evidence.

■ RECOMMENDATION 22

Glucocorticoid therapy is not recommended for thyroid autoantibody-positive euthyroid women undergoing ART.

Weak recommendation, moderate-quality evidence.

Question 30: Does ovarian hyperstimulation alter thyroid function?

Although multiple ART protocols are currently in use, protocols for IVF or IVF with ICSI typically begin by inducing controlled ovarian hyperstimulation through the use of gonadotrophins, gonadotrophin-releasing hormone analogues, or gonadotrophin-releasing hormone antagonists in combination with gonadotrophins. Follicular development is monitored by ultrasound, and when leading follicles are large enough, hCG is administered to produce fully mature oocytes. Controlled ovarian hyperstimulation causes serum estradiol to rapidly rise to supraphysiologic levels (4000–6000 ng/L), comparable to those detected in late pregnancy (246). These hormonal manipulations may alter thyroid function. The presumed mechanism for this effect relates to the rise in TBG associated with high estrogen levels, which reduce free thyroid hormone concentrations and in turn feed back to cause serum TSH elevations. In addition, the administered hCG can directly stimulate thyroidal TSH receptors, causing increases in thyroid hormone and subsequent

decreases in TSH. A systematic review (247) found inconsistent effects of ovarian stimulation on serum thyroid hormones. For example, during and up to 1 month after ovarian stimulation serum TSH was increased in three of five studies, while FT4 was increased in two studies, decreased in one study, and unchanged in another. In some women (248–250), serum TSH levels increased to ≥ 2.5 mU/L following ovarian stimulation. Gracia and colleagues (251) reported increases in both serum TSH and FT4 during ovarian stimulation that peaked 1 week after hCG administration, with 44% of the women with baseline TSH ≤ 2.5 mU/L having increases to > 2.5 mU/L. Reinblatt *et al.* (252) reported that serum TSH increased in parallel with serum estradiol during the course of ovarian stimulation, with more marked TSH increases seen among TPOAb-positive women.

Ovarian hyperstimulation syndrome (OHSS) is a complication of controlled ovarian hyperstimulation in which increased vascular permeability results in fluid shifts from intravascular to third space compartments. Mild OHSS, manifested by mild ascites and abdominal symptoms, occurs in up to 20%–33% of all IVF cycles. Severe OHSS, which occurs in 0.1%–2% of IVF cycles, may be associated with thrombosis and respiratory distress, and is occasionally fatal (253). A recent case report described a patient who had subclinical hypothyroidism prior to IVF that was adequately treated with LT4. The patient rapidly developed overt hypothyroidism in the setting of severe OHSS (254). However, Poppe *et al.* (246) prospectively studied thyroid function in women undergoing controlled ovarian hyperstimulation and found no differences in serum TSH or FT4 levels between 25 women who developed OHSS and 52 women who did not.

Among women with adequately treated hypothyroidism (i.e., receiving LT4 therapy), ovarian stimulation has been reported to induce hypothyroidism by the time of egg retrieval (255). In one cohort study, increases in serum TSH during ovarian stimulation were more marked in LT4-treated women than in women without underlying hypothyroidism 1 week after hCG administration. However, TSH levels no longer differed between the groups by the time of pregnancy testing (251). In a prospective cohort study in 72 treated hypothyroid women with serum TSH < 2.5 mU/L prior to ovarian stimulation, serum TSH rose sequentially from 1.7 ± 0.7 mU/L at baseline to 2.9 ± 1.3 mU/L at the time of hCG administration, and to 3.2 ± 1.7 mU/L 16 days after hCG, with serum hCG > 2.5 mU/L in 46 (64%) of subjects by the time hCG was administered (256). A retrospective cohort study demonstrated that among treated hypothyroid women who undergo IVF, 84% required LT4 dose increases, most within the first 5–7 weeks of gestation (257). However, a small prospective study found that treated hypothyroid patients who conceived with the help of gonadotrophin therapy did not require larger LT4 dose increases than treated hypothyroid women who conceived spontaneously (258).

■ RECOMMENDATION 23

When possible, thyroid function testing should be performed either before or 1–2 weeks after controlled ovarian hyperstimulation because results obtained during the course of controlled ovarian stimulation may be difficult to interpret.

Weak recommendation, moderate-quality evidence.

■ RECOMMENDATION 24

In women who achieve pregnancy following controlled ovarian hyperstimulation, TSH elevations should be treated according to the recommendations outlined in Section VII. In nonpregnant women with mild TSH elevations following controlled ovarian stimulation, serum TSH measurements should be repeated in 2–4 weeks because levels may normalize.

Weak recommendation, moderate-quality evidence.

VII. HYPOTHYROIDISM AND PREGNANCY

Primary overt maternal hypothyroidism is generally defined as the presence of an elevated TSH and a decreased serum FT4 concentration during gestation, with both concentrations outside the (trimester-specific) reference ranges. In very rare cases, it is important to exclude other causes of abnormal thyroid function such as TSH-secreting pituitary tumors, thyroid hormone resistance, or central hypothyroidism with biologically inactive TSH. Several investigations report that at least 2%–3% of healthy, nonpregnant women of childbearing age have an elevated serum TSH (259,260). The prevalence may be higher in areas of iodine insufficiency. When iodine nutrition is adequate, the most frequent cause of hypothyroidism is autoimmune thyroid disease (Hashimoto's thyroiditis). Therefore, not surprisingly, thyroid autoantibodies can be detected in approximately 30%–60% of pregnant women with an elevated TSH concentration (19,260,261).

In the 2011 ATA guidelines, the upper reference limit for serum TSH concentration during pregnancy was defined as 2.5 mU/L in the first trimester, and 3.0 mU/L in the second and third trimesters. These cutoffs were predominantly based on the published reference ranges obtained from six pregnancy studies together comprising a total cohort of approximately 5500 subjects (12,13,18,262–264). Since that publication, additional much larger cohorts have published center-specific and trimester-specific pregnancy reference ranges. These analyses combine data from over 60,000 subjects (17,24,265–268). Importantly, this larger analysis demonstrates substantial population differences in the TSH upper reference limit (Table 4) (17,24,142,262,264–274). These differences may be partly attributable to differences in the iodine status between populations as well as the TSH assays used for analysis. However, these data also demonstrate important influences of BMI, geography, and ethnicity upon “normalcy” of TSH concentrations in pregnant women. In summary, substantial variation exists between populations, with many recent investigations confirming a more liberal upper TSH reference range in healthy pregnant women with no thyroid disease (269). Equally important, recent studies have also demonstrated an important additive influence of TPOAb positivity upon maternal thyroid status. Increasingly, there appears to be a greater risk for adverse events in women who are TPOAb positive compared to those who are TPOAb negative, even when thyroid function is identical. The reasons for this difference remain unclear. As described in Section V, some studies suggest that TPOAb-positive euthyroid women may be at increased risk for adverse clinical outcomes not observed in TPOAb-negative comparators (19). As a consequence, it is difficult to precisely define a universal TSH cutoff above which LT4 therapy should be initiated for all pregnant women. Rather, decisions about LT4

treatment must be based upon both measurement of thyroid function and TPOAb status.

Question 31: What is the definition of hypothyroidism in pregnancy?

Elevations in serum TSH concentrations during pregnancy should ideally be defined using pregnancy- and population-specific reference ranges. It is important to note that detection of an increased TSH concentration is not always synonymous with decreased FT4 concentrations. Frequently, elevated maternal TSH is detected when FT4 concentrations are normal. Conversely, low FT4 concentrations can be detected despite normal TSH concentrations. The latter situation is referred to as isolated hypothyroxinemia. Excepting the very rare scenarios noted above, serum TSH measurement remains the principal determinant of maternal thyroid status at the present time and should be used to guide treatment decisions and goals. Because substantial differences exist in the upper reference limit for TSH between different populations (Table 4), each practitioner and hospital should ideally seek to determine their own trimester-specific reference ranges, obtained from analysis of healthy, TPOAb-negative, and iodine-sufficient women. However, the task force recognizes that this goal is frequently not feasible. Therefore, this guideline also proposes a stepwise approach to evaluation, providing a practical guide for clinicians.

■ **RECOMMENDATION 25**

In the setting of pregnancy, maternal hypothyroidism is defined as a TSH concentration elevated beyond the upper limit of the pregnancy-specific reference range.

Strong recommendation, high-quality evidence.

■ **RECOMMENDATION 26**

The pregnancy-specific TSH reference range should be defined as follows:

- When available, population- and trimester-specific reference ranges for serum TSH during pregnancy should be defined by a provider's institute or laboratory and should represent the typical population for whom care is provided. Reference ranges should be defined in healthy TPOAb-negative pregnant women with optimal iodine intake and without thyroid illness.

Strong recommendation, high-quality evidence.

- When this goal is not feasible, pregnancy-specific TSH reference ranges obtained from similar patient populations and performed using similar TSH assays should be substituted (Table 4).

Strong recommendation, high-quality evidence.

- If internal or transferable pregnancy-specific TSH reference ranges are not available, an upper reference limit of ~ 4.0 mU/L may be used. For most assays, this limit represents a reduction in the nonpregnant TSH upper reference limit of ~ 0.5 mU/L.

Strong recommendation, moderate-quality evidence.

Question 32: How is isolated hypothyroxinemia defined in pregnancy?

Isolated hypothyroxinemia is typically defined as a FT4 concentration in the lower 2.5th–5th percentile of a given

population in conjunction with a normal maternal TSH concentration.

Question 33: What adverse outcomes are associated with overt hypothyroidism during pregnancy?

Overt maternal hypothyroidism has consistently been shown to be associated with an increased risk of adverse pregnancy complications (275) as well as detrimental effects upon fetal neurocognitive development (276). Specific adverse outcomes associated with overt maternal hypothyroidism include increased risks of premature birth, low birth weight, pregnancy loss, and lower offspring IQ. Abalovich *et al.* (277) demonstrated that women with overt hypothyroidism carry an estimated 60% risk of fetal loss when not adequately treated. Separately, Leung *et al.* (278) demonstrated a 22% risk of gestational hypertension in pregnant women with overt maternal hypothyroidism. Finally, Allan *et al.* (260) similarly described an increased risk of fetal death among pregnant women with overt disease. Together, these data demonstrate a clear association between overt maternal hypothyroidism and risk to the maternal–fetal unit.

Question 34: What adverse outcomes are associated with subclinical hypothyroidism during pregnancy?

Subclinical hypothyroidism is variably associated with an increased risk of adverse pregnancy outcomes in most, but not all studies, partly because separate studies use differing cutoffs to define an elevated TSH concentration. Also, many studies do not account for TPOAb status. Table 5 provides an overview of all available observational and prospective studies (inclusive of at least 400 subjects) investigating the effect of maternal subclinical hypothyroidism upon pregnancy (19,192,193, 259,279–285,287–289,291,294,295,306,314,318). Separately, Table 6 describes the additive adverse impact of TPOAb status upon maternal hypothyroidism (19,192,279,284,287–289,291, 295,314).

For the purposes of discussion, the investigations that have studied the association between elevated maternal TSH concentration and adverse clinical endpoints can be broadly grouped into three categories below based upon adverse endpoints. These include adverse effects on pregnancy outcome (i.e., pregnancy loss), adverse perinatal outcomes (i.e., premature delivery, hypertensive disorders), and adverse neurocognitive outcomes (IQ) in offspring.

Pregnancy loss. Early fetal loss naturally occurs in approximately 30% of pregnancies. Since the majority of pregnancy losses occur even before pregnancy is clinically recognized, pregnancy loss is a difficult study endpoint (158). Nevertheless, different studies have suggested a relationship between higher levels of maternal TSH and pregnancy loss. Negro *et al.* (286) reported a significantly higher pregnancy loss rate in TPOAb-negative women with TSH concentrations between 2.5 and 5.0 mU/L compared to those with TSH concentrations below 2.5 mU/L (6.1% vs. 3.6%). Similarly, Benhadi *et al.* (287) performed a prospective cohort study investigating the risk of pregnancy loss (defined as miscarriage or fetal or neonatal death) in 2497 Dutch women. In this cohort of pregnant women without overt hypothyroidism, the

TABLE 5. STUDIES INVESTIGATING SUBCLINICAL HYPOTHYROIDISM AND ADVERSE MATERNAL OUTCOMES

	Population selection	TFTs	Definition of SCH (TSH value in mU/L)	Outcome	Adjusted analysis	Finding	Comments
Casey <i>et al.</i> N = 17,298, (261)	Nonselected, population-based	<20 weeks	TSH >97.5th % (2.74–5.09 mU/L; gestational age specific)	Perinatal outcomes	No	Preterm delivery: unadjusted RR 1.8 [1.1–2.9] Abrupton: unadjusted RR 3.3 [1.1–8.2] No association	
Cleary-Goldman <i>et al.</i> N = 10,990 (289)	Nonselected, population-based	1st and 2nd trimesters (two measurements)	TSH >97.5th % (1st 4.28 mU/L; 2nd 3.93 mU/L)	Perinatal outcomes	Yes	No association	
Benhadi <i>et al.</i> N = 2497 (287)	Nonselected, population-based	Mean 13 weeks, up to end 2nd trimester	TSH >5.6 mU/L	Pregnancy loss	Yes	OR 1.8 [1.07–3.03] per TSH doubling	Small number of subjects in the high TSH cohort
Mannisto <i>et al.</i> N = 5805 (295)	Nonselected, population-based	Mean 10 weeks, <20 weeks	TSH >95th % (3.6 mU/L)	Perinatal outcomes	No	No association	Analysis of only 5808 out of 9247 subjects TPOAb and TgAb associated with perinatal mortality, after adjustment for age and parity TPOAb RR 3.2 [1.5–7.1]
Negro <i>et al.</i> N = 4123 (286)	Nonselected, population-based	<11 weeks	TSH 2.5–5.0 mU/L	Pregnancy loss Preterm delivery	No multivariate analysis; adjusted only for smoking	Pregnancy loss OR 1.16/mIU [1.00–1.34] Preterm delivery: no association	Study inclusive of only TPOAb-negative women
Mannisto <i>et al.</i> N = 5,805 (294)	Nonselected, population-based	Mean 11 weeks	TSH >95th % (3.6 mU/L)	Pregnancy complications	Yes	No association	Study primarily focused on endpoints of preeclampsia and gestational diabetes
Sahu <i>et al.</i> N = 633 (279)	Nonselected, population-based	13–26 weeks	TSH >5.5 mU/L (not pregnancy adjusted)	Pregnancy and perinatal complications	No	Cesarean section rate for fetal distress: OR 2.8 [1.2–6.6] ($p=0.04$)	
Su <i>et al.</i> N = 1017 (291)	Nonselected, population-based	Until 20 weeks	TSH >95% (3.77–4.35 mU/L; gestational age specific)	Pregnancy, perinatal and child development outcomes	Yes	Fetal distress: OR 3.65 [1.44–9.26] Preterm delivery: OR 3.32 [1.22–9.05], Poor vision development OR 5.34 [1.09–26.16] Neurodevelopmental delay OR 10.49 [1.01–119.19]	No data on TPOAb status. Small numbers of affected individuals leading to low power for statistical analysis
Karakosta <i>et al.</i> N = 1170 (192)	Nonselected, population-based	Mean 14 weeks	TSH >97.5%, Trimester specific (1st >2.53 mU/L; 2nd >2.73 mU/L)	GDM and perinatal outcomes	Yes	Low birth weight, RR 3.1 [1.2–8] GDM (only if TPOAb positive), RR 4.3 [2.1–8.9]	
Lazarus <i>et al.</i> N = 21,846 (306)	Nonselected, population-based	Median 12 weeks	TSH >97.5% Variable in time (UK >3.65 mU/L; Italy >3.50 mU/L)	Child IQ	N/A	No association	

(continued)

TABLE 5. (CONTINUED)

	Population selection	TFTs	Definition of SCH (TSH value in mU/L)	Outcome	Adjusted analysis	Finding	Comments
Korevaar <i>et al.</i> N=5971 (19)	Nonselected, population-based	Mean 13.2 weeks	TSH >97.5% (>4.04 mU/L)	Premature delivery	Yes	OR 1.87 (1.11–1.34)	Not significant finding after exclusion of TPOAb-positive subjects
Ong <i>et al.</i> N=2411 (280)	Selected population of women attending private clinic for screening	Mean 11.1 weeks	TSH >97.5% (TSH >2.15 mU/L)	Pregnancy and fetal outcomes	Yes	No association	
Chen <i>et al.</i> N=8012 (281)	Nonselected, population-based	14% 1st trimester, 33% 2nd trimester, 53% 3rd trimester	TSH >97.5% (1st >3.47 mU/L; 2nd >3.81 mU/L; 3rd >4.99 mU/L)	Hypertension, PROM, low birth weight, IUGR	Yes	Hypertension: OR 2.24 [1.25–4.02], PROM: 6.01 [3.98–9.10], IUGR 3.34 [1.75–6.38], and LBW 2.92 [1.65–5.16]	Majority of subjects assessed late in pregnancy (3rd trimester)
Godoy <i>et al.</i> N=5646 (282)	Nonselected, population-based	Median 13.2 weeks	TSH >97.5% (TSH >4.04 mU/L)	Childhood growth, body composition and cardiovascular characteristics.	Yes	No association	
Medici <i>et al.</i> N=5153 (285)	Nonselected, population-based	13.5 weeks	TSH >97.5% (TSH >4.04 mU/L)	Blood pressure and hypertensive disorders	Yes	No association	
Liu <i>et al.</i> N=3147 (288)	Selected healthy population, population-based	4–8 weeks	TSH >97.5% (TSH 5.22)	Miscarriage	Yes	Increased risk for women when TSH >5.22 (OR 3.4 [1.6–7.2]), or women with thyroid autoimmunity (OR 2.7 [1.4–5.1]). Synergistic worsening when both SCH and autoimmunity together.	Thyroid autoimmunity defined as either TPOAb or TgAb positivity
Leon <i>et al.</i> N=2170 (283)	Nonselected, population-based	Mean 13.5 weeks	TSH >95% (TSH >3.5 mU/L)	Pregnancy and fetal outcomes	Yes	No association	No analysis of subclinical hypothyroidism
Kumru <i>et al.</i> N=497 (193)	Selected, healthy population; population-based	10–12 weeks	TSH >95% (TSH level not reported)	Pregnancy and perinatal outcomes	Yes	Preterm delivery: OR 4.8 [1.89–12.42] No association with other maternal or perinatal outcomes.	Synergistic effect of thyroid hormone status and TPOAb status analyzed
Päkkilä <i>et al.</i> N=5131 (284)	Nonselected, population-based	Mean 10.5 weeks	TSH >3.1 mU/L in 1st or >3.5 mU/L in 2nd	ADHD	Yes	SCH was not associated with an increase in ADHD symptoms TSH was continuously associated with ADHD symptoms in girls	
Taylor <i>et al.</i> N=769 (318)	Women prescribed levothyroxine, population-based	1st trimester	3 endpoints: >2.5, >4.5, >10 mU/L	Miscarriage	Partial	TSH >4.5 associated with an increased risk of miscarriage. Graded effect with increasing TSH	

SCH, subclinical hypothyroidism; GDM, gestational diabetes mellitus; IUGR, intrauterine growth restriction; PROM, premature rupture of membranes; ADHD, attention deficit hyperactivity disorder; LBW, low birth weight.

TABLE 6. THYROID PEROXIDASE ANTIBODY POSITIVITY AND ADVERSE MATERNAL OUTCOMES

	Population selection	TFTs	Outcome	Adjusted analysis	Association with antibodies	Additive effects SCH	Comments
Cleary-Goldman <i>et al.</i> N = 10,990 (289)	Nonselected, population-based	1st and 2nd trimester (two measurements)	Perinatal outcomes	Yes	TPOAb positivity associated with PROM (OR 2.4–3.1)	Not investigated	
Benhadi <i>et al.</i> N = 2497 (287)	Nonselected, population-based	Mean 13 weeks, up to end of 2nd trimester	Pregnancy loss	Yes	TPOAb positivity not associated with miscarriage.		Adjustment for TSH did not change significant continuous association for TSH. Absolute risk very low
Mannisto <i>et al.</i> N = 5805 (295)	Nonselected, population-based	Mean 10 weeks, up to 20 weeks	Perinatal outcomes	No	If TgAb positive, a larger proportion of noncephalic presentation at birth (3.6 vs. 6.3%), as well as perinatal mortality (0.8 vs. 1.2%) If TPOAb positive, larger proportion of LGA (5.0 vs. 2.4%) and perinatal mortality (2.4 vs. 0.8%)	Not investigated	
Karakosta <i>et al.</i> N = 1170 (192)	Nonselected, population-based	Mean 14 weeks	GDM and perinatal outcomes	Yes	OR 1.7 for spontaneous preterm delivery.	TAI + TSH >97.5%: OR 4.3 for GDM TAI + TSH >97.5%: OR 3.1–3.7 for LBW	Thyroid autoimmunity defined as positive TPOAb and/or TgAb
Korevaar <i>et al.</i> N = 5971 (19)	Nonselected, population-based	Mean 13.2 weeks	Premature delivery	Yes	TPO pos. + TSH >97.5%: OR 2.53 TPO pos. + TSH >2.5 mU/L: OR 2.18 TPO pos.+ TSH >2mU/L: OR 1.75 (ns) TPO pos. + TSH>median: OR 1.4 (ns)		
Medici <i>et al.</i> N = 5153	Nonselected, population-based	13.5 weeks	Blood pressure and hypertensive disorders	Yes	No association for TPOAb pos.		
Liu <i>et al.</i> N = 3147 (288)	Selected healthy population, population-based	4–8 weeks	Miscarriage	Yes	Thyroid autoimmunity alone OR 2.71 TAI with TSH 2.5–5.22: OR 4.96 TAI with TSH >5.22: OR 9.56		TAI defined as positive TPOAb and/or TgAb with normal TSH and FT4
Päkkilä <i>et al.</i> N = 5131 (284)	Nonselected, population-based	Mean 10.5 weeks	ADHD symptoms	Yes	TPOAb positivity not associated with ADHD symptoms	Not investigated	Exclusion of TPOAb-positive women did not change the results

risk of child loss increased with higher levels of maternal TSH, although results should be interpreted with caution given the very small number of 27 cases studied as well as the heterogeneity of the study's endpoint. In an Australian cohort, early-pregnancy TSH levels >95th percentile were associated with an increased risk of miscarriage (OR 3.66) although subclinical and overt hypothyroid cases were pooled (292). A separate retrospective study that determined thyroid parameters in early-pregnancy samples obtained from 202 pregnancies that subsequently miscarried showed higher mean TSH and lower FT4 concentrations as well as a higher prevalence of TSH concentrations >97.5th percentile and FT4 concentrations <2.5th percentile compared to 3592 normal pregnancies (137). More recently, Liu and colleagues (288) demonstrated a graded increase in miscarriage risk as maternal TSH concentrations increased. This effect was augmented by the presence of TPOAb positivity.

Negro *et al.* (286) published data suggesting that subclinical hypothyroidism increases the risk of pregnancy complications in TPOAb-positive women. In this prospective trial of more than 4600 subjects, women were randomized to universal screening versus case finding (high-risk screening) during pregnancy, with subsequent LT4 treatment of TPOAb-positive women with TSH >2.5 mU/L. Low-risk women in the unscreened group had serum collected and stored for analysis post partum. Within the subset of women classified as low risk for hypothyroidism, treatment of TPOAb-positive women with TSH >2.5 mU/L resulted in a significant reduction in a composite endpoint of pregnancy complications compared to no treatment. The composite endpoint remains a significant study limitation because many variables were subjective in nature. Furthermore, it is critical to note that the primary study endpoint was nonsuperior, showing no benefit of universal screening and treatment (286) compared with screening of high-risk women only, because the primary, predefined endpoint analyzed the effects of LT4 treatment on both low-risk and high-risk subjects together. Importantly, all high-risk women in the study were tested and treated for elevated TSH values. Therefore, when combining both groups, the treatment effect on the low-risk group was diluted, leading to the conclusion of no superiority of universal screening. As such, this study should be viewed as providing important and provocative data that are worthy of further study (286). However, its conclusion that universal screening did not confer a benefit, combined with the difficulty in drawing conclusions from a composite endpoint, makes it challenging to translate into clinical practice.

Premature delivery and other pregnancy complications. The largest study investigating the association of maternal hypothyroidism and premature delivery was performed by Casey *et al.* (261) in a cohort of 17,298 pregnant women presenting for prenatal care. Subclinical hypothyroidism was associated with an increased risk of premature delivery at <34 weeks (4% vs. 2.5%, $p=0.01$), but not with premature delivery at <32 weeks (2.5% vs. 1%, $p=0.07$) or at <36 weeks (7% vs. 6%, $p=0.39$). This lack of continuous effect raises questions about the 34-week finding. A later study by Cleary-Goldman *et al.* (289) showed no association of an elevated TSH with prematurity at <37 weeks. Separately, other studies have also investigated this potential adverse relationship, albeit with conflicting results (192,260,266,286,290–292).

This variation can in part be explained by some studies pooling overt and subclinical hypothyroid cases together (260,292), while others used different TSH cutoff values (286,290), and yet others enrolled a very limited number of subjects (192,291). A recent study compared the value of using population-based reference range limits (TSH >97.5 percentile, defined as 4.0 mU/L) versus a fixed TSH cutoff of 2.5 mU/L (19). While a TSH >2.5 mU/L was not associated with premature delivery, 1.9 and 2.5 times increased risks of prematurity at <37 and <34 weeks, respectively, were observed among women with TSH >4.0 mU/L. Interestingly, this association no longer persisted after exclusion of TPOAb-positive women or women with comorbidities. This shows that these factors may be important confounders in various studies and underlines the importance of performing in-depth analyses of observed associations.

The majority of large studies focusing upon pre-eclampsia and hypertensive disorders, with one exception one (293), did not find associations of other pregnancy complications with elevated TSH (260,289,294). Mannisto and colleagues (294,295) evaluated the relationship between pregnancy outcomes and thyroid function tests obtained at 12 weeks gestation in 5805 women. No adverse association between thyroid function and perinatal mortality was noted. A separate large study investigating the relationship between subclinical hypothyroidism and birth weight showed no effect on very low (<2500 g) or high (>4000 g) birth weights (289). This finding was also confirmed by Mannisto *et al.* (295). A recent meta-analysis broadly analyzed pregnancy outcomes in relation to maternal thyroid status (296). The authors found an increasing risk of pregnancy complications (pregnancy loss, preterm delivery, and placental abruption) in relation to maternal subclinical hypothyroidism during early pregnancy, although subclinical hypothyroidism was variably defined across studies (296).

Together, despite some differences in study design, biochemical cutoffs applied and slightly differing endpoints, the above studies overall indicate an increasing risk of pregnancy-specific complications, most notably pregnancy loss and preterm delivery, in relation to elevated maternal TSH concentrations. Importantly, however, this effect is exacerbated by the presence of elevated TPOAb, such that any additive risk is apparent in TPOAb-positive women when TSH exceeds 2.5 mU/L. However, in TPOAb-negative women similar adverse risk is not consistently apparent until maternal TSH exceeds 5–10 mU/L.

Adverse neurocognitive effect on the offspring. The detrimental effects of maternal thyroid hypofunction on fetal neurocognitive development are less clear. In support of an adverse impact attributable to maternal hypothyroidism, data from a large case-control study demonstrated a seven-point reduction in IQ among children born to untreated overtly hypothyroid women compared to euthyroid controls (276). Findings also supported a delay in motor skill development, language development, and attention at 7–9 years of age. Subsequent studies have shown similar impact on children born to women with isolated hypothyroxinemia, (19,283, 297–304,314,317). Separately, three small studies analyzing only TPOAb positivity appear to similarly show an effect on neurocognitive outcome in the offspring but need to be confirmed in larger samples (199,201,305).

In contrast, the Controlled Antenatal Thyroid Screening (CATS) study was a large prospective RCT investigating the

benefit of population screening for elevated TSH concentrations and low FT4 concentrations in pregnant women. This study demonstrated no improvement in cognitive function when children of treated hypothyroid or hypothyroxinemic mothers were evaluated at 3 years of age. In this cohort, detection of either an elevated TSH or low FT4 triggered initiation of 150 µg/d LT4 therapy, at a mean of 13 weeks and 3 days gestation (306). Preliminary results of another large multicenter, randomized, controlled trial, the Randomized Trial of Thyroxine Therapy for Subclinical Hypothyroidism or Hypothyroxinemia Diagnosed During Pregnancy, have recently become available. This study screened 97,226 pregnant women in order to randomize 677 women with subclinical hypothyroidism and 526 women with isolated maternal hypothyroxinemia to LT4 treatment versus placebo at a mean time point of 17 weeks. Similarly, this study demonstrated no significant effect of treatment on offspring IQ at the age of 5 years (21). Other smaller studies support this conclusion (307–309).

Taken together, these prospective results provide insufficient evidence to conclude that treatment of subclinical hypothyroidism is associated with improved neurocognitive outcomes in offspring.

It should be noted, however, that the lack of positive data does not rule out a potential harmful effect, nor does it rule out the theoretical effectiveness of any intervention. It is important to note that such studies are highly difficult to complete and often enroll subjects with great heterogeneity with regard to important study parameters. For example, the timing of LT4 intervention during gestation likely plays an important role in the effect of any intervention. The two randomized controlled studies described above initiated LT4 treatment only at the completion of the first trimester or later—which may be too late to significantly impact neurodevelopment. Animal studies have suggested any window of opportunity is likely earlier in gestation (310,311). Similarly, the duration and severity of maternal hypothyroidism are likely important, yet virtually never controlled for, because all studies include only a single baseline measurement of TSH concentration during pregnancy. It therefore must again be emphasized that overt maternal hypothyroidism during pregnancy should be considered dangerous, and logic suggests that moderate (or even mild) maternal hypothyroidism may similarly impart risk. What remains uncertain is the nuanced understanding of how this risk is reduced or abated as the extent of maternal hypothyroidism is normalized or other variables are modified. This point should be emphasized as we consider that the most common cause of maternal hypothyroidism has dramatically changed over the last century. Formerly, severe iodine deficiency was common, while more recently, the principal cause of maternal hypothyroidism is maternal Hashimoto's disease. These disorders are physiologically different, though both may impart a similar phenotype demonstrating elevated maternal TSH concentrations.

Question 35: What adverse outcomes are associated with isolated hypothyroxinemia in pregnancy?

Pop and colleagues (312) initially reported a decrease in psychomotor test scores among offspring born to women

with FT4 indices in the lowest 10th percentile, despite having normal serum TSH concentrations. Similarly, Li *et al.* (199) observed a reduction in IQ among offspring of mothers suffering from either hypothyroidism or isolated hypothyroxinemia during the first trimester. In recent years, additional prospective, nonrandomized studies have similarly reported adverse child outcomes in children born to mothers with isolated hypothyroxinemia (19,192,193,259,279,280–284, 287–289,291,294,295,306,313,314,318) (see Tables 5 and 6). Adverse outcomes include lower IQ, language delay, worsened motor function, smaller head circumference, and an increased risk of autism. These data are derived from different populations across the world (China, Belgium, the Netherlands, Spain) with known differences in iodine status. In contrast to those studies investigating the association of elevated TSH concentrations with adverse pregnancy outcome, however, very few studies have investigated isolated hypothyroxinemia and adverse pregnancy outcomes (see Table 7), excepting birth weight (314–316) and premature delivery (19). Available data suggest an association with higher birth weight and higher risk of premature delivery. Interestingly, many large-scale studies demonstrate that the populations of women with elevated TSH concentrations are generally exclusive from those identified with low FT4 concentrations. For example, in the CATS study, approximately the same proportion of screened mothers were identified in the hypothyroxinemic and subclinically hypothyroid groups, with little overlap.

Overall, available evidence appears to show an association between hypothyroxinemia and cognitive development of the offspring, with uncertain effects on prematurity (314–316) and low birth weight (19). However, no studies exist in which LT4 administration has been shown to ameliorate such harmful effects. In the CATS trial (303), 499 women were identified with low FT4 concentrations and 242 randomized to treatment with LT4. This intervention failed to demonstrate any improvement in neurocognitive outcomes in the offspring at 3 years of age. However, women in this study received a fairly high dose (150 µg) of LT4 treatment and 10% needed a dose reduction because of biochemical or clinical signs of overtreatment. This is interesting because a recent study by Korevaar *et al.* (317) showed that both low and high FT4 concentrations may be associated with a decrease in child IQ and reduction of cerebral gray matter volume as assessed by magnetic resonance imaging.

Question 36: Should women with overt hypothyroidism be treated in pregnancy?

As mentioned above, numerous retrospective and case-control studies confirm the detrimental effects of overt hypothyroidism on both pregnancy and fetal health (318). A recent retrospective study of more than 1000 pregnant women on chronic LT4 replacement, showed that the risk of pregnancy loss increased proportionally to the degree of TSH elevation, with no increased risk associated with TSH normalization (318). Although no prospective, randomized investigation of LT4 intervention to improve obstetric outcomes or child development has occurred in pregnant women with overt hypothyroidism, such an investigation would be unethical to complete. Nonetheless, available data confirm the benefits of treating severe hypothyroidism during pregnancy.

TABLE 7. HYPOTHYROIDISM AND ADVERSE OBSTETRICAL AND NEONATAL OUTCOMES^a

	Population selection	TFTs	Definition of hypothyroxinemia	Outcome	Adjusted analysis	Main result	Comments
<i>Obstetrical outcomes</i>	Korevaar <i>et al.</i> N=5971 (19)	<18 weeks	TSH 2.5–97.5th percentile and FT4 < 2.5th percentile	Premature delivery	Yes	Hypothyroxinemia was associated with a 2.5-fold increased risk of premature delivery and a 3.6-fold increased risk of very premature delivery, for spontaneous premature delivery this was 3.4- and 4.2-fold, respectively.	Hypothyroxinemia also associated with the risk of PROM (2.4-fold increased risk), also in spontaneous deliveries (2.7-fold increased risk).
	Medici <i>et al.</i> N=5153	<18 weeks	TSH 2.5–97.5th percentile and FT4 < 2.5th percentile	Blood pressure and hypertensive disorders	Yes	No association	Hyperthyroidism and high normal FT4 were associated with hypertensive disorders.
	Leon <i>et al.</i> N=2170 (283)	<24 weeks	TSH 5–95th percentile and FT4 < 5th percentile	Birth weight	Yes	Hypothyroxinemia was associated with higher mean birth weight. FT4 was negatively associated with birth weight.	
<i>Neurocognitive outcomes</i>	Henrichs <i>et al.</i> N=3659 (297)	<18 weeks	TSH < 2.5 mU/L with FT4 < 10th percentile (mild) or FT4 < 5th percentile (severe)	Language delay (18 and 30 months) and non-verbal cognitive delay (30 months)	Yes	A 1.4 to 1.8-fold increased risk of expressive language delay (mild and severe hypothyroxinemia respectively). A 2-fold increased risk of non-verbal cognitive delay (severe hypothyroxinemia)	Outcomes of delay defined as <15th age- and sex-specific percentile.
	Craig <i>et al.</i> N=5560 (298)	Mean 17.3 and 16.8 weeks (cases and controls)	FT4 < 3rd percentile	Bayley Scale Scores at 2 years	Yes	Hypothyroxinemia was not associated with a lower score on cognitive, language or motor development domain.	After final selection, this study only had very low statistical power to detect a difference. Also the matching was suboptimal.
	Roman <i>et al.</i> N=4039 (299)	<18 weeks	TSH > 0.03 and < 2.5 mU/L and FT4 < 5th percentile	Combination of the Pervasive Developmental Scale of the CBCL checklist and the Social Responsiveness Scale	Yes	Hypothyroxinemia was associated with a higher mean autistic symptoms score and a 2-fold higher risk of pervasive development problems.	No association for mild hypothyroxinemia (<10th FT4 percentile) Results were more apparent in boys.

(continued)

TABLE 7. (CONTINUED)

<i>Population selection</i>	<i>TFTs</i>	<i>Definition of hypothyroxinemia</i>	<i>Outcome</i>	<i>Adjusted analysis</i>	<i>Main result</i>	<i>Comments</i>
Finken <i>et al.</i> <i>N</i> = 1765 (300)	<15 weeks	FT4 < 10th percentile	Computerized assessment utilizing Amsterdam Neuropsychological Tasks	Yes	Hypothyroxinemia was associated with a 41 millisecond slower response speed. FT4 was also negatively associated with reaction time.	Reaction time is associated with brain morphologic and neurocognitive outcomes.
Julvez <i>et al.</i> <i>N</i> = 1761 (301)	<20 weeks	Gestational age standardized FT4 < 10th percentile	Bayley Scale Scores at 14 months	Yes	Hypothyroxinemia was associated with a 2.2 point lower mean IQ <5th percentile showed a statistically borderline result of 3.4 point lower mean IQ.	Also an association for trend when the trend of <10th, <5th and <2.5th percentiles was investigated.
Ghassabian <i>et al.</i> <i>N</i> = 3727 (302)	<18 weeks	FT4 < 5th percentile	Non-verbal IQ (median age 6)	Yes	Hypothyroxinemia was associated with a 4.3 point lower mean IQ as compared to the rest of the population.	Hypothyroxinemia was not associated with brain morphology (on MRI), later paper showed this was inadequately analyzed.
Gyllenberg <i>et al.</i> <i>N</i> = 1010 (303)	Mean 11.0 and 10.8 weeks (cases and controls)	TSH 5th–95th percentile and FT4 < 10th percentile	Schizophrenia	Yes	Hypothyroxinemia was 1.75-fold as likely to be present in cases as compared to controls (11.8% vs. 8.6%, respectively).	Subclinical hyperthyroidism was 1.9-fold more likely to be present in cases versus controls.
Modesto <i>et al.</i> <i>N</i> = 3873 (304)	<18 weeks	TSH 0.1–2.5 mU/L and FT4 < 5th percentile	Conners' Parent Rating Scale–Revised Short Form (median age 8).	Yes	Hypothyroxinemia was associated with higher parent-reported ADHD scores.	
Korevaar <i>et al.</i> <i>N</i> = 3839 (317)	<18 weeks	Various FT4 cutoffs	Nonverbal IQ (median age 6), gray matter and cortex volume (MRI scans at median age 8)		For FT4 < 9th percentile an association as seen with lower mean offspring IQ. The peak was at FT4 < 5th percentile. Low FT4 was associated with lower gray matter and lower cortex volume (<i>N</i> = 646)	Similar effects for high FT4

^aAvailable prospective observational studies since 2010. MRI, magnetic resonance imaging.

■ RECOMMENDATION 27

Treatment of overt hypothyroidism is recommended during pregnancy.

Strong recommendation, moderate-quality evidence.

Question 37: Should women with subclinical hypothyroidism be treated in pregnancy?

Many prospective and retrospective studies have demonstrated an increased risk of pregnancy complications associated with mildly elevated maternal TSH concentrations, especially in TPOAb-positive women. However, only a small number of studies have investigated the impact of LT4 treatment on pregnancy complications in such women.

As noted above, a single RCT has demonstrated a potential benefit of LT4 intervention at ~9 weeks gestation. Importantly, this study documented a reduction in the adverse pregnancy composite outcome only in TPOAb-positive women with mild hypothyroidism (defined as a TSH >2.5 mU/L) (286). It should again be noted that the majority of women with subclinical hypothyroidism detected in this investigation were TPOAb negative, for whom no intervention or treatment was provided. This study also used a composite endpoint that included subjective endpoints such as cesarean section rates and postdelivery admission to the neonatal intensive care unit. Importantly, the authors' conclusion for their primary endpoint stated that universal screening for elevated TSH concentration in a broad population of pregnant women did not improve outcomes compared to a high risk screening strategy. Outcomes of universal screening compared to no screening were not assessed.

A separate RCT demonstrated a decrease in preterm delivery and pregnancy loss in euthyroid (defined as TSH <4.2 mU/L) TPOAb-positive women who were treated with LT4 beginning in the first trimester of pregnancy (28). However, the majority of pregnancy losses in the control group occurred before the average start of LT4 therapy.

Many studies have stratified the risk imparted by hypothyroidism according to TPOAb status and consistently show that this risk is higher in TPOAb-positive women (288). These data also suggest that the adverse impact associated with maternal TSH levels is apparent at lower TSH elevations in women known to be TPOAb positive compared to women who are TPOAb negative. Furthermore, two studies suggest a reduction in pregnancy loss when TPOAb-positive women are treated with LT4, even when biochemically euthyroid (28,180). Intervention trials have not been performed in TPOAb-negative women.

However, despite the limitations of available interventional trials of LT4 therapy in this subclinically hypothyroid group, the data taken in aggregate appear to suggest a benefit of treatment, especially as it applies to reducing miscarriage in TPOAb-positive women. Therefore, it seems reasonable to recommend or consider LT4 treatment for specific subgroups of pregnant women with subclinical hypothyroidism. The strength of such recommendations, however, should differ depending on TPOAb status, as will the strength of evidence supporting treatment for each subgroup. This recommendation also necessitates that any pregnant women with an elevated TSH concentration must also be evaluated for TPOAb status. In making the recommendation, the task force acknowledges the very low risk inherent in initiating low-dose

LT4 treatment. A dose of only 50 µg/d is typically required for effective treatment of subclinically hypothyroid women.

■ RECOMMENDATION 28

Pregnant women with TSH concentrations >2.5 mU/L should be evaluated for TPOAb status.

■ RECOMMENDATION 29

Subclinical hypothyroidism in pregnancy should be approached as follows:

- (a) LT4 therapy is recommended for
 - TPOAb-positive women with a TSH greater than the pregnancy-specific reference range (see Recommendation 1).

Strong recommendation, moderate-quality evidence.

- TPOAb-negative women with a TSH greater than 10.0 mU/L.

Strong recommendation, low-quality evidence.

- (b) LT4 therapy may be considered for
 - TPOAb-positive women with TSH concentrations >2.5 mU/L and below the upper limit of the pregnancy-specific reference range.

Weak recommendation, moderate-quality evidence.

- TPOAb-negative women and TPOAb-negative women with TSH concentrations greater than the pregnancy-specific reference range and below 10.0 mU/L.

Weak recommendation, low-quality evidence.

- (c) LT4 therapy is not recommended for
 - TPOAb-negative women with a normal TSH (TSH within the pregnancy-specific reference range or <4.0 mU/L if unavailable).

Strong recommendation, high-quality evidence.

Question 38: Should women with isolated hypothyroxinemia be treated with LT4 in pregnancy?

Although several studies have reported adverse outcomes in children born to mothers with isolated hypothyroxinemia, no interventional data have yet been published that demonstrate beneficial effects of LT4 therapy. One observational study analyzing women at 12, 24, and 32 weeks of pregnancy demonstrated delayed infant neurodevelopment in women with persistent hypothyroxinemia. However, when FT4 concentrations increased during pregnancy, infant development was not improved (312). Nevertheless, at present there are only two randomized, prospective, intervention trials in which women with a low FT4 were treated with LT4, at 13 and 17 weeks gestation, respectively (21,306). Both investigations failed to show any beneficial effect on cognitive development following LT4 administration, though a major limitation of the studies was the late timing of the intervention, after completion of the first trimester. Nevertheless, given the existing interventional data, treatment of isolated hypothyroxinemia cannot be recommended at this time.

■ RECOMMENDATION 30

Isolated hypothyroxinemia should not be routinely treated in pregnancy.

Weak recommendation, low-quality evidence.

Question 39: What is the optimal method of treating hypothyroidism in pregnant women?

Delivery of T4 is crucial for the developing fetal brain (313). The ratio of T4 to T3 in desiccated thyroid preparations is 4.2:1, which is significantly lower than the 14:1 ratio of secretion by the human thyroid gland. This relative excess of T3 leads to supraphysiologic maternal levels of T3 and relatively low levels of T4 (319). Patients using either desiccated thyroid or a treatment regimen combining T3 and T4 (320) are likely at risk for having insufficient transfer of maternal T4 to the fetal brain. It is notable that the majority of fetal T3 present in the CNS during pregnancy is derived from maternal T4 actively transported into this space. The fetal CNS is relatively impermeable to T3, which therefore argues against use of exogenous T3 during pregnancy. For these reasons, the task force feels that any T3-containing preparation should be avoided for the treatment of maternal hypothyroidism during pregnancy.

■ **RECOMMENDATION 31**

The recommended treatment of maternal hypothyroidism is administration of oral LT4. Other thyroid preparations such as T3 or desiccated thyroid should not be used in pregnancy.

Strong recommendation, low-quality evidence.

Question 40: What is the biochemical goal when treating hypothyroidism in pregnant women?

■ **RECOMMENDATION 32**

In parallel to the treatment of hypothyroidism in a general population, it is reasonable to target a TSH in the lower half of the trimester-specific reference range. When this is not available, it is reasonable to target maternal TSH concentrations below 2.5 mU/L.

Weak recommendation, moderate-quality evidence.

Question 41: Are there circumstances in which euthyroid women are at risk for hypothyroidism once pregnant?

In 1994, Glinoe *et al.* (147) performed a prospective study in 87 TPOAb-positive euthyroid women evaluated before and during early pregnancy. Twenty percent of women in the study developed a TSH >4 mU/L during gestation despite a normal TSH and no requirement for LT4 prenatally. This occurred despite the expected decrease in TPOAb titers during pregnancy (147). Negro *et al.* (28) demonstrated similar results in a prospective study. The authors found that in TPOAb-positive euthyroid women, TSH levels increased progressively as gestation progressed, from a mean of 1.7 mU/L (12th week) to 3.5 mU/L (term), with 19% of women having a supranormal TSH concentration at delivery. These findings confirm that an increased requirement for thyroid hormone occurs during gestation. In women who are TPOAb positive, both overt and subclinical hypothyroidism may occur because of a lack of ability of the thyroid to augment production when needed during pregnancy. Similarly, patients who undergo hemithyroidectomy or receive radioactive iodine and are euthyroid before preg-

nancy are at risk for developing elevated serum TSH levels during gestation.

In summary, euthyroid patients who are antithyroid Ab positive, post-hemithyroidectomy, or treated with radioactive iodine have an increased propensity for the development of hypothyroidism in gestation and should be monitored regularly.

Question 42: How should women with hypothyroidism or at risk for hypothyroidism be monitored through pregnancy?

In women at risk for hypothyroidism (TPOAb- or TgAb-positive, post-hemithyroidectomy, and/or post-radioactive iodine) increased surveillance is recommended. Based on findings extrapolated from investigations of treated hypothyroid women from early pregnancy onwards (321), it is reasonable to evaluate these women for TSH elevation approximately every 4 weeks during pregnancy. Serial testing is preferably continued through midpregnancy because the increased T4 demand continues throughout the first half of gestation.

A study by Yassa and colleagues (322) investigated the optimal timing of subsequent assessment of thyroid function following dose modification though this was in patients consuming LT4 prenatally and may not be generalizable to patients not taking LT4 but at risk for hypothyroidism.

■ **RECOMMENDATION 33**

Women with overt and subclinical hypothyroidism (treated or untreated) or those at risk for hypothyroidism (e.g., patients who are euthyroid but TPOAb or TgAb positive, post-hemithyroidectomy, or treated with radioactive iodine) should be monitored with a serum TSH measurement approximately every 4 weeks until midgestation and at least once near 30 weeks gestation.

Strong recommendation, high-quality evidence.

Question 43: How do treated hypothyroid women (receiving LT4) differ from other patients during pregnancy? What changes can be anticipated in such patients during gestation?

The major physiologic thyroid changes during pregnancy have been thoroughly studied. Total body T4 requirements are not static throughout gestation. Rather, data demonstrate that the total body T4 pool must increase by approximately 40%–50% to maintain a euthyroid state (321,323). In a healthy woman who becomes pregnant, the pregnancy hormone hCG plays a major role as a stimulus of maternal thyroid hormone production, especially throughout the first half of pregnancy. Together with pituitary TSH, placental hCG stimulates endogenous thyroid hormone production when an intact thyroid is present, and it helps to maintain a euthyroid state during gestation.

In women with known hypothyroidism, serum hCG and TSH cannot stimulate adequate T4 production. If exogenous LT4 is not adjusted, the increased demand of pregnancy will outstrip supply, and maternal hypothyroidism will occur. Clinical studies have confirmed that the increased requirement for thyroxine (or exogenous LT4) occurs as early as 4–6 weeks of pregnancy (321). Such requirements gradually

increase through 16–20 weeks of pregnancy and plateau thereafter until the time of delivery. These data provide the basis for recommending adjustments of LT4 dosage when affected women become pregnant and also for the timing of follow-up intervals for TSH in treated patients.

Question 44: What proportion of treated hypothyroid women (receiving LT4 prenatally) require changes in their LT4 dose during pregnancy?

Between 50% and 85% of LT4-treated hypothyroid women need to increase exogenous LT4 dosing during pregnancy (277,321,323). The incremental increase largely depends on the underlying etiology of the hypothyroidism. There is a greater likelihood that dose increases will be required in those patients without functional thyroid tissue (e.g., due to radioablation or surgery) in comparison with patients with Hashimoto's thyroiditis (324,325). The preconception level of TSH as well as other factors can also influence the rapidity and extent of LT4 augmentation necessary to maintain a euthyroid state during pregnancy. For example, variation and changes in maternal estrogen levels during pregnancy correlate with variations in the gestational requirements for LT4 (321).

The LT4 adjustment should be made as soon as possible after pregnancy is confirmed to reduce the probability of hypothyroidism. Normalization of TSH concentrations throughout gestation is the goal. For women receiving LT4 preconception, a prospective, randomized study has provided evidence that supports a single dose-adjustment strategy rather than a stepwise approach for LT4 dosage adjustment postconception (322). For euthyroid women receiving once-daily dosing of LT4 (regardless of the dosage), a recommendation to increase by two additional tablets weekly (nine tablets per week instead of seven tablets per week, giving a 29% increase) can effectively mimic gestational physiology and thus prevent maternal hypothyroidism during the first trimester (322). Another option is to increase the dosage of daily LT4 by approximately 25%–30%. Dosage augmentation should occur as soon as possible when a missed menstruation or suspected pregnancy occurs, and this should be discussed with every patient in the prepregnancy setting. Confirmatory biochemical testing should also occur simultaneously.

Question 45: How should preconception LT4 be adjusted in treated hypothyroid women (receiving LT4) planning pregnancy?

The difficulties inherent to achieving rapid, postconceptional TSH normalization have also focused attention upon preconception TSH modulation. Different cutoff values for preconception TSH, ranging from <1.2 to <2.5 mU/L have been advocated. In one study, only 17% of women with TSH <1.2 mU/L had to increase LT4 dose later during pregnancy (326). Given this, it is recommended that all treated hypothyroid women (currently receiving LT4) optimize thyroid parameters preconception. A maternal serum TSH concentration <2.5 mU/L is a reasonable goal for such women. Even lower preconception TSH values (<1.5 mU/L) could reduce the risk of TSH elevation during the first trimester (322), but a lower treatment target may not improve outcomes because

the LT4 dose can be immediately increased upon a positive pregnancy test. Furthermore, the process of achieving a TSH concentration at the lower end of the reference range could induce subnormal TSH concentrations in some patients. Though generally safe for any developing fetus, potential such effects upon conception and/or successful implantation are unknown.

■ **RECOMMENDATION 34**

Treated hypothyroid women of reproductive age should be counseled regarding the likelihood of increased demand for LT4 during pregnancy. Such women should also be counseled to contact their caregiver immediately upon a confirmed or suspected pregnancy.

Strong recommendation, high-quality evidence.

■ **RECOMMENDATION 35**

In hypothyroid women treated with LT4 who are planning pregnancy, serum TSH should be evaluated preconception, and LT4 dose adjusted to achieve a TSH value between the lower reference limit and 2.5 mU/L.

Strong recommendation, moderate-quality evidence.

■ **RECOMMENDATION 36**

Hypothyroid patients receiving LT4 treatment with a suspected or confirmed pregnancy (e.g., positive home pregnancy test) should independently increase their dose of LT4 by ~20%–30% and urgently notify their caregiver for prompt testing and further evaluation. One means of accomplishing this is to administer two additional tablets weekly of the patient's current daily LT4 dosage.

Strong recommendation, high-quality evidence.

Question 46: How should LT4 be adjusted postpartum?

The increased LT4 dose requirements during gestation are a function of pregnancy itself. Therefore, following delivery, maternal LT4 dosing should be reduced to prepregnancy levels, and a serum TSH assessed 6 weeks thereafter. However, a study demonstrated that more than 50% of women with Hashimoto's thyroiditis required an increase in the pregestational thyroid hormone dose in the postpartum period, presumably due to an exacerbation of autoimmune thyroid dysfunction post partum (327). In women started on LT4 during pregnancy for thyroid autoimmunity in the absence of TSH elevation, the LT4 can be stopped at delivery, with serum TSH assessment at 6 weeks post partum.

■ **RECOMMENDATION 37**

Following delivery, LT4 should be reduced to the patient's preconception dose. Additional thyroid function testing should be performed at approximately 6 weeks post partum.

Strong recommendation, moderate-quality evidence.

■ **RECOMMENDATION 38**

Some women in whom LT4 is initiated during pregnancy may not require LT4 post partum. Such women are candidates for discontinuing LT4, especially when the LT4

dose is $\leq 50 \mu\text{g/d}$. The decision to discontinue LT4, if desired, should be made by the patient and their caregiver. If LT4 is discontinued, serum TSH should be evaluated in approximately 6 weeks.

Weak recommendation, moderate-quality evidence.

Question 47: What is the outcome and long-term prognosis when maternal hypothyroidism is effectively treated through gestation?

Although untreated (or incompletely treated) hypothyroidism can adversely affect pregnancy, no data suggest that women with adequately treated subclinical or overt hypothyroidism have an increased risk of any obstetrical complication. Consequently, there is no indication for any additional obstetric testing or surveillance in pregnancies of women with either subclinical or overt hypothyroidism who are being monitored and treated appropriately.

Question 48: Except for measurement of maternal thyroid function, should additional maternal or fetal testing occur in treated hypothyroid women during pregnancy?

■ RECOMMENDATION 39

In the care of women with adequately treated hypothyroidism, no other maternal or fetal testing (such as serial fetal ultrasounds, antenatal testing, and/or umbilical blood sampling) is recommended beyond measurement of maternal thyroid function unless needed due to other circumstances of pregnancy. An exception to this is women with GD effectively treated with ^{131}I ablation or surgical resection, who require TSH receptor antibody (TRAb) monitoring.

Strong recommendation, moderate-quality evidence.

VIII. THYROTOXICOSIS IN PREGNANCY

Question 49: What are the causes of thyrotoxicosis in pregnancy?

Thyrotoxicosis is the clinical syndrome of hypermetabolism and hyperactivity that results when a person is exposed to supraphysiological amounts of thyroid hormones. The most common cause of thyrotoxicosis is hyperfunction of the thyroid gland (hyperthyroidism), and the most common cause of hyperthyroidism in women of childbearing age is autoimmune GD occurring before pregnancy in 0.4%–1.0% of women and in approximately 0.2% during pregnancy (328).

Less common non-autoimmune causes of hyperthyroidism in pregnancy include toxic multinodular goiter and toxic adenoma. Subacute painful or painless thyroiditis with passive release of thyroid hormones from a damaged thyroid gland are less common causes of thyrotoxicosis in pregnancy, and a number of other conditions such as a TSH-secreting pituitary adenoma (329), struma ovarii (330), functional thyroid cancer metastases, or germline TSH receptor mutations (331) are very rare. A special cause of thyrotoxicosis is overtreatment with or factitious intake of thyroid hormone.

More frequent than GD as the cause of thyroid function tests demonstrating hyperthyroxinemia is “gestational transient thyrotoxicosis,” which is limited to the first half of

pregnancy. This condition, characterized by elevated FT4 and suppressed serum TSH, is diagnosed in about 1%–3% of pregnancies. This frequency depends on the geographic area and is secondary to elevated hCG levels (332,333). Often it is associated with hyperemesis gravidarum, defined as severe nausea and vomiting in early pregnancy with more than 5% weight loss, dehydration, and ketonuria. Hyperemesis gravidarum occurs in 3–10 per 1000 pregnancies (334,335). Other conditions associated with hCG-induced thyrotoxicosis include multiple gestation, hydatidiform mole, and choriocarcinoma (336,337). Most cases present with marked elevations of serum hCG (14). A TSH receptor mutation leading to functional hypersensitivity to hCG also has been recognized as a rare cause of pregnancy-associated hyperthyroidism (338).

Question 50: What is the appropriate initial evaluation of a suppressed serum TSH concentration during the first trimester of pregnancy?

Serum TSH may decrease in the first trimester of normal pregnancy as a physiological response to the stimulating effect of hCG upon the TSH receptor. A peak hCG level typically occurs between 7 and 11 weeks gestation (339). In particular, a serum TSH below 0.1 mU/L (in some cases even undetectable) may be present in approximately 5% of women by week 11 of pregnancy (270). Any subnormal serum TSH value should be evaluated in conjunction with serum TT4 (or FT4) and T3 values. The biochemical diagnosis of overt hyperthyroidism is confirmed in the presence of a suppressed or undetectable serum TSH and inappropriately elevated serum TT4/FT4, or T3 (Section III).

Question 51: How can gestational transient thyrotoxicosis be differentiated from Graves' hyperthyroidism in pregnancy?

Diagnosing the cause of the disease is essential in any patient with thyrotoxicosis. In early pregnancy, the differential diagnosis in the majority of cases is between Graves' hyperthyroidism and gestational transient thyrotoxicosis (332,333). In both situations, common clinical manifestations include palpitations, anxiety, tremor, and heat intolerance. A careful history and physical examination is of utmost importance in establishing the etiology. The findings of no prior history of thyroid disease, no stigmata of GD (goiter, orbitopathy), a self-limited mild disorder, and symptoms of emesis favor the diagnosis of gestational transient thyrotoxicosis.

If other causes for thyrotoxicosis are suspected, measurement of TRAb is indicated. If this is negative or thyroid nodules are suspected based on clinical examination, a thyroid ultrasound should be performed to evaluate nodularity. Serum hCG is higher on average in gestational transient thyrotoxicosis than in patients with GD, but overlap is considerable and the clinical usefulness of such measurement is limited (340). No study has demonstrated usefulness of thyroid ultrasonography for differentiating between gestational transient thyrotoxicosis and GD. In the presence of a nodular goiter, a serum TT3 determination is helpful in assessing the possibility of the “T3 toxicosis” syndrome. TT3 determination may also be of benefit in diagnosing T3 thyrotoxicosis

caused by GD. In general, serum T3 tends to be disproportionately elevated more than T4 in cases of thyrotoxicosis caused by direct thyroid hyperactivity. In comparison, T4 tends to be disproportionately elevated beyond T3 when thyrotoxicosis is caused by destructive processes such as thyroiditis (341).

■ RECOMMENDATION 40

When a suppressed serum TSH is detected in the first trimester (TSH less than the reference range), a medical history, physical examination, and measurement of maternal serum FT4 or TT4 concentrations should be performed. Measurement of TRAb and maternal TT3 may prove helpful in clarifying the etiology of thyrotoxicosis.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 41

Radionuclide scintigraphy or radioiodine uptake determination should not be performed in pregnancy.

Strong recommendation, high-quality evidence.

Question 52: What is the appropriate management of gestational transient thyrotoxicosis?

The management of women with gestational transient thyrotoxicosis depends on the severity of symptoms. In women with hyperemesis gravidarum, control of vomiting and treatment of dehydration with intravenous fluids is the customary treatment. Women with severe hyperemesis gravidarum need frequent medical visits for management of dehydration and electrolyte abnormalities. In some cases, hospitalization is required. Antithyroid drugs (ATDs) are not indicated because the serum T4 returns to normal by 14–18 weeks gestation and ATD use in early pregnancy increases risk of birth defects. Importantly, obstetrical outcome was not

improved in isolated cases in which gestational transient thyrotoxicosis was treated with ATDs (342). However, there are no studies reported in the literature comparing ATD therapy versus supportive therapy. In situations in which symptomatic therapy is indicated, small amounts of β -blockers given over a limited time period may be useful, and close follow-up with repeat investigation for the cause of disease should be performed.

■ RECOMMENDATION 42

The appropriate management of abnormal maternal thyroid tests attributable to gestational transient thyrotoxicosis and/or hyperemesis gravidarum includes supportive therapy, management of dehydration, and hospitalization if needed. ATDs are not recommended, though β -blockers may be considered.

Strong recommendation, moderate-quality evidence.

Question 53: How should women with GD seeking future pregnancy be counseled?

The planning of therapy in relation to possible future pregnancy should be discussed with all women of child-bearing age who develop thyrotoxicosis. In general, pregnancy should be postponed until a stable euthyroid state is reached. As a guide, two sets of thyroid function test within the reference range, at least 1 month apart, and with no change in therapy between tests, can be used to define a stable euthyroid state. The use of contraception until the disease is controlled is strongly recommended.

A hyperthyroid patient who desires future pregnancy may be offered ablative therapy using ^{131}I , thyroid surgery, or medical therapy. Each therapeutic option carries advantages and disadvantages, as detailed below (see Table 8). This is further discussed in the ATA guidelines for the management of thyrotoxicosis (343).

TABLE 8. ADVANTAGES AND DISADVANTAGES OF THERAPEUTIC OPTIONS FOR WOMEN WITH GRAVES' DISEASE SEEKING FUTURE PREGNANCY

Therapy	Advantages	Disadvantages
Antithyroid drugs	Effective treatment to euthyroid state within 1–2 months Often induces gradual remission of autoimmunity (decreasing antibody titers)	Medication adverse effects (mild 5%–8%; severe 0.2%) Birth defects associated with use during pregnancy (MMI 3%–4%; PTU 2%–3% though less severe) Relapse after drug withdrawal likely in 50%–70%
Radioactive iodine	Easily discontinued or modified. Treatment easy to take. Relatively inexpensive Easy oral administration Reduction in goiter size	Repeat therapy at times necessary Rising antibody titers following treatment may contribute to worsening orbitopathy or fetal risk Lifelong need of levothyroxine therapy following ablation
Thyroidectomy	Definitive therapy of hyperthyroidism. Stable euthyroid state easily achieved on replacement levothyroxine therapy Post surgery, gradual remission of autoimmunity occurs Goiter disappears	Life-long need for levothyroxine supplementation Surgical complications occur in 2%–5% Healing and recovery from surgery Permanent neck scar

MMI, methimazole; PTU, propylthiouracil.

Ablative therapy. If the patient opts for radioactive iodine ablative therapy prior to pregnancy, the following recommendations should be provided. First, TRAb levels tend to increase following ^{131}I therapy and may remain elevated for many months following ^{131}I therapy. Therefore, patients with high TRAb levels or severe hyperthyroidism may favor consideration of other therapeutic options such as surgery (344). Second, a subset of young patients with severe GD may not become stably euthyroid within the first year after ^{131}I therapy (345,346,347,348). Third, if ^{131}I therapy is planned, a pregnancy test should be performed 48 hours before ^{131}I ablation to confirm absence of unexpected pregnancy. Fourth, conception should be delayed 6 months and until a stable euthyroid state is reached after ablation and initiation of LT4 replacement therapy.

Antithyroid drugs. If the patient chooses ATD therapy, the following recommendations should be given. First, the increased risk of birth defects associated with both PTU and MMI use during early pregnancy should be reviewed. Second, if possible, ATDs should be avoided in the first trimester of pregnancy, but when necessary PTU is generally favored. Third, consideration can be given to discontinuing PTU after the first trimester and switching to MMI to decrease the risk of liver failure in the mother.

■ RECOMMENDATION 43

In all women of childbearing age who are thyrotoxic, the possibility of future pregnancy should be discussed. Women with GD seeking future pregnancy should be counseled regarding the complexity of disease management during future gestation, including the association of birth defects with ATD use. Preconception counseling should review the risks and benefits of all treatment options and the patient's desired timeline to conception.

Strong recommendation, high-quality evidence.

■ RECOMMENDATION 44

Thyrotoxic women should be rendered stably euthyroid before attempting pregnancy. Several treatment options exist, each of which are associated with risks and benefits. These include ^{131}I ablation, surgical thyroidectomy, or ATD therapy.

Strong recommendation, moderate-quality evidence.

Question 54: What is the management of patients with Graves' hyperthyroidism during pregnancy?

Several studies have shown that obstetric and medical complications are directly related to control of maternal hyperthyroidism, and the duration of the euthyroid state throughout pregnancy (339,342,344,349). Poor control of thyrotoxicosis is associated with pregnancy loss, pregnancy-induced hypertension, prematurity, low birth weight, intrauterine growth restriction, stillbirth, thyroid storm, and maternal congestive heart failure (350). Moreover, some studies suggest fetal exposure to excessive levels of maternal thyroid hormone may program the offspring to develop diseases such as seizure disorders and neurobehavioral disorders in later life (351).

Thionamide ATDs (MMI, carbimazole [CM], and PTU) are the mainstays of treatment for hyperthyroidism during pregnancy (352,353). They reduce iodine organification and coupling of monoiodotyrosine and diiodotyrosine, therefore inhibiting thyroid hormone synthesis. Because the block is not absolute and the thyroid contains a depot of thyroid hormone bound to Tg, the normalization of thyroid function tests takes place gradually over weeks. The initial dose of ATD depends on the severity of the symptoms and the degree of hyperthyroxinemia. In general, initial doses of ATDs during pregnancy are as follows: MMI, 5–30 mg/d (typical dose in average patient 10–20 mg); CM, 10–40 mg/d; and PTU, 100–600 mg/d (typical PTU dose in average patient 200–400 mg/d). The equivalent potency of MMI to PTU is approximately 1:20 (e.g., 5 mg MMI = 100 mg of PTU) (354–356). Ten milligrams of CM is rapidly metabolized to approximately 6 mg of MMI (328). Because the half-life of PTU is shorter than that of MMI, PTU dosing should generally be split into two or three daily doses. In comparison, MMI can generally be given in one daily dose. In rare cases of severe hyperthyroidism, twice or three-times daily dosing may be of benefit (357,358).

Importantly, side effects occur in 3%–5% of patients taking thionamide drugs, the majority of which are allergic reactions such as skin rash (352), whereas the severe side effects of agranulocytosis (0.15%) (359,360) and liver failure (<0.1%) (92) are rare. Most side effects develop within the first months following initiation (359) or re-initiation (361) of therapy. In 2010 the U.S. Food and Drug Administration (FDA) called attention to the risk of hepatotoxicity in patients exposed to PTU because PTU had been found to be third on the list of drugs leading to liver transplantation in the United States (362,363). An advisory committee recommended limiting the use of PTU to the first trimester of pregnancy (364). Exceptions to this are patients with MMI allergy or those with thyroid storm. Monitoring hepatic enzymes during administration of PTU may be considered. However, no prospective data exist that have demonstrated that the monitoring of liver enzymes is effective in preventing fulminant PTU-induced hepatotoxicity.

However, the greatest risk surrounding the use of ATDs in pregnancy is related to their potential teratogenic effects (365). As early as 1972, exposure to MMI had been associated with aplasia cutis (366). Subsequently, several other types of congenital malformations have been associated with MMI use. A “syndrome of methimazole/carbimazole embryopathy” was described, which also includes dysmorphic facies (367, 368). Apart from aplasia cutis, defects with a statistically significant association with the use of MMI include choanal or esophageal atresia; various types of abdominal wall defects including umbilicocele; and eye, urinary system, and ventricular septal defects (369,370,371). Recent studies have shown that these complications are more common than previously thought, and they affect 2%–4% of children who have been exposed to MMI in early pregnancy, especially during gestational weeks 6–10 (369,370,372).

PTU was previously considered a safe medication for use during gestation (364,367). Recently, however, a Danish study revealed that 2%–3% of children exposed to PTU developed birth defects associated with this therapy (370). The defects were primarily face and neck cysts (often considered to be minor birth defects) and urinary tract abnormalities (in males). Prior to the recent investigation, such abnormalities

were not commonly associated with PTU exposure, likely because they were diagnosed later in life when complications ensued. Importantly, however, most affected patients received surgery for the abnormality (373). Thus, PTU-associated birth defects appear less severe than MMI-associated birth defects but occur with similar incidence.

Beta-adrenergic blocking agents, such as propranolol 10–40 mg every 6–8 hours may be used for controlling hypermetabolic symptoms until patients have become euthyroid on ATD therapy. The dose should be reduced as clinically indicated. In the vast majority of cases the drug can be discontinued in 2–6 weeks. Long-term treatment with β -blockers has been associated with intrauterine growth restriction, fetal bradycardia, and neonatal hypoglycemia (374). One study suggested a higher rate of spontaneous pregnancy loss when both drugs were taken together, as compared with patients receiving only MMI (375). However, it was not clear that this difference was attributable to the medication as opposed to the underlying condition. Beta-blocking drugs may be used as preparation for thyroidectomy.

Question 55: Should antithyroid medication be withdrawn or modified in early pregnancy?

Many patients receiving ATD therapy for GD gradually enter remission of the autoimmune abnormality when made euthyroid. Though approximately half of patients eventually experience a relapse of hyperthyroidism when the antithyroid medication is withdrawn after 1–2 years of therapy, only a small fraction of patients who have become TRAb negative during therapy will become hyperthyroid within the first months. In support of this, in a prospective Norwegian study of 218 GD patients treated with ATD for 12 months, only 5% of TRAb-negative patients became hyperthyroid within 8 weeks after ATD withdrawal (376).

Thus, one option when pregnancy is diagnosed in a woman receiving ATD therapy for GD and who, based on clinical and biochemical findings appears to be in remission, is to withdraw ATD medication and perform repeated thyroid function testing during the first trimester of pregnancy. If ATD therapy is needed during the first trimester, PTU is preferred over MMI because the risk for severe birth defects is lower. Cessation of medication has to be recommended early in gestation, before the major teratogenic periods (gestational weeks 6–10) (372,373,376,377).

Gestational weeks are calculated from the first day of the last normal menstrual period; it is typically at week 5 when the next normal menstruation does not appear in a pregnant woman. Sensitive pregnancy tests are widely available and should detect pregnancy by this time. Optimally, women receiving ATD should test for pregnancy within the first days of missing or unusually weak menstruation. If the test is positive, the woman should contact the caregiver responsible for thyroid therapy to receive instruction regarding withdrawing or modifying ATD therapy and to discuss thyroid function testing.

The risk of rapid relapse of hyperthyroidism after medication withdrawal in early pregnancy varies among patients. The risk is high in patients who have been treated for a short period (<6 months), who have suppressed or low serum TSH while on medication prepregnancy, who require >5–10 mg of MMI per day to stay euthyroid, who have active orbitopathy or large goiter, and those who have high levels of TRAb (378). Because

this assessment is based upon clinical judgment, no specific single parameter may be used to assign risk. However, if the risk is considered high, medication should not be withdrawn, and PTU should be administered as the drug of choice.

Even if conventional ATDs are effective in achieving a euthyroid state, the risk of birth defects has raised the question of alternative types of drug therapy for hyperthyroidism. Iodine in pharmacological doses was widely used to treat hyperthyroidism before the thionamide drugs (and radioiodine) became available. It was (and is) especially effective as a preparation for subsequent thyroid surgery, as introduced by Plummer in 1923 (379). However, escape from the effect during prolonged therapy was not uncommon, especially in patients with severe hyperthyroidism (380), and iodine therapy is now mostly used to reduce thyroid blood flow before surgery and as part of the combination of therapies given to patients with thyrotoxic crises.

An exception to this narrow indication for the treatment of hyperthyroidism is Japan, where there is considerable experience with high-dose iodine therapy in pregnancy (381,382). The doses of iodine used have varied between 5 and 75 mg/d, adjusted according to thyroid function tests. In a recent retrospective nonrandomized study, substituting MMI with potassium iodine in early pregnancy reduced the risk of birth defects. However, a proportion of such pregnant women became hyperthyroid following the change in treatment (382). Important to note, Japan is a high iodine intake country, and the Japanese results may not be reproducible in other countries. Cases of iodine-induced congenital hypothyroidism have been reported in children of U.S. women treated with high-dose iodine during pregnancy (383), and the use of iodine-containing disinfectants during labor is a well-established cause of elevated TSH during screening for congenital hypothyroidism (384). Thus, at present, such therapy cannot be recommended outside Japan until more evidence on safety and efficacy is available.

Theoretically, other possible treatments can also be considered. Cholestyramine is not absorbed from the gut (and thus not transferred to the fetus) and may moderately reduce circulating thyroid hormones by binding the hormones during their enterohepatic circulation (385–387). Cholestyramine has been used in pregnancy to treat obstetric cholestasis (388). Gastrointestinal discomfort with nausea, vomiting, and diarrhea may be a problem (388), and there is a risk of binding in the gut of vitamins and other substances important in pregnancy. Perchlorate competitively inhibits the active transport of iodide into the thyroid (and into breast milk), and it has been used to treat hyperthyroidism (389,390). Limited evidence suggests it has no teratogenic effects as long as normal maternal thyroid function is maintained (391). In summary, more data are needed before the use of any such drugs can be recommended. Lithium has thyroid-inhibiting effects somewhat similar to pharmacological doses of iodine (392), but it may be teratogenic (393) and therefore should not be used to treat hyperthyroidism in pregnancy.

■ RECOMMENDATION 45

Women taking MMI or PTU should be instructed to confirm potential pregnancy as soon as possible. If the pregnancy test is positive, pregnant women should contact their caregiver immediately.

Strong recommendation, high-quality evidence.

■ RECOMMENDATION 46

- (a) In a newly pregnant woman with GD, who is euthyroid on a low dose of MMI (≤ 5 – 10 mg/d) or PTU (≤ 100 – 200 mg/d), the physician should consider discontinuing all antithyroid medication given potential teratogenic effects. The decision to stop medication should take into account the disease history, goiter size, duration of therapy, results of recent thyroid function tests, TRAb measurement, and other clinical factors.

Weak recommendation, low-quality evidence.

- (b) Following cessation of antithyroid medication, maternal thyroid function testing (TSH, and FT4 or TT4) and clinical examination should be performed every 1–2 weeks to assess maternal and fetal thyroid status. If the pregnant woman remains clinically and biochemically euthyroid, test intervals may be extended to 2–4 weeks during the second and third trimester.

Weak recommendation, low-quality evidence.

- (c) At each assessment, the decision to continue conservative management (withholding antithyroid medication) should be guided both by the clinical and the biochemical assessment of maternal thyroid status.

Weak recommendation, low-quality evidence.

■ RECOMMENDATION 47

In pregnant women with a high risk of developing thyrotoxicosis if antithyroid drugs were to be discontinued, continued antithyroid medication may be necessary. Factors predicting high clinical risk include being currently hyperthyroid, or requirement of >5 – 10 mg/d MMI or >100 – 200 mg/d PTU to maintain a euthyroid state. In such cases,

- (a) PTU is recommended for the treatment of maternal hyperthyroidism through 16 weeks of pregnancy.

Strong recommendation, moderate-quality evidence.

- (b) Pregnant women receiving MMI who are in need of continuing therapy during pregnancy should be switched to PTU as early as possible.

Weak recommendation, low-quality evidence.

- (c) When shifting from MMI to PTU, a dose ratio of approximately 1:20 should be used (e.g., MMI 5 mg/d = PTU 50 mg twice daily).

Strong recommendation, moderate-quality evidence.

- (d) If ATD therapy is required after 16 weeks gestation, it remains unclear whether PTU should be continued or therapy changed to MMI. As both medications are associated with potential adverse effects and shifting potentially may lead to a period of less-tight control, no recommendation regarding switching antithyroid drug medication can be made at this time.

No recommendation, insufficient evidence.

Question 56: What are the principles of thyroid testing and ATD administration when treating Graves' hyperthyroidism during pregnancy?

Thyroid stimulating antibodies, ATDs, and most maternal thyroid hormones can effectively cross the placenta barrier. When the fetal thyroid is functional, it can respond to TRAb antibodies, causing excess fetal production of thyroid hormone. Furthermore, if the mother has an intact thyroid and is hyperthyroid from GD, the fetus will also be exposed to the

hyperthyroxinemia produced by the mother's thyroid during gestation. Typically, fetal hyperthyroidism due to cross-placental passage of TRAb develops at or after week 20 of pregnancy. However, one case has been published in which excessive maternal TRAb production led to fetal hyperthyroidism by week 18 (394).

As mentioned, MMI, PTU, and CM also effectively cross the placenta, and therefore ATD therapy for maternal hyperthyroidism also modulates fetal thyroid function. Importantly, all ATDs tend to be more potent in the fetus than in the mother. Thus, when the mother is made euthyroid, the fetus is often overtreated (395). Therefore, in order to avoid a deleterious fetal impact, the aim of treatment is to maintain maternal TT4/FT4 values at, or just above the pregnancy-specific upper limit of normal. As a general rule, the smallest possible dose of ATDs should be used whenever possible.

In the setting of hyperthyroidism during pregnancy, maternal TT4/FT4 and TSH (and in cases of severe hyperthyroidism, also serum T3) should be measured approximately every 2–4 weeks following initiation of therapy, and every 4–6 weeks after achieving the target value (396–398). When trimester-specific FT4 values are not available, use of the reference range for nonpregnant patients is recommended. Separately, a TT4 measurement with reference value 1.5 times the nonpregnancy range may be used in second and third trimesters. Overtreatment should be avoided because of the possibility of inducing fetal goiter and/or fetal hypothyroidism (399).

In the first trimester of pregnancy some women with GD experience an exacerbation of symptoms (400), which is parallel to the moderate increase in incidence of GD in early pregnancy (401). By the third trimester the incidence of GD becomes very low (401) corresponding to the general decrease in thyroid autoimmunity with a decrease in TRAb.

Discontinuation of all ATD therapy is feasible in 20%–30% of patients in the last trimester of gestation (402). Maternal serum TSH well within the reference range is a sign that the ATD dose has to be reduced to avoid fetal overtreatment. If this is not done, fetal hypothyroidism and goiter may develop from overtreatment with ATDs. Disappearance of maternal TRAb in late pregnancy indicates a high likelihood of successful ATD withdrawal.

■ RECOMMENDATION 48

- (a) In women being treated with ATDs in pregnancy, FT4/TT4 and TSH should be monitored approximately every 4 weeks.

Strong recommendation, moderate-quality evidence.

- (b) Antithyroid medication during pregnancy should be administered at the lowest effective dose of MMI or PTU, targeting maternal serum FT4/TT4 at the upper limit or moderately above the reference range.

Strong recommendation, high-quality evidence.

Because the fetal thyroid responds more strongly to ATD therapy than the maternal thyroid, mothers on an ATD in the second half of pregnancy, who by nonpregnancy standards would be considered euthyroid, should have the ATD dose reduced to protect the fetus.

In occasional patients, GD stays very active and serum TT3 may remain elevated even if TT4/FT4 becomes normal

or even low (403). An increase in ATD dose to normalize maternal serum TT3 will cause elevated serum TSH in the infants at birth (398), and a balance in ATD dosing with careful clinical evaluation of the fetus and the mother is needed.

Worsening of disease activity with a need for an increase in ATD dose or relapse of previously remitted disease often occurs after delivery (400,404). Women should be informed about this risk, and appropriate monitoring performed. A single Japanese study has suggested that relapse may be prevented by low-dose ATD during the postpartum period (405), but more studies on this are needed.

The combination of LT4 and an ATD (block-replace therapy) has in general been shown not to improve GD remission rates (406), and it results in a larger dose of ATD required in order to maintain the FT4 within the target range. The placenta is readily permeable to the ATD but not to the LT4 given to the mother, and the fetal thyroid is relatively more sensitive to the effect of ATDs than the maternal thyroid. Therefore, block-replacement therapy given to the mother in the second half of pregnancy will generally lead to fetal goiter and hypothyroidism (403). The only indication for such combination therapy during pregnancy is in the treatment of isolated fetal hyperthyroidism caused by maternal TRAb production in a mother who previously received ablative therapy for GD (407). The ATD will pass the placenta and treat the fetal hyperthyroidism, whereas the LT4 is necessary to keep the mother euthyroid.

■ RECOMMENDATION 49

A combination regimen of LT4 and an ATD should not be used in pregnancy, except in the rare situation of isolated fetal hyperthyroidism.

Strong recommendation, high-quality evidence.

Question 57: What are the indications and timing for thyroidectomy in the management of GD during pregnancy?

Thyroidectomy should be considered in cases of allergies/contraindications to both ATDs, in the patient who is not compliant with drug therapy, and in women in whom euthyroidism cannot be achieved even on large doses of ATDs. If surgery is indicated, the second trimester is the optimal time. Thyroidectomy is often followed by a gradual, but not immediate disappearance of TRAb, and withdrawal of ATD in the mother after thyroidectomy may lead to isolated fetal hyperthyroidism (403). High serum TRAb values before surgery indicate a risk for isolated fetal hyperthyroidism, and after maternal ATD withdrawal a program of careful fetal monitoring and possible therapy should be planned (403). In hyperthyroid patients not tolerating or nonresponsive to ATD therapy, preparation for surgery with β -blocking agents and a short course of potassium iodide solution (50–100 mg/d) are recommended (381). Potassium iodide preparation before surgery is also recommended in patients with other signs of active disease to reduce bleeding during surgery.

■ RECOMMENDATION 50

Thyroidectomy in pregnancy may be indicated for unique scenarios. If required, the optimal time for thyroidectomy

is in the second trimester of pregnancy. If maternal TRAb concentration is high (>3 times the upper reference for the assay) the fetus should be carefully monitored for development of fetal hyperthyroidism throughout pregnancy, even if the mother is euthyroid post thyroidectomy.

Strong recommendation, high-quality evidence.

Question 58: How should pregnant patients with GD be prepared for urgent nonthyroid surgery?

■ RECOMMENDATION 51

We concur with the American College of Obstetricians and Gynecologists' Committee on Obstetric Practice consensus guidelines (written in 2011 and revised in 2015) (408) which state the following: "1) A pregnant woman should never be denied indicated surgery, regardless of trimester. 2) Elective surgery should be postponed until after delivery. 3) If possible, nonurgent surgery should be performed in the second trimester when preterm contractions and spontaneous abortion are least likely."

In the setting of a patient with GD undergoing urgent, nonthyroid surgery, if the patient is well controlled on ATD, no other preparation is needed. Beta-blockade should also be utilized if needed.

Strong recommendation, moderate-quality evidence.

Question 59: What is the value of TRAb measurement in the evaluation of a pregnant woman with Graves' hyperthyroidism?

Fetal risks in women with previous or current Graves' hyperthyroidism include (a) fetal hyperthyroidism, (b) neonatal hyperthyroidism, (c) fetal hypothyroidism, (d) neonatal hypothyroidism, and (e) central hypothyroidism. The above potential complications depend on several factors: (a) poor control of hyperthyroidism throughout pregnancy may induce transient central hypothyroidism (409,410); (b) excessive amounts of ATDs may be responsible for fetal and neonatal hypothyroidism, even if the mother is biochemically euthyroid (399,403); and (c) high levels of thyroid stimulating antibodies in the second half of pregnancy may induce fetal and neonatal hyperthyroidism (411,412,413,414). TRAb is measurable in around 95% of patients with active Graves' hyperthyroidism, and levels may remain high following ablation therapy, even more so after radioiodine treatment than surgical removal (415).

Indications for ordering a TRAb test in pregnant women with GD include (a) mothers with untreated or ATD-treated hyperthyroidism in pregnancy, (b) a previous history of GD with past treatment with radioiodine or total thyroidectomy, (c) a previous history of delivering an infant with hyperthyroidism, or (d) a known history of thyroidectomy for the treatment of hyperthyroidism in pregnancy (416). In the majority of patients, maternal TRAb concentrations decrease with the progression of pregnancy; however, as in nonpregnant patients, the course of GD is variable.

The incidence of fetal and neonatal hyperthyroidism is between 1% and 5% in all women with active or a past history of Graves' hyperthyroidism and is associated with increased fetal/neonatal morbidity and mortality if

unrecognized and untreated (417). In a recent French study (418), follow-up of 47 newborns of mothers harboring measurable levels of TRAb in pregnancy showed that nine developed transient neonatal hyperthyroidism and five of these (9% overall) had to be treated with ATD for a median duration of 60 days. A maternal TRAb serum concentration >5 IU/L (approximately 3 times the upper limit of normal for the assay) in the second and third trimester predicted neonatal hyperthyroidism with 100% sensitivity and 43% specificity. A similar TRAb level risk cutoff was found in a separate French study (419).

A determination of serum TRAb in early pregnancy is helpful in detecting pregnancies at risk. A value >5 IU/L or 3 times the upper limit of normal in a mother who previously received ablative therapy for GD is an indication for establishing close follow-up of the fetus in collaboration with a maternal–fetal medicine physician. Cases of overlooked isolated fetal hyperthyroidism leading to fetal loss in such women have repeatedly been published.

A determination of serum TRAb in late pregnancy in a mother who is still in need of ATD therapy to remain euthyroid, is helpful for detecting risk for neonatal hyperthyroidism. A value >5 IU/L or 3 times the upper limit of normal in the mother indicates that the fetal thyroid may be strongly stimulated by TRAb passing through the placenta. After birth, any ATD from the mother is much more rapidly cleared in the neonate than are the TRAb, and the neonate may become hyperthyroid.

If TRAb becomes undetectable in a pregnant woman taking ATD, it may be feasible to reduce or withdraw the ATD to protect the fetus against hypothyroidism and goiter.

In women who are in remission and euthyroid after a previous course of ATD therapy for GD, measurement of TRAb in pregnancy is not required (416).

■ RECOMMENDATION 52

- (a) If the patient has a past history of GD treated with ablation (radioiodine or surgery), a maternal serum determination of TRAb is recommended at initial thyroid function testing during early pregnancy.

Strong recommendation, moderate-quality evidence.

- (b) If maternal TRAb concentration is elevated in early pregnancy, repeat testing should occur at weeks 18–22.

Strong recommendation, moderate-quality evidence.

- (c) If maternal TRAb is undetectable or low in early pregnancy, no further TRAb testing is needed.

Weak recommendation, moderate-quality evidence.

- (d) If a patient is taking ATDs for treatment of Graves' hyperthyroidism when pregnancy is confirmed, a maternal serum determination of TRAb is recommended.

Weak recommendation, moderate-quality evidence.

- (e) If the patient requires treatment with ATDs for GD through midpregnancy, a repeat determination of TRAb is again recommended at weeks 18–22.

Strong recommendation, moderate-quality evidence.

- (f) If elevated TRAb is detected at weeks 18–22 or the mother is taking ATD in the third trimester, a TRAb measurement should again be performed in late pregnancy (weeks 30–34) to evaluate the need for neonatal and postnatal monitoring.

Strong recommendation, high-quality evidence.

Question 60: Under what circumstances should additional fetal ultrasound monitoring for growth, heart rate, and goiter be performed in women with Graves' hyperthyroidism in pregnancy?

Serial ultrasound examinations may be performed for the assessment of gestational age, fetal viability, amniotic fluid volume, fetal anatomy, and detection of malformations. Fetal well-being may be compromised in the presence of elevated TRAb, uncontrolled hyperthyroidism, and pre-eclampsia (349,420–422). Signs of potential fetal hyperthyroidism that may be detected by ultrasonography include fetal tachycardia (heart rate >170 bpm, persistent for over 10 minutes), intrauterine growth restriction, presence of fetal goiter (the earliest sonographic sign of fetal thyroid dysfunction), accelerated bone maturation, signs of congestive heart failure, and fetal hydrops (411,421–423). A team approach to the management of these patients is required, including an experienced obstetrician or maternal–fetal medicine specialist, neonatologist, and anesthesiologist. In most cases, the diagnosis of fetal hyperthyroidism should be made on clinical grounds based on maternal history, interpretation of serum TRAb levels, and fetal ultrasonography (396,411,422,423).

■ RECOMMENDATION 53

Fetal surveillance should be performed in women who have uncontrolled hyperthyroidism in the second half of pregnancy, and in women with high TRAb levels detected at any time during pregnancy (greater than 3 times the upper limit of normal). A consultation with an experienced obstetrician or maternal–fetal medicine specialist is recommended. Monitoring may include ultrasound to assess heart rate, growth, amniotic fluid volume, and the presence of fetal goiter.

Strong recommendation, moderate-quality evidence.

Question 61: When should umbilical blood sampling be considered in women with GD in pregnancy?

Umbilical cord blood sampling (cordocentesis) is associated with both fetal mortality and morbidity (424,425). It has been utilized when a mother is TRAb positive and treated with ATDs, a fetal goiter is present, and the thyroid status of the fetus is unclear (403,411,426). The presence of elevated TRAb alone is not an indication for cordocentesis (427).

■ RECOMMENDATION 54

Cordocentesis should be used in rare circumstances and performed in an appropriate setting. It may occasionally be of use when fetal goiter is detected in women taking ATDs to help determine whether the fetus is hyperthyroid or hypothyroid.

Weak recommendation, low-quality evidence.

Recently, biological assays able to distinguish between stimulating and blocking TRAb have become commercially available (428). The use of such assays for characterization of TRAb biological activity may theoretically make cordocentesis unnecessary, though the complexity of differential clearance of stimulatory and blocking antibodies should be considered (see Section X, Fetal and Neonatal Considerations).

Question 62: How should hyperthyroidism caused by autonomous thyroid nodules be handled in pregnancy?

One or more autonomous thyroid nodules is a common cause of hyperthyroidism in populations with current or previous mild to moderate iodine deficiency (429). However, even in such areas this type of hyperthyroidism is quite rare under the age of 40 years (430). Hyperthyroidism caused by autonomous nodules tends to develop more insidiously and be less severe than in GD.

In pregnancy, a major difference from GD is that no TRAb is produced by the mother, and consequently the fetal thyroid is not stimulated as it is in GD. Therefore, ATD therapy to make the mother euthyroid would significantly increase the risk of hypothyroidism and goiter in the fetus. Given the limited data on this subject, if ATDs are required in this setting, careful monitoring of the fetus should occur. Furthermore, the dose of ATD should be kept low, and maternal surgical therapy with removal of autonomous nodule(s) considered if signs of fetal hypothyroidism develop. Generally, if possible, ablative therapy should be considered before conception for hyperthyroidism caused by thyroid autonomy in women seeking future pregnancy.

If only low-grade autonomous thyroid hormone production with subclinical hyperthyroidism is present in a non-pregnant woman, the physiological increase in thyroid hormone needs may ameliorate any hyperthyroidism in pregnancy. On the other hand, the early pregnancy high hCG level may theoretically activate nonaffected normal thyroid tissue and increase thyroid secretion in early pregnancy. Furthermore, hormone production in autonomous thyroid nodules will depend on the available amount of iodine. Therefore, women with such disease should probably refrain from taking iodine-containing supplements in pregnancy.

■ RECOMMENDATION 55

If ATD therapy is given for hyperthyroidism caused by autonomous nodules, the fetus should be carefully monitored for goiter and signs of hypothyroidism during the second half of pregnancy. A low dose of ATD should be administered with the goal of maternal FT₄ or TT₄ concentration at the upper limit or moderately above the reference range.

Strong recommendation, low-quality evidence.

Question 63: What are the etiologies of thyrotoxicosis in the postpartum period?

The most common cause of thyrotoxicosis in the postpartum period is postpartum thyroiditis (PPT), with a thyrotoxicosis prevalence of 4% (431,432) (see Section XII). Many cases are mild and of short duration and spontaneously revert to euthyroidism. More severe cases in need of a short course of β -blockers are seen in patients with high levels of TPOAb, and they are often followed by a period of hypothyroidism. The majority of women return to euthyroidism by 1 year post partum (433,434).

The postpartum increase in thyroid autoimmunity is also associated with a 3- to 4-fold increase in the incidence of new GD (401), in addition to the increase in the risk of relapse of GD in remission after previous ATD therapy. In one study, the overall relapse rate of GD following a pregnancy was

84%, as compared to a relapse rate of 56% in women who did not become pregnant (404).

Question 64: How should Graves' hyperthyroidism be treated in lactating women?

See Section XI.

IX. THYROID NODULES AND THYROID CANCER DURING PREGNANCY

Thyroid nodules and thyroid cancer discovered during pregnancy present unique challenges to both the clinician and the mother. A careful balance is required between making a definitive diagnosis and instituting treatment while avoiding interventions that may adversely impact the mother, the health of the fetus, or the maintenance of the pregnancy.

Question 65: What is the prevalence of thyroid nodules during pregnancy?

Three studies have evaluated the prevalence of thyroid nodules during pregnancy, the impact of pregnancy on nodular size, and the proportion of women who have new nodules detected during pregnancy. All three studies were performed in areas with mild to moderate iodine deficiency (Brussels, China, and Germany), with the majority using ultrasound examination of the gland. The prevalence of thyroid nodules varied between 3% and 21% (187,435,436) and increased with increasing parity (9.4% without a prior pregnancy, 20.7% with one prior pregnancy, 20.7% with two prior pregnancies, and 33.9% with three or more prior pregnancies) (435). In the Belgian study, 60% of nodules doubled in size during pregnancy, yet remained between 5 and 12 mm (187). The maximum diameter of the dominant nodule did not increase during pregnancy in the Chinese study (mean diameter was 5.1 mm, 5.1 mm, and 5.5 mm, in the first, second, and third trimester, respectively). However, an increase in nodular volume was reported during pregnancy, with a return to first-trimester volumes by the third postpartum month (436). The studies in Belgium and Germany reported that 11%–20% of women with a nodule detected in the first trimester of pregnancy developed a second nodule through the course of pregnancy (187,436). Increasing age is associated with an increase in the proportion of pregnant women who have thyroid nodules (187,436).

Question 66: What is the prevalence of thyroid cancer in women with thyroid nodules discovered during pregnancy?

Data regarding the prevalence of thyroid cancer derive from three retrospective studies performed at three tertiary referral centers (Mayo Clinic, George Washington University Hospital, and Mount Sinai Hospital-Toronto), one prospective study, and a retrospective study of the California Cancer Registry. The research performed at the referral centers revealed a 15% (6/40, Mayo Clinic) (437), 12% (7/57, George Washington Hospital) (438), and 43% (7/16, Mount Sinai Hospital-Toronto) (439) rate of thyroid malignancy. All three studies are limited by two major methodological flaws. The first, and most problematic, is selection bias. The population studies consisted of women referred for diagnosis and treatment at major referral centers. As such, they are not representative of the population of pregnant women with

thyroid nodules detected during pregnancy. Instead, they represent a select group of women referred to a tertiary medical center because of physician concern for thyroid malignancy. Consequently, they likely represent an overrepresentation of the prevalence of thyroid malignancy during pregnancy. Secondly, each study was retrospective in nature and therefore neither the accuracy of the diagnosis nor the completeness of case identification within the database could be verified. The lone prospective study investigated the prevalence of thyroid cancer during pregnancy in 212 Chinese women. The study found a 15.3% (34/221) rate of thyroid nodules and a 0% rate of malignancy. Interpreting these data is hampered by the limited number of women enrolled in the study (436). The final study consisted of a population-based retrospective analysis of all obstetrical deliveries in California from 1991 through 1999 identified by cross-referencing maternal/neonatal hospital discharges in California and the California Cancer Registry ($n=4,846,505$ women). A prevalence of thyroid cancer in pregnancy of 14.4/100,000 was reported, with papillary cancer being the most frequent pathological type (440). Timing of diagnosis of the thyroid malignancy was as follows: 3.3/100,000 cases diagnosed before delivery, 0.3/100,000 at delivery, and 10.8/100,000 within 1 year post partum.

Question 67: What is the optimal diagnostic strategy for thyroid nodules detected during pregnancy?

History and physical examination. The patient with a thyroid nodule should be asked about a family history of benign or malignant thyroid disease, familial medullary thyroid carcinoma, multiple endocrine neoplasia type 2 (MEN 2), familial papillary thyroid carcinoma, and a familial history of a tumor syndrome predisposing to thyroid cancer syndrome (e.g., **phosphatase and tensin homolog [PTEN]** hamartoma tumor syndrome [Cowden's disease], familial

adenomatous polyposis, Carney complex, or Werner syndrome) (441,442,443). The malignancy risk is higher for nodules detected in both adult survivors of childhood cancers in which treatment involved head, neck, and/or cranial radiation and those exposed to ionizing radiation before 18 years of age (444,445). Thorough palpation of the thyroid and neck inspection for cervical nodes is essential (446).

Ultrasound. Thyroid ultrasound is the most accurate tool for detecting thyroid nodules, determining their sonographic features and pattern, monitoring growth, and evaluating cervical lymph nodes. The recent 2015 ATA Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer should be referenced for diagnostic use and performance of thyroid and neck sonography as well as for decision-making regarding fine-needle aspiration (FNA) for thyroid nodules (443). Recent reports have validated that the identification of defined nodule sonographic patterns representing constellations of sonographic features is more robust for malignancy risk correlation than that associated with individual ultrasound characteristics (447,448). Hence, a high-suspicion sonographic pattern that includes solid hypoechoic nodules with irregular borders and microcalcifications correlates with a >70% chance of cancer compared to the very low-suspicion pattern of a noncalcified mixed cystic solid or spongiform nodule (<3% cancer risk). Recommendation 9 in the 2015 ATA guidelines (443) indicates different FNA size cutoffs based upon five defined sonographic patterns and their associated risk stratification for thyroid cancer (Table 9).

Thyroid function tests. All women with a thyroid nodule should have a TSH measurement performed (449,450). Thyroid function tests are usually normal in women with thyroid cancer. In nonpregnant women, a subnormal serum TSH level may indicate a functioning nodule, which is then evaluated with scintigraphy because functioning nodules are

TABLE 9. ATA SONOGRAPHIC PATTERNS AND ESTIMATED RISK OF MALIGNANCY FOR THYROID NODULES (ADAPTED FROM ATA 2015 GUIDELINES WITH PERMISSION)

<i>Sonographic pattern</i>	<i>Ultrasound features</i>	<i>Estimated risk of malignancy</i>	<i>FNA size cutoff (largest dimension)</i>
High suspicion	Solid hypoechoic nodule or solid hypoechoic component of a partially cystic nodule with one or more of the following features: irregular margins (infiltrative, microlobulated), microcalcifications, taller than wide shape, rim calcifications with small extrusive soft tissue component, evidence of extrathyroidal extension	>70%–90%	Recommend at >1 cm
Intermediate suspicion	Hypoechoic solid nodule with smooth margins without microcalcifications, extrathyroidal extension, or taller than wide shape	10%–20%	Recommend at >1 cm
Low suspicion	Isoechoic or hyperechoic solid nodule, or partially cystic nodule with eccentric solid areas, without microcalcification, irregular margin or extrathyroidal extension, or taller than wide shape.	5%–10%	Recommend at >1.5 cm
Very low suspicion	Spongiform or partially cystic nodules without any of the sonographic features described in low, intermediate or high suspicion patterns	<3%	Consider at >2 cm; observation without FNA is also a reasonable option
Benign	Purely cystic nodules (no solid component)	< 1%	No biopsy

Ultrasound-guided fine-needle aspiration (FNA) is recommended for lateral neck cervical lymph nodes that are sonographically suspicious for thyroid cancer.

so rarely malignant that cytologic evaluation is not indicated. However, pregnancy produces two hurdles for following this algorithm. First, the lower limit of the TSH reference range decreases, especially during early gestation, making it difficult to differentiate what is normal for pregnancy from potential nodular autonomous function. Second, scintigraphy with either technetium pertechnetate or ^{123}I is contraindicated in pregnancy (see below).

Calcitonin and thyroglobulin. As within the general population, the routine measurement of calcitonin remains controversial (451). Calcitonin measurement may be performed in pregnant women with a family history of medullary thyroid carcinoma or MEN 2 or a known *RET* gene mutation. However, the utility of measuring calcitonin in all pregnant women with thyroid nodules has not been evaluated. The pentagastrin stimulation test is contraindicated in pregnancy (452).

In the presence of an *in situ* thyroid gland, serum Tg measurements are neither sensitive nor specific for thyroid cancer and can be elevated in many benign thyroid disorders (453). Thus, serum Tg measurement is not recommended.

Fine-needle aspiration. Fine needle aspiration is a safe diagnostic tool in pregnancy and may be performed in any trimester (454–463). Two retrospective case series of FNAs performed during pregnancy, involving a total of 94 patients, have been published. In those cases in which surgery was performed, pathological examination of the specimens confirmed the diagnosis of all FNAs classified by cytology as either benign or malignant. Six of the 16 (37.5%) cases reported by cytology as suspicious for malignancy were found to be malignant at pathological examination (437,438). Pregnancy does not appear to alter a cytological diagnosis of thyroid tissue obtained by FNA, but there have been no prospective studies to evaluate potential differences in FNA cytology obtained during pregnancy versus in the nonpregnant state. Since overall survival does not differ if surgery is performed during or after gestation in patients diagnosed with thyroid cancer during pregnancy (464–466), patient preference for timing of FNA (during pregnancy or post partum) should be considered.

Radionuclide scanning. Iodine ^{131}I readily crosses the placenta and the fetal thyroid begins to accumulate iodine by 12–13 weeks gestation (467). There are reports of inadvertent administration of therapeutic ^{131}I therapy for treatment of hyperthyroidism during unsuspected pregnancy. If ^{131}I is given after 12–13 weeks gestation, it accumulates in the fetal thyroid resulting in fetal/neonatal hypothyroidism. In this scenario, the International Atomic Energy Agency (IAEA) recommends intervention with 60–130 mg of stable potassium iodide given to the mother only if the pregnancy is discovered within 12 hours of ^{131}I administration. This will partially block the fetal thyroid, hence reducing fetal thyroid ^{131}I uptake (468). However, if maternal treatment occurs prior to 12 weeks, the fetal thyroid does not appear to be damaged (469–471). Rather, the issue is the fetal whole-body radiation dose due to gamma emissions from ^{131}I in the maternal bladder, which is in the range of 50–100 mGy/GBq of administered activity. This dose is decreased by hydrating the mother and by encouraging frequent voiding (468). No studies have specifically examined whether scanning doses of ^{123}I or technetium pertechnetate have adverse fetal effects if used during gesta-

tion. In general, these are contraindicated because all maternal radionuclides are associated with a fetal irradiation resulting from both placental transfer and external irradiation from maternal organs, specifically the bladder. Again, both maternal hydration and frequent voiding reduce fetal exposure (468).

The optimal diagnostic strategy for thyroid nodules detected during pregnancy is based on risk stratification. All women should have the following: a complete history and clinical examination, serum TSH measurement, and ultrasound of the neck.

■ RECOMMENDATION 56

For women with suppressed serum TSH levels that persist beyond 16 weeks gestation, FNA of a clinically relevant thyroid nodule may be deferred until after pregnancy. At that time, if serum TSH remains suppressed, a radionuclide scan to evaluate nodule function can be performed if not breastfeeding.

Strong recommendation, low-quality evidence.

■ RECOMMENDATION 57

The utility of measuring calcitonin in pregnant women with thyroid nodules is unknown. The task force cannot recommend for or against routine measurement of serum calcitonin in pregnant women with thyroid nodules.

No recommendation, insufficient evidence.

■ RECOMMENDATION 58

Thyroid nodule FNA is generally recommended for newly detected nodules in pregnant women with a nonsuppressed TSH. Determination of which nodules require FNA should be based upon the nodule's sonographic pattern as outlined in Table 9. The timing of FNA, whether during gestation or early postpartum, may be influenced by the clinical assessment of cancer risk or by patient preference.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 59

Radionuclide scintigraphy or radioiodine uptake determination should not be performed during pregnancy.

Strong recommendation, high-quality evidence.

Question 68: Does pregnancy impact the prognosis of newly diagnosed thyroid carcinoma?

Six studies have compared the diagnostic features and prognosis of women diagnosed with differentiated thyroid cancer (DTC) either during gestation or within the first year post partum to nonpregnant women (464–466,472–474). None of the studies were RCTs, all were retrospective, and the size of many of the studies was limited. Four studies either predate or do not use the contemporary tools for recurrence detection, including sensitive serum Tg assays and neck sonography (464–466,473). Using large database evaluation (464–466), the analyzed outcomes were either overall survival or disease-free survival, neither of which differed between the two groups. This outcome is not unexpected because the vast majority (>99%) of both the pregnant and nonpregnant populations had stage 1 disease. In addition, all studies consistently found that timing of surgery, either during pregnancy or deferred until postpartum, did not affect survival.

Recently, using both sonography and sensitive Tg assays, two studies from referral centers in Italy reported higher persistence or recurrence rates in women diagnosed with thyroid cancer during pregnancy or within 12–24 months of delivery compared to both nulliparous women and those diagnosed after a longer interval post partum (472,474). DTC persistence or recurrence was defined as (a) either stimulated or suppressed serum Tg >2 ng/mL, (b) increasing levels of Tg autoantibodies, (c) sonography detection of metastatic lymph nodes, or (d) uptake outside the thyroid bed on scintigraphy. Subjects were followed for a median of about 6 years after definitive therapy. Vannucchi *et al.* (472) observed a much higher than expected persistence/recurrence rate (60%) in the pregnant/early postpartum group than otherwise anticipated for Stage 1 disease, compared with 4% in women diagnosed more than 12 months after delivery and 13% in nulliparous women. The majority of the recurrences (67%) were identified by Tg or TgAb criteria without identification of structural disease, so it is unclear if Tg/TgAb detection is associated with other confounding factors such as potentially less-aggressive surgery if done during gestation. Messuti *et al.* (474) reported a recurrence rate of 10% that was more typical for early stage disease, but this was statistically higher than the 1.3% for women diagnosed 2 years after delivery and 4.3% for nulliparous women. No details for recurrence classification, biochemical versus structural, were provided. Importantly, delaying surgery until after delivery did not further affect recurrence rates among women diagnosed with DTC during gestation in either study. Although several potential explanations exist for the poorer prognosis of pregnant/early postpartum women versus the comparator groups, findings are not consistent in the two studies. Metastatic cervical adenopathy at diagnosis was more frequent in pregnant/postpartum women compared to the other groups in one study (63% vs. ~40%) (472) but not in the other (474). Vannucchi *et al.* (472) documented more frequent presence of estrogen receptor α in the tumors of women diagnosed during pregnancy/early post partum as compared to the other two groups, which may indicate that the poorer prognosis is related to estrogen-mediated growth and stimulation of the MAPK pathway. However, this was not confirmed in the study by Messuti *et al.* (474). The discrepancy may be due to methodologic issues because a less-dilute estrogen receptor Ab was employed in the study with the positive findings. Hence, there is a recent observation of higher persistence/recurrence rates for women diagnosed with DTC during or immediately after pregnancy, without elucidation of pathophysiologic mechanisms. Further studies may help to clarify causality. Importantly, survival is not compromised.

The impact of pregnancy on women with medullary or anaplastic carcinoma is unknown.

Question 69: What are the perioperative risks to mother and fetus of thyroid surgery during pregnancy?

Surgery is the treatment of choice for DTC. As deferring surgery until postpartum has not been associated with a worse prognosis, it is imperative to assess maternal and neonatal complications before recommending an operation during pregnancy. Between 1986 and 2008, nine studies evaluated the impact of thyroidectomy during pregnancy on a total of

113 patients (study size ranged from 1 to 96) (437,441,442, 444,446,475–478). The majority, but not all, of the operations were performed in the second trimester. There were no maternal or fetal complications in any of the studies. A 2009 population-based study of nonfederal hospitals in the United States compared 201 pregnant women who underwent thyroid and parathyroid surgery during pregnancy with 31,155 similarly treated nonpregnant women. One hundred sixty-five operations were thyroid related and 46% of the women had thyroid cancer. Pregnant patients had a higher rate of endocrine and general complications, longer lengths of stay, and higher hospital costs. The fetal and maternal complication rates were 5.5% and 4.5%, respectively (479). Interpretation of the results of this study is difficult because there were substantial baseline differences between the two groups. Pregnant women were more likely to have either urgent or emergent admissions and had a higher percentage of government insurance. However, surgical volume was also an independent predictor of poor outcomes, so that if surgery is performed during pregnancy, an experienced surgeon is preferred. Recently, a study from Japan focused on 45 patients with DTC, of whom 24 had thyroidectomy during pregnancy (19 during the second trimester) and 21 had surgery after delivery. The groups were compared with regard to disease and other factors. There was no difference in recurrence rates in the mothers and there were no pregnancy losses or birth defects. The authors concluded that although thyroidectomy can be performed safely in the second trimester, surgery after delivery is recommended for most patients with nonaggressive DTC (480). Thus, in instances in which surgery during pregnancy is indicated or desired, it should be performed in the second trimester in order to minimize complications to both the mother and fetus (altered organogenesis and spontaneous abortion in the first trimester, preterm labor and delivery in the third trimester) (481), preferably by an experienced thyroid surgeon. The risk of post-thyroidectomy maternal hypothyroidism and hypoparathyroidism should also be considered.

Question 70: How should cytologically benign thyroid nodules be managed during pregnancy?

Although pregnancy is a risk factor for modest progression of nodular thyroid disease, there is no evidence demonstrating that LT4 is effective in decreasing the size or arresting the growth of thyroid nodules during pregnancy. Hence, LT4 suppressive therapy for thyroid nodules is not recommended during pregnancy. Nodules that were benign on FNA but show rapid growth or ultrasound changes suspicious for malignancy should be evaluated with a repeat FNA and be considered for surgical intervention. In the absence of rapid growth, nodules with biopsies that are benign do not require surgery during pregnancy (482).

RECOMMENDATION 60

Pregnant women with cytologically benign thyroid nodules do not require special surveillance strategies during pregnancy and should be managed according to the 2015 ATA Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer.

Strong recommendation, moderate-quality evidence.

Question 71: How should cytologically indeterminate nodules be managed during pregnancy?

There have been no prospective studies evaluating the outcome and prognosis of women with an FNA that is interpreted as either atypia of undetermined significance/follicular lesion of undetermined significance (AUS/FLUS), suspicious for follicular neoplasm (SFN), or suspicious for malignancy (SUSP). The reported malignancy rates associated with these cytologic diagnoses range from 6% to 48% for AUS/FLUS, 14% to 34% for SFN, and 53% to 87% for SUSP (483). Although molecular testing for cytologically indeterminate nodules is now being considered, no validation studies address application of these tests in pregnant women. It is theoretically possible that thyroid gestational stimulation may alter a nodule's gene expression and change diagnostic performance of molecular tests based upon RNA expression, whereas testing based upon either single base-pair DNA mutations or translocations would be less likely to be affected.

Since prognosis for DTC diagnosed during pregnancy is not adversely affected by performing surgery post partum, it is reasonable to defer surgery until following delivery. Because the majority of these women will have benign nodules, LT4 therapy during pregnancy is not recommended.

■ **RECOMMENDATION 61**

Pregnant women with cytologically indeterminate (AUS/FLUS, SFN, or SUSP) nodules, in the absence of cytologically malignant lymph nodes or other signs of metastatic disease, do not routinely require surgery while pregnant.

Strong recommendation, moderate-quality evidence.

■ **RECOMMENDATION 62**

During pregnancy, if there is clinical suspicion of an aggressive behavior in cytologically indeterminate nodules, surgery may be considered.

Weak recommendation, low-quality evidence.

■ **RECOMMENDATION 63**

Molecular testing is not recommended for evaluation of cytologically indeterminate nodules during pregnancy.

Strong recommendation, low-quality evidence.

Question 72: How should newly diagnosed thyroid carcinoma be managed during pregnancy?

The 2015 ATA guidelines (443) recommend that a nodule with cytology indicating papillary thyroid carcinoma discovered early in pregnancy should be monitored sonographically and, if either it grows substantially by 24 weeks gestation (50% in volume and 20% in diameter in two dimensions), or if metastatic cervical lymph nodes are present, surgery should be considered in the second trimester. However, if it remains stable by midgestation, or if it is diagnosed in the second half of pregnancy, surgery may be performed after delivery. Surgery in the second trimester is an option if the DTC is advanced stage at diagnosis or if the cytology indicates medullary or anaplastic carcinoma.

If surgery is not performed, the utility of thyroid hormone therapy targeted to lower serum TSH levels to improve the prognosis of DTC diagnosed during gestation is not known. Because higher serum TSH levels may be correlated with a more advanced stage of cancer at surgery (484), if the patient's serum TSH is >2 mU/L, it may be reasonable to initiate thyroid hormone therapy to maintain the TSH between 0.3 and 2.0 mU/L for the remainder of gestation.

■ **RECOMMENDATION 64**

PTC detected in early pregnancy should be monitored sonographically. If it grows substantially before 24–26 weeks gestation, or if cytologically malignant cervical lymph nodes are present, surgery should be considered during pregnancy. However, if the disease remains stable by midgestation, or if it is diagnosed in the second half of pregnancy, surgery may be deferred until after delivery.

Weak recommendation, low-quality evidence.

■ **RECOMMENDATION 65**

The impact of pregnancy on women with newly diagnosed medullary carcinoma or anaplastic cancer is unknown. However, a delay in treatment is likely to adversely affect outcome. Therefore, surgery should be strongly considered, following assessment of all clinical factors.

Strong recommendation, low-quality evidence.

Question 73: What are the TSH goals for pregnant women with previously treated thyroid cancer receiving LT4 therapy?

Based on studies which have demonstrated a lack of maternal or neonatal complications from subclinical hyperthyroidism, it is reasonable to assume that the preconception degree of TSH suppression can be safely maintained throughout pregnancy. The appropriate level of TSH suppression depends upon preconception risk of residual or recurrent disease. According to the 2009 and 2015 ATA management guidelines for DTC (443,482), and the ETA consensus (485), the serum TSH should be maintained indefinitely below 0.1 mU/L in patients with persistent structural disease. Target TSH values are first based upon initial risk of recurrence determined postoperatively (Table 12 of the 2015 ATA guidelines) and then are modified based upon classification of the patient's response to therapy as defined in Table 13 of the 2015 ATA guidelines. Hence, for a patient at initial high risk for recurrence, TSH suppression at or below 0.1 mU/L is recommended. If the patient then demonstrates an excellent response to therapy at one year with an undetectable suppressed serum Tg and negative imaging, the TSH target may rise to the lower half of the reference range.

The main challenge in caring for women with previously treated DTC is maintaining the TSH level within the preconception range. In a recent report (325), thyroid cancer patients were reported to require smaller LT4 dose increases than patients who had undergone thyroid ablation for benign thyroid disorders or patients with primary hypothyroidism. However, this outcome was because the nonpregnant serum level TSH was <0.1 mU/L in 83% of the thyroid cancer patients prior to pregnancy and this target was not achieved during gestation. It is likely that if the prepregnancy TSH

target level is maintained throughout gestation, the LT4 dosage increase will be similar to the ~49% required in ablated hypothyroidism. Patients require careful monitoring of thyroid function tests in order to avoid hypothyroidism.

Thyroid function should be evaluated as soon as pregnancy is confirmed. The adequacy of LT4 treatment should be checked 4 weeks after any LT4 dose change. The same laboratory should be utilized to monitor TSH and Tg levels before, during, and after pregnancy.

■ RECOMMENDATION 66

Pregnant women with thyroid cancer should be managed at the same TSH goal as determined preconception. TSH should be monitored approximately every 4 weeks until 16–20 weeks of gestation, and at least once between 26 and 32 weeks of gestation.

Strong recommendation, moderate-quality evidence.

Question 74: What is the effect of therapeutic radioactive iodine treatment on subsequent pregnancies?

Following surgery for DTC, many patients will receive an ablative dose of radioactive iodine. The possible deleterious effect of radiation on gonadal function and the outcome of subsequent pregnancies has been evaluated by Sawka *et al.* (486) and Garsi *et al.* (487); the latter collected 2673 pregnancies, 483 of which occurred after radioiodine treatment. Neither study found an increased risk of infertility, pregnancy loss, stillbirths, neonatal mortality, congenital malformations, preterm births, low birth weight, or death during the first year of life or cancers in offspring.

Radioiodine treatment may lead to suboptimal thyroid hormonal control during the month following administration. The potential adverse effect upon the pregnancy is described in Section VII. It therefore seems reasonable to wait a minimum of 6 months to ensure that thyroid hormonal control is stable before conceiving following radioiodine ablative therapy.

Iodine 131 may affect spermatogenesis. In one study following men after ¹³¹I therapy, there was a dose-dependent increase in follicle-stimulating hormone levels and a reduction in normokinetic sperm (488). Therefore, it seems prudent for a man who has received ¹³¹I to wait 120 days (the lifespan of sperm) after ¹³¹I therapy before attempting to conceive.

■ RECOMMENDATION 67

Pregnancy should be deferred for 6 months after a woman has received therapeutic radioactive iodine (¹³¹I) treatment.

Strong recommendation, low-quality evidence.

Question 75: What is the effect of tyrosine kinase inhibitors on pregnancy?

Several tyrosine kinase inhibitor (TKI) medications are now FDA approved for the therapy of metastatic differentiated and medullary thyroid cancer. These include sorafenib, lenvatinib, and cabozantinib. All three drugs have demonstrated both teratogenicity and embryo toxicity in animals when administered at doses below the recommended human dose (489). There are no human studies. The FDA recom-

mends that women be advised of the potential risk to the fetus with TKI medications. However, specific advisories vary by medication. Furthermore, the use of any TKI must be guided by an assessment of risks to benefits that is also affected by the stage of disease and recommended drug indications. For sorafenib, the FDA prescribing information counsels to avoid pregnancy while taking the drug. For lenvatinib, contraception is explicitly recommended. For cabozantinib, no additional warnings are listed (489).

Question 76: Does pregnancy increase the risk of DTC recurrence?

Five studies have evaluated the impact of pregnancy after a woman has been treated for DTC. Two were published five decades ago, predating contemporary tools for thyroid cancer surveillance (490,491). Still, Rosvoll *et al.* (490) demonstrated pregnancy did not accelerate tumor growth in the 22 women who had stable or slowly progressive metastatic disease. Hill *et al.* (491) found no difference in clinically detected thyroid cancer recurrence rates in 70 women who had one or more pregnancies following initial diagnosis of DTC and 109 women who had no subsequent pregnancies. A more recent study from Memorial Sloan Kettering reported on 36 women who became pregnant between 1997 and 2006, a median of 4.3 years after initial treatment for DTC (492). Although all had serum Tg testing, only 16 (44%) had ultrasound imaging before and after delivery. Disease progression occurred in one of the two women who had metastatic cervical lymph nodes on baseline imaging (the lymph node grew from 2.3 cm to 2.7 cm). The mean suppressed Tg after delivery was not significantly different than the prepartum value. However, eight women had postpartum Tg values more than 20% higher than before pregnancy (three with known disease, five with no clinical evidence of disease). No evidence of recurrence was detected in the early postpartum period in women with negative prepregnancy neck ultrasound and a suppressed serum Tg <1 ng/mL. Two more recent studies (493,494) routinely used neck sonography and sensitive Tg assays to evaluate cancer progression during pregnancy. No recurrence was observed in 64 Brazilian women (48 papillary cancer and 16 follicular cancer) who had suppressed Tg levels <2 ng/mL and negative neck sonography prepregnancy. However, follow-up was limited to only 6 months after delivery (493). Using a definition of disease-free as a suppressed Tg <0.9 ng/mL and negative neck ultrasound, Hirsch and colleagues (494) followed 63 women who had given birth after receiving treatment for DTC (all papillary thyroid cancer) for a mean of 4.8 years. None of the 50 women categorized as disease-free had progression of disease. However, 6 of the 13 women (46%) with persistent disease prepregnancy experienced progression. Metastatic cervical lymph nodes grew in 50% of those with known prepregnancy nodal disease, and new nodal metastases appeared in two of the seven women with negative baseline sonography.

Thus, pregnancy does not pose a risk for tumor recurrence in women without structural or biochemical disease present prior to the pregnancy. Therefore, women with an excellent response to therapy as defined by the 2015 ATA guidelines (443) do not require additional monitoring during gestation. However, pregnancy may represent a stimulus to thyroid

cancer growth in patients with known structural (ATA 2015 structural incomplete response to therapy) or biochemical (ATA 2015 biochemical incomplete response to therapy) disease present at the time of conception and requires monitoring.

Question 77: What type of monitoring should be performed during pregnancy in a patient who has already been treated for DTC prior to pregnancy?

■ RECOMMENDATION 68

Ultrasound and Tg monitoring during pregnancy is not required in women with a history of previously treated DTC with undetectable serum Tg levels (in the absence of Tg autoantibodies) classified as having no biochemical or structural evidence of disease prior to pregnancy.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 69

Ultrasound and Tg monitoring should be performed during pregnancy in women diagnosed with well-differentiated thyroid cancer and a biochemically or structurally incomplete response to therapy, or in patients known to have active recurrent or residual disease.

Strong recommendation, moderate-quality evidence.

Question 78: What type of monitoring should be performed during pregnancy in a patient who is under active surveillance for papillary thyroid microcarcinoma?

Although FNA of subcentimeter nodules in the absence of suspicious cervical adenopathy is not recommended by the 2015 ATA guidelines (443), there are times when this has been performed with subsequent diagnosis of a papillary thyroid microcarcinoma (PTMC). In some situations, women with this diagnosis may have chosen active surveillance. Observation with active surveillance has been demonstrated to be a viable option for patients cytologically diagnosed with PTMC (size <1 cm) in the absence of suspicious lymph nodes or extrathyroidal invasion by sonographic imaging (495). Although tumor growth >3 mm and appearance of novel metastatic lymph nodes occurred respectively overall in 8% and 3.8% of a cohort of 1235 Japanese patients, progression was more likely in patients younger than 40 years of age. Therefore, the same group published their observations of nine women with PTMC who became pregnant during active surveillance and compared outcomes to 27 age-matched nonpregnant women. During pregnancy, growth ≥3 mm occurred more frequently in pregnant (44%) than nonpregnant (11%) women ($p=0.0497$). There was no correlation between growth and gestational TSH or Tg levels, and novel metastatic cervical lymph nodes were not observed. Three of the four patients who experienced tumor growth had surgery post partum, but the tumor decreased in size post partum in the fourth patient who is still under active surveillance. Furthermore, the PTMC enlarged by less than 3 mm in the other five pregnant women, but then decreased in three postpartum (496). This supports the hypothesis that pregnancy is either permissive for or promotes growth of papillary thyroid cancer. While this may have no impact on overall

health, it seems reasonable to monitor such women with serial sonographic evaluations.

■ RECOMMENDATION 70

Ultrasound monitoring of the maternal thyroid should be performed each trimester during pregnancy in women diagnosed with PTMC who are under active surveillance.

Weak recommendation, low-quality evidence.

Question 79: What special considerations should be followed for women with medullary cancer due to germline RET mutations?

Hereditary medullary thyroid cancer (MTC) may occur as a result of activating germline mutations of the *RET* oncogene. Over 100 mutations, duplications, insertions, or deletions involving *RET* have been identified in patients with hereditary MEN2a (including familial MTC) or MEN2b syndromes. The newly Revised American Thyroid Association Guidelines for the Management of Medullary Thyroid Carcinoma have graded the most common mutations according to risk of aggressive MTC and likelihood of concomitant pheochromocytoma and hyperparathyroidism (497). For a woman with a given *RET* mutation with or without clinical MTC, who is either pregnant or contemplating conception, the associated risks of MTC aggressiveness combined with her own history may inform her about whether to seek genetic counseling. Both prenatal and preimplantation genetic testing are available, and the reader is referred to the excellent discussion in the 2015 ATA guidelines that culminates in Recommendation 12: “The duty to warn of genetic risk extends to both preconception and prenatal contexts. Genetic counseling about the options of pre-implantation or prenatal diagnostic testing should be considered for all *RET* mutation carriers of childbearing age, particularly those with MEN2B. Parents who do not wish to have prenatal *RET* gene mutation testing should be offered genetic counseling and informed of the availability of genetic testing of their child to detect a mutated *RET* allele. This is particularly important for mutations associated with the onset of MTC before 5 years of age.” In addition, testing to exclude a pheochromocytoma should be done prior to pregnancy for all women with MEN2. No studies address the effect of pregnancy on MTC progression in patients with biochemical or structural evidence of residual or metastatic disease.

X. FETAL AND NEONATAL CONSIDERATIONS

Question 80: What is the relationship between maternal and fetal thyroid hormone status?

Under normal circumstances the fetal hypothalamic-pituitary-thyroid system develops relatively independently of maternal influence because of the presence of the placenta, which regulates the passage of many substances, including T4, to the fetus. Circulating fetal thyroid hormone levels therefore largely reflect the stage of fetal maturation.

Thyroid hormone receptors are present in the fetal brain at low concentrations up to week 10 of gestation. Thyroid hormone receptor concentrations then rapidly increase 10-fold through week 16 of pregnancy (498).

Serum TT4 and FT4 are first measurable in low levels in fetal serum at 12–14 weeks of development (499–501).

Based on values obtained by fetal cord sampling in normal pregnant women, it has been estimated that the mean fetal serum TT4 is 2 $\mu\text{g/dL}$ (26 nmol/L) at 12 weeks, and a relatively larger proportion is in the free form. Beginning in midgestation, the fetal TT4 concentration begins to increase and typically reaches values comparable to nonpregnant females (10 $\mu\text{g/dL}$ [138 nmol/L]) by 36 weeks. This rise in TT4 follows an increase in the serum TBG concentration due primarily to the stimulatory effects of maternal estrogen on the fetal liver. In addition, owing to up-regulation of the TSH receptor, there is an increase in the fetal FT4 concentration from a mean of approximately 0.1 ng/dL (1.3 pmol/L) to 2 ng/dL (25.7 pmol/L), beginning early in the third trimester (502).

In contrast to T4, the fetal circulating levels of its active metabolite T3 remain significantly lower in fetal life than postnatally, whereas the inactive metabolites reverse T3 and T3 sulfate are elevated. The fetal serum T3 concentration is approximately 6 ng/dL (0.09 nmol/L) at 12 weeks, rising to 45 ng/dL (0.68 nmol/L) at 36 weeks. Despite the low levels of circulating T3, brain T3 levels are 60%–80% those of the adult by fetal age 20–26 weeks (503). This reflects the importance of local conversion of T4 to T3 in the brain itself. During the second trimester, expression of type 2 deiodinase, which converts T4 to T3, increases in the fetal cerebral cortex in parallel to the T3 concentrations (504). T3 sulfate may also serve as an alternate source of T3 for the pituitary (505).

The serum concentration of TSH rises modestly from ~ 4 mU/L at 12 weeks to 8 mU/L at term and is always higher than the corresponding maternal levels (500,506). The reason for this is not completely understood but may be a consequence of incomplete maturation of the hypothalamic-pituitary-thyroid axis and/or the high TRH levels in the fetus.

Question 81: What is the role of placental transfer?

Thyroid hormone. The placenta plays a key role in regulating the exchange of products between the mother and fetus. The amount of placental transfer and the consequent effect on fetal thyroid function varies widely.

Under normal circumstances, the human placenta has modest permeability for T4 because of the predominance of placental deiodinase (D3), which serves to inactivate most of the thyroid hormone presented from the maternal to the fetal circulation. The iodide released by this process can be used as a substrate for fetal thyroid hormone synthesis. The placenta also expresses some D2 (an activating deiodinase), but placental D3 activity is approximately 200-fold greater than D2; the activity of both D2 and D3 falls as pregnancy progresses. The placenta also expresses a range of thyroid hormone transporters, binding proteins, sulfotransferases, and sulfatases. The balance of all of these enzymatic processes determines the amount of T4 transfer that occurs (507).

In the first half of pregnancy, before fetal thyroid maturation, T4 is detectable in the fetal brain, indicating that mechanisms exist for small but significant transfer of maternal T4 to the fetal circulation (503,507). This transfer of maternal thyroid hormone is particularly important when the fetus is hypothyroid and probably explains the normal or near-normal cognitive outcomes of babies diagnosed with severe congenital hypothyroidism at birth and treated adequately and sufficiently early following birth. For example, even in infants with congenital absence of thyroid peroxidase

(TPO), the cord serum concentration of T4 is between 25% and 50% of normal (508). This maternal T4 disappears rapidly from the newborn circulation, with a half-life of approximately 3–4 days.

Iodine. Iodine is actively transported across the placenta from the maternal circulation to the fetus. The placenta actively concentrates iodine through expression of the sodium iodide transporter, the concentration of which increases with gestational age (509).

Immunoglobulin G. Maternal autoantibodies of the immunoglobulin G (IgG) class are transported across the placenta from approximately midgestation onward by a cellular receptor termed the neonatal Fc receptor, thereby providing the fetus and neonate with humoral immune protection until the infant starts producing its own IgG (510). In mothers with autoimmune thyroid disease, autoantibodies to TPO, Tg, and the TSH receptor can be transmitted to the fetus in a similar fashion. Whereas TPO antibodies and Tg antibodies do not significantly affect fetal or neonatal thyroid function, antibodies to the TSH receptor can stimulate or block thyroid hormonogenesis or be biologically neutral (511) (see sections VII and VIII on Hypothyroidism and Thyrotoxicosis). Maternal transmission of antibodies to thyroid hormone has also been reported as a cause of congenital hypothyroidism (512).

Hyperthyroidism generally develops only in babies born to mothers with the most potent (or highest concentrations of) stimulatory TRAb activity (513,514). This corresponds to 1%–2% of mothers with GD, or 1 in 50,000 newborns, an incidence that is approximately 4 times higher than that for transient neonatal hypothyroidism due to maternal TSH receptor blocking antibodies (515). Some mothers have mixtures of stimulating and blocking antibodies in their circulation, the relative proportion of which may change over time. Not surprisingly, the clinical picture in the fetus and neonate of these mothers is complex and depends not only on the relative proportion of each activity in the maternal circulation at any one time but on the rate of their clearance from the neonatal circulation postpartum. Thus, one affected mother gave birth, in turn, to a normal infant, a baby with transient hyperthyroidism, and one with transient hypothyroidism (516). In another neonate, the onset of hyperthyroidism did not become apparent until 1–2 months postpartum when the higher-affinity blocking antibodies had been cleared from the neonatal circulation (517). In the latter case, multiple TSH receptor stimulating and blocking antibodies were isolated from maternal peripheral lymphocytes. Each monoclonal antibody recognized different antigenic determinants (epitopes) on the receptor and had different functional properties (518). Occasionally, neonatal hyperthyroidism may occur in infants born to hypothyroid mothers. In these situations, the maternal thyroid has been destroyed by prior radioablation or surgery so that potent thyroid stimulating antibodies, present in the maternal circulation, affect the fetal thyroid but not the maternal thyroid.

TSH receptor blocking, Ab-induced congenital hypothyroidism, is much less common than neonatal GD but should be suspected in any child with congenital hypothyroidism born to a mother with severe autoimmune thyroid disease, especially if there is a history of a similarly affected sibling (515). Blocking Ab-induced congenital hypothyroidism is

important to identify because it identifies a child who is likely to have transient disease (515,519). Affected infants may have significantly impaired cognitive outcomes despite early and adequate postnatal treatment if maternal hypothyroidism was present and untreated during gestation (520). The half-life of IgG is 1–2 weeks, and the duration of the Ab-induced hypothyroidism is usually about 3 months, but it may be longer.

Significant transplacental transfer of maternal TRH occurs, but not maternal TSH, hCG, or Tg. Importantly, both ATDs (PTU and MMI) as well as TSH receptor antibodies can pass from the mother and the fetus. This may have significant effects on fetal and neonatal thyroid function, as discussed elsewhere.

Question 82: When maternal thyroid abnormalities are detected, how and when should information be provided to the neonatologist or pediatrician?

Question 83: What information should be provided to the neonatologist or pediatrician?

Most infants born to women with known thyroid illness are healthy. Furthermore, all newborn infants in the United States are screened for thyroid dysfunction as part of universal screening mandates. Nonetheless, maternal thyroid illness, abnormal maternal thyroid function, the presence of maternal TSH receptor antibodies, and/or the use of antithyroid medications during gestation can each contribute to adverse effects in the newborn. Therefore, knowledge of these facts is important and should be documented in the newborn infant's medical record. In many cases, direct communication with the neonatologist or pediatrician is recommended, especially in the setting of severe maternal hyperthyroidism or maternal use of ATDs at any time throughout pregnancy. The etiology, timing, severity, and treatment of maternal thyroid disease should all be conveyed. In women with GD (even if previously ablated with ^{131}I or athyreotic following surgery) concentrations of circulating TRAb should be assessed at initial thyroid function testing in early pregnancy, and if TRAb is positive, again at weeks 18–22 (see Section VIII). Such information should be documented and conveyed to the pediatrician because maternally derived thyroid antibodies as well as antithyroid medication can transfer to the fetus and affect newborn thyroid health.

■ **RECOMMENDATION 71**

A history of maternal thyroid illness, use of antithyroid medications (PTU, MMI) during gestation, or measurements of abnormal maternal thyroid function or TRAb during gestation should be communicated to the newborn's neonatologist or pediatrician.

Strong recommendation, moderate-quality evidence.

■ **RECOMMENDATION 72**

The severity of maternal and fetal thyroid illness should guide the timing of communication. Severe, progressive, or complex thyroid illness during pregnancy mandates communication with the neonatologist or pediatrician before birth and consideration of consultation with a pedi-

atric endocrinologist. Most other illness is optimally communicated shortly after birth.

Strong recommendation, moderate-quality evidence.

Question 84: Should all neonates be screened for thyroid dysfunction?

Congenital hypothyroidism is one of the most frequent treatable causes of intellectual impairment, and newborn screening for the detection of congenital hypothyroidism is performed routinely in all 50 states of the United States, as well as in Canada, Europe, Israel, Japan, Australia, and New Zealand (521). Screening programs are also being introduced in many developing countries, accounting overall for approximately one third of the world's population (521). Blood spot specimens eluted from filter paper are utilized. The optimal timing of measurement is at 2–5 days of age in order to avoid confounding by the physiologic surge in neonatal TSH, which occurs shortly after birth. Some programs employ primary T4 screening with TSH measurement when specimens demonstrate a value below a specified cutoff. Others use a primary TSH determination with reflex T4 strategy. The introduction of newborn screening for congenital hypothyroidism has led to the virtual elimination of intellectual impairment due to hypothyroidism so long as early and adequate postnatal LT4 treatment is initiated immediately (522,523).

■ **RECOMMENDATION 73**

All newborns should be screened for hypothyroidism by blood spot analysis typically 2–5 days after birth.

Strong recommendation, high-quality evidence.

Question 85: How is neonatal hypothyroidism treated?

In women receiving ATDs at the time of delivery, transplacental transfer of the medication can potentially induce neonatal hypothyroidism. In such cases, neonatal metabolism removes the remaining MMI or PTU from the newborn circulation. This results in the return of normal thyroid function, typically within 3–5 days. For neonates with congenital hypothyroidism, LT4 must be administered. The recommended starting dose for full-term infants is 10–15 $\mu\text{g}/\text{kg}/\text{d}$, and depends on the severity of the initial hypothyroidism (522,523). In premature hypothyroid infants a lower dose is generally utilized. For optimal cognitive outcomes, therapy should be initiated within 2 weeks of life. Both inadequate and excess thyroid hormone replacement may be harmful, so close follow-up is important, particularly during the first 3 years of life. During this time, brain development is thyroid hormone dependent. Age-dependent normative values for TSH and T4 should be used to guide therapy. Until further information becomes available, infants with severe congenital hypothyroidism (i.e., initial serum TSH concentration $>100\text{ mU/L}$) should remain on a single formulation of LT4 without substitution (524). Treatment should be guided by a pediatric endocrinologist or experienced pediatrician.

The presence of a goiter in a newborn should prompt referral to a pediatric endocrinologist. An ultrasound examination should be performed to evaluate the patency of the trachea. A goiter may reflect a hypothyroid or hyperthyroid

state in response to maternal thyroid illness or overtreatment with antithyroid medications. Regardless, close observation and urgent treatment may be required to achieve a euthyroid state and avoid airway compromise in the newborn.

For a more detailed discussion of this and other aspects of neonatal hypothyroidism, the reader is referred to several recent excellent reviews and guidelines (522,523,525).

Question 86: How is neonatal hyperthyroidism treated?

Most neonatal hyperthyroidism is caused by maternal transfer of TRAb to the fetus. Typically, neonatal GD does not present until the end of the first week of life when maternal ATD, but not the TRAb, have been cleared from the neonatal circulation. This may be delayed in babies born to mothers with a mixture of stimulating and blocking antibodies. Thus, results on newborn screening may be paradoxically normal. A high degree of suspicion for infants at risk is important.

The conventional treatment for neonatal hyperthyroidism is PTU 5–10 mg/kg administered in divided doses. However recent evidence of rare hepatotoxicity due to this agent has led to a switch to MMI therapy (0.5–1 mg/d) by many pediatric endocrinologists. Propranolol (2 mg/kg) may be added if the hyperthyroidism is severe. Close follow-up of the affected newborn is important, with downward adjustment in the dose of ATD required as the hyperthyroidism resolves. The usual duration of neonatal GD is 1–3 months, but it depends on antibody potency. Separate from neonatal GD (which is self-limited), neonatal hyperthyroidism may rarely be caused by a gain-of-function mutation in the TSH receptor, or by McCune–Albright syndrome. In such cases, newborn hyperthyroidism may be permanent (526).

XI. THYROID DISEASE AND LACTATION

Question 87: Does maternal thyroid hormone status impact lactation?

Question 88: In the breastfeeding mother, should maternal hypothyroidism be treated to improve lactation?

Question 89: In the breastfeeding mother, should maternal hyperthyroidism be treated to improve lactation?

Abnormal maternal thyroid hormone concentrations (both hypothyroidism and hyperthyroidism) can impact milk let-down and the ability to successfully breastfeed. While there are no controlled or randomized human trials, several case series and cohort analyses have been reported. One study reported 26 human pregnancies complicated by hypothyroidism in New Delhi, India. Deficient lactation was reported in 19.2% of subjects (527). Other case reports describe impaired lactation, which improved upon successful treatment of hypothyroidism (528). Interventional trials in nonhuman mammals corroborate these findings. Hapon and colleagues (529) investigated the effects of PTU-induced hypothyroidism in rats at days 10–12 of lactation. After 30 minutes of suckling, milk excretion was reduced compared to euthyroid controls. The authors suggested mammary function was maintained while a reduced milk ejection reflex was likely

responsible. A separate investigation of Brahman cows demonstrated reduced milk production in PTU-induced hypothyroid cows compared to euthyroid controls. Following cessation of PTU and normalization of thyroid status, lactation in these animals returned to baseline (530). Together, these data strongly suggest maternal hypothyroidism can adversely affect lactation. However, with limited human data and no controlled or randomized intervention studies, it is unclear what proportion of hypothyroid mothers would be affected. Furthermore, it is possible that many mothers experiencing only mild to moderate hypothyroidism could experience only minimal difficulties with breastfeeding and milk production. Importantly, the available data confirm that maternal hypothyroidism itself is not harmful to the breastfeeding infant *per se*, so long as adequate nutrition (milk) is provided. Therefore, it appears most reasonable to recommend evaluation of thyroid function in women who demonstrate poor lactation, with the goal of effectively returning any hypothyroid mother to a euthyroid state.

Maternal hyperthyroidism may also impair lactation, although the data are less clear. No controlled human trial or large cohort analysis has been performed. Two rat studies demonstrated adverse effects of hyperthyroidism on lactation. In one study, moderate to severe hyperthyroidism was induced in pregnant rats during and after pregnancy. Complete lactation failure was noted (531). A second study in which moderate hyperthyroidism was induced demonstrated continued ability of rats to lactate, but impairment of milk ejection. Ultimately, this resulted in death of the offspring due to malnutrition (532). Such limited data, obtained from only nonhuman populations, make it impossible to draw firm conclusions. Therefore, no recommendation can be made to treat maternal hyperthyroidism on the grounds that it will improve lactation.

■ **RECOMMENDATION 74**

As maternal hypothyroidism can adversely impact lactation, women experiencing poor lactation without other identified causes should have TSH measured to assess for thyroid dysfunction.

Weak recommendation, low-quality evidence.

■ **RECOMMENDATION 75**

Given its adverse impact upon milk production and let-down, subclinical and overt hypothyroidism should be treated in lactating women seeking to breastfeed.

Weak recommendation, low-quality evidence.

■ **RECOMMENDATION 76**

The impact of maternal hyperthyroidism upon lactation is not well understood. Therefore, no recommendation to treat maternal hyperthyroidism on the grounds of improving lactation can be made at this time.

No recommendation, insufficient evidence.

Question 90: Is maternal thyroid hormone (thyroxine or LT4) transferred to the newborn via breast milk?

A small amount of maternal thyroid hormone is present in the breast milk of lactating women, and also detected in cow,

goat, and sheep milk. A study investigated T4 and T3 in the breast milk of mothers who delivered very premature infants. Thyroxine concentration in breast milk ranged from 0.17 to 1.83 $\mu\text{g/L}$ (mean 0.83 $\mu\text{g/L}$) (533). Assuming an average consumption of $\sim 750\text{ mL/d}$ of breast milk, this equates to approximately 0.62 μg of T4 per day. Assuming total T4 requirements of 40–60 $\mu\text{g/d}$ for most athyreotic infants, the amount of thyroxine transferred to the infant via the human lactating breast is $\sim 1\%$ of the total daily requirement. Thus, maternal transfer of thyroid hormone does not have a meaningful impact on the infant's thyroid hormone status.

Studies in cows (534), sheep (535), and donkeys (536) similarly find a small but detectable concentration of T4 and T3 in the milk supply. One study demonstrated secretion of 200–300 pg/mL of T3 in cow milk. This translates into a total daily consumption of 0.2–0.4 $\mu\text{g/d}$ depending the age and feeding habits of the infant. Similarly, T4 concentrations in cows' milk demonstrate an average concentration of 0.97 ng/mL, translating into the consumption of approximately 0.75–1.5 μg T4 per day. Thus, dairy milk obtained from cows, sheep, goats, or donkeys has little impact on the thyroid health of the infant or young child.

Question 91: Should diagnostic or therapeutic radiopharmaceuticals be administered to breastfeeding or lactating women?

Because radioactive iodine is concentrated in breast milk, and the half-life of ^{131}I is relatively long (~ 8 days), use of ^{131}I for scanning or treatment is absolutely contraindicated in the breastfeeding mother. The half-life of ^{123}I is 13 hours. If circumstances necessitate the use of ^{123}I for of thyroid uptake measurement or scans while lactating, breast milk should be pumped and discarded for several days until the radioactive iodine has cleared from the body (537).

■ **RECOMMENDATION 77**

The use of ^{131}I is contraindicated during lactation. If required, ^{123}I can be used if breast milk is pumped and discarded for 3–4 days before breastfeeding is resumed. Similarly, Tc-99m pertechnetate administration requires breast milk to be pumped and discarded during the day of testing.

Strong recommendation, moderate-quality evidence.

Question 92: Are ATDs (PTU, MMI) transferred into breast milk, and what are the clinical consequences to the breastfed infant?

Both PTU and MMI can be detected in the breast milk of treated hyperthyroid women. This finding raised initial concern that consumption of these medications could prove detrimental to the health of the breastfeeding infant. However, studies first performed using PTU confirmed that only a very small amount of the drug is transferred from maternal serum into breast milk. In a study of nine women given 200 mg PTU orally, milk PTU concentration was measured for 4 hours thereafter and only 0.007%–0.077% of the ingested dose was detected. The authors calculated that a lactating mother consuming PTU 200 mg three times daily would transmit only 149 μg (0.149 mg) of PTU daily to her infant (538). This is well below a therapeutic dose and deemed to pose no risk to the breastfeeding infant. Separate studies confirm normal thyroid

function in breastfed infants of mothers consuming PTU (539,540). In these studies, two infants were found to have an elevated TSH within 1 week of birth, however, these values normalized thereafter despite continued breastfeeding. This suggests the cause of neonatal hypothyroidism in these infants was transplacental passage of PTU before birth, as opposed to any adverse effect of PTU transferred via breast milk. The remaining children who consumed breast milk expressed from mothers taking PTU had normal thyroid function.

Studies of MMI or CM, confirm a 4- to 7-fold higher proportion of the medication transferred into maternal milk in comparison to PTU. Approximately 0.1%–0.2% of an orally administered MMI/CM dose is excreted into breast milk (541,542). Johansen and colleagues (541) calculated that a single 40 mg dose of MMI could result in delivery of 70 μg (0.07 mg) to the breastfeeding infant. Several studies have separately investigated the effect of maternally ingested MMI or CM upon the thyroid status of the breastfeeding infant (543–545). Virtually all have documented normal neonatal thyroid function. Furthermore, several women overtreated with MMI/CM were found to have elevated TSH concentrations (TSH 19–102 mU/L). Even in these situations, normal neonatal thyroid function was nonetheless confirmed in breastfeeding infants.

The largest study investigating the effects of maternal MMI consumption during lactation was performed by Azizi and colleagues (545). Importantly, this study assessed both neonatal thyroid function in the breastfeeding offspring, but also intellectual development and physical growth in a subset of infants. Verbal and performance IQ scores were measured in 14 children who breastfed from MMI-treated mothers, with comparison to 17 control children. Testing was performed between 48 and 74 months of age. No difference was detected in the IQ or physical development of the breastfeeding children compared to the control children (543).

Together, these data have led experts to confirm the safety of low to moderate doses of both PTU and MMI/CM in breastfeeding infants. However, given the relatively small size of the studied population, maximal daily doses of 20 mg MMI or 450 mg PTU are advised (352).

Question 93: What is the approach to the medical treatment of maternal hyperthyroidism in lactating women?

■ **RECOMMENDATION 78**

Excepting treatment decisions specifically made on the grounds of improving lactation (discussed above), the decision to treat hyperthyroidism in lactating women should be guided by the same principles applied to non-lactating women.

Strong recommendation, low-quality evidence.

Question 94: When medical treatment of maternal hyperthyroidism is indicated, what medications should be administered?

■ **RECOMMENDATION 79**

When antithyroid medication is indicated for women who are lactating, both MMI (up to maximal dose of 20 mg/d)

and PTU (up to maximal dose of 450 mg/d) can be administered. Given a small, but detectable amount of both PTU and MMI transferred into breast milk, the lowest effective doses of MMI/CM or PTU should always be administered.

Strong recommendation, moderate-quality evidence.

Question 95: How should breastfeeding children of mothers who are treated with antithyroid medications be monitored?

■ RECOMMENDATION 80

Breastfed children of women who are treated with ATDs should be monitored for appropriate growth and development during routine pediatric health and wellness evaluations. Routine assessment of serum thyroid function in the child is not recommended.

Weak recommendation, moderate-quality evidence.

Question 96: What are the iodine nutritional considerations in lactating women?

Iodine is an essential nutrient required for thyroid hormone production and is primarily derived from the diet. For most breastfeeding infants, breast milk is the sole (or primary) source of nutrition and thus of dietary iodine. Therefore, adequate iodine intake in the lactating mother positively affects infant health. Iodine requirements increase during pregnancy to accommodate the maternal–fetal unit. Post partum, maternal iodine requirements increase even further, as iodine is transferred to the breastfeeding infant. The dietary goals for individual total daily iodine intake (dietary and supplement) recommended by the Institute of Medicine are 290 µg/d for women who are breastfeeding (546). The WHO recommends 250 µg/d for both pregnant women and lactating women (547).

As discussed in Section IV, dietary sources of iodine vary regionally, and the dietary iodine intake of individuals cannot be reliably ascertained either by patient history or by any laboratory measure. Owing to concerns that a portion of women in the United States are mildly to moderately iodine deficient (which can be exacerbated during lactation due to increased demand), iodine supplementation for breastfeeding mothers is recommended. The ATA has previously recommended 150 µg/d as iodine supplementation for all North American women who are breastfeeding (117). The goal of supplementation is to augment rather than replace dietary iodine intake.

Unfortunately, recommendations for iodine supplementation when breastfeeding have not seen widespread adoption. In the NHANES 2001–2006 survey, only 15% of lactating women reported ingesting iodine-containing supplements (121). Furthermore, one analysis confirmed that only 51% of U.S. prenatal vitamins contained iodine (122). Therefore, physician counseling at the time of postbirth hospital discharge is paramount.

■ RECOMMENDATION 81

All breastfeeding women should ingest approximately 250 µg of dietary iodine daily.

Strong recommendation, high-quality evidence.

■ RECOMMENDATION 82

Breastfeeding women should supplement their diet with a daily oral supplement that contains 150 µg of iodine. This is optimally delivered in the form of potassium iodide (present in a multivitamin) because kelp and other forms of seaweed do not provide a consistent delivery of daily iodine.

Strong recommendation, moderate-quality evidence.

■ RECOMMENDATION 83

In severely iodine-deficient, low-resource regions, where universal salt iodization is lacking and daily supplementation is not feasible, lactating women should receive one dose of 400 mg iodine as oral iodized oil soon after delivery.

Strong recommendation, high-quality evidence.

■ RECOMMENDATION 84

As is the case during pregnancy, sustained iodine intake while breastfeeding that exceeds 500–1100 µg/d should be avoided because of concerns about the potential for inducing hypothyroidism in the infant.

Strong recommendation, moderate-quality evidence.

XII. POSTPARTUM THYROIDITIS

Question 97: What is the definition of PPT and what are its clinical implications?

Postpartum thyroiditis is the occurrence of thyroid dysfunction, excluding GD, in the first postpartum year in women who were euthyroid prior to pregnancy (548). This is an inflammatory autoimmune condition. In the classic form, transient thyrotoxicosis is followed by transient hypothyroidism with a return to the euthyroid state by the end of the initial postpartum year (2). The clinical course of PPT varies, with approximately one quarter of patients presenting with the classical form, one quarter with isolated thyrotoxicosis, and one half presenting with isolated hypothyroidism (431). The thyrotoxic phase of PPT typically occurs between 2 and 6 months post partum, but episodes have been reported as late as 1 year following delivery. All episodes of thyrotoxicosis resolve spontaneously. The hypothyroid phase of PPT occurs from 3 to 12 months postpartum with 10%–20% of cases resulting in permanent hypothyroidism. It should be noted, however, that a prospective study reported that 50% of women with PPT remained hypothyroid at the end of the first postpartum year (549).

Question 98: What is the etiology of PPT?

Postpartum thyroiditis is an autoimmune disorder associated with the presence of thyroid antibodies (TPOAb and TgAb), lymphocyte abnormalities, complement activation, increased levels of IgG1, increased NK cell activity, and specific HLA haplotypes (550–552). Women who are thyroid Ab positive in the first trimester have a high risk of developing PPT, ranging from 33% to 50% (553). Women with the highest Ab titers also have the highest risk of PPT (431). The occurrence of PPT reflects the rebound of the immune system in the postpartum period after the relative immune suppression of pregnancy.

Question 99: How should the etiology of new thyrotoxicosis be determined in the postpartum period?

The major challenge is to differentiate thyrotoxicosis caused by PPT from thyrotoxicosis caused by GD. This is an important distinction because the two disease entities require different treatments and have markedly different clinical courses. The timing of onset provides some clues about etiology. In a Japanese hospital study of 42 patients who developed thyrotoxicosis within the first year after pregnancy, 86% of patients who had debut of disease within the first 3 months after delivery had thyroiditis, whereas all who developed thyrotoxicosis after 6.5 months had GD (554).

TSH receptor antibodies are positive in GD in nearly all cases and are typically negative in PPT, although some mixed-type disease is seen. An elevated T4:T3 ratio suggests the presence of PPT. Physical stigmata of GD, such as goiter with a bruit or ophthalmopathy, are diagnostic when present. The radioiodine uptake is elevated or normal in GD and low in the thyrotoxic phase of PPT, but the use of radioactive diagnostic procedures in lactating patients is rarely indicated. Due to their short half-lives, ^{123}I or technetium scans (Tc-99m) may be used in women who are breastfeeding if breast milk is pumped and discarded for several days after the scan pending the isotope used (see Section XI). As described in Section XI, the use of ^{131}I is contraindicated in women who are breastfeeding.

Question 100: What is the prevalence of PPT?

The prevalence of PPT is approximately 5%, although it has varied markedly in different studies, with frequencies reported from 1.1% to 16.7% (555). Women with other autoimmune disorders have an increased risk of PPT. Specifically, the prevalence of PPT is 3–4 times higher in women with diabetes mellitus type 1 compared to unselected populations (556,557). The frequency of PPT is 25% in women with chronic viral hepatitis (558), 14% in women with systemic lupus erythematosus (559), 44% in women with a prior history of GD (560), and 27% in patients with antipituitary antibodies (561). Individuals who recover fully from PPT have a 70% chance of developing PPT in each subsequent pregnancy (562). Women on LT4 therapy secondary to Hashimoto's thyroiditis may develop PPT if their thyroid gland is not completely atrophic (563). Cases of PPT have been reported following pregnancy loss, but the prevalence of PPT following pregnancy loss is unknown (564).

Question 101: What symptoms are associated with PPT?

Postpartum thyroiditis is a painless condition and most women are asymptomatic or only mildly symptomatic during the thyrotoxic phase. This is because the degree of increase in thyroid hormones is typically mild, and T4 levels are usually more elevated than T3. Nevertheless, in prospective studies, reported symptoms include irritability, heat intolerance, fatigue, and palpitations (431,565–567). The hypothyroid phase of PPT is more frequently symptomatic. Symptoms during the hypothyroid phase of PPT may include cold intolerance, dry skin, fatigue, impaired concentration, and paresthesias (566,567). One study demonstrated that patients

with TPO antibodies and PPT had more symptoms than those who were TPOAb negative (568).

Question 102: Is PPT associated with depression?

Studies evaluating the relationship of PPT to postpartum depression have been inconsistent (569–573). Two studies have reported a significant association between the presence of thyroid antibodies and depression (568,574), irrespective of thyroid function, whereas another study showed no association between the presence of microsomal antibodies and postpartum depression (575). A prospective trial of LT4 treatment versus placebo in postpartum TPOAb-positive women resulted in no difference in rates of postpartum depression between the two groups (576).

■ **RECOMMENDATION 85**

All patients with depression, including postpartum depression, should be screened for thyroid dysfunction.

Strong recommendation, low-quality evidence.

Question 103: What is the treatment for the thyrotoxic phase of PPT?

There have been no prospective studies evaluating when and how to treat PPT. Treatment of the thyrotoxic phase is guided by its transitory nature. ATDs (PTU and MMI) are ineffective in treating the thyrotoxic phase of PPT because it is a destructive thyroiditis in which thyroid hormone synthesis is not increased. Symptoms are typically mild. In rare cases when symptoms are clinically significant, propranolol at the lowest possible dose to alleviate symptoms may be used. The thyrotoxic phase of PPT must be differentiated from recurrent or *de novo* GD.

■ **RECOMMENDATION 86**

During the thyrotoxic phase of PPT, symptomatic women may be treated with β -blockers. A β -blocker that is safe for lactating women, such as propranolol or metoprolol, at the lowest possible dose to alleviate symptoms is the treatment of choice. Therapy is typically required for a few weeks.

Strong recommendation, moderate-quality evidence.

■ **RECOMMENDATION 87**

ATDs are not recommended for the treatment of the thyrotoxic phase of PPT.

Strong recommendation, high-quality evidence.

Question 104: Once the thyrotoxic phase of PPT resolves, how often should TSH be measured to screen for the hypothyroid phase?

■ **RECOMMENDATION 88**

Following the resolution of the thyrotoxic phase of PPT, serum TSH should be measured in approximately 4–8 weeks (or if new symptoms develop) to screen for the hypothyroid phase.

Strong recommendation, high-quality evidence.

Question 105: What is the treatment for the hypothyroid phase of PPT?

In women with significant symptoms, those currently lactating or women who are actively attempting pregnancy, treatment should be started (431). LT4 treatment should be considered during the hypothyroid phase of PPT if the patient is mildly symptomatic, or recommended if the patient is considering another conception. If treatment is delayed, thyroid function should be rechecked every 4–8 weeks until euthyroidism is restored and women should be counseled to use contraception.

■ RECOMMENDATION 89

LT4 should be considered for women with symptomatic hypothyroidism due to PPT. If treatment is not initiated, their TSH level should be checked every 4–8 weeks until thyroid function normalizes. LT4 should also be started in hypothyroid women who are attempting pregnancy or who are breastfeeding.

Weak recommendation, moderate-quality evidence.

Question 106: How long should LT4 be continued once initiated?

The length of time that LT4 should be continued has not been systematically evaluated. Guiding principles are to maintain a euthyroid state in women who are attempting pregnancy or pregnant. Tapering LT4 doses in order to determine whether the hypothyroid phase of PPT was transitory or permanent can begin by 12 months post partum. Tapering should be gradual and TSH should be monitored every 6–8 weeks.

■ RECOMMENDATION 90

If LT4 is initiated for PPT, discontinuation of therapy should be attempted after 12 months. Tapering of LT4 should be avoided when a woman is actively attempting pregnancy or is pregnant.

Weak recommendation, low-quality evidence.

Question 107: How often should thyroid function testing be performed after the hypothyroid phase of PPT resolves?

The impact of PPT on long-term thyroid function has been evaluated in seven studies (549,562,577–581). These data demonstrate that 10%–50% of women in whom the hypothyroid phase of PPT initially resolves will ultimately go on to develop permanent hypothyroidism. Factors associated with an increased risk of developing permanent hypothyroidism are multiparity, thyroid hypoechogenicity on ultrasound, greater severity of the initial hypothyroidism, higher TPOAb titers, greater maternal age, and a history of pregnancy loss.

■ RECOMMENDATION 91

Women with a prior history of PPT should have TSH testing annually to evaluate for the development of permanent hypothyroidism.

Strong recommendation, high-quality evidence.

Question 108: Does treatment of thyroid Ab-positive euthyroid women during pregnancy prevent PPT?

Two randomized placebo-controlled trials have evaluated the efficacy of iodine or LT4 treatment during pregnancy to prevent the development of PPT in thyroid Ab-positive women. Neither intervention decreased the incidence of PPT (154,582). A single trial has suggested a benefit of selenium supplementation for the prevention of PPT in TPOAb-positive pregnant women, though this has not been independently validated. Furthermore, selenium use has been associated with an increased risk of type 2 diabetes (156). Currently there is not sufficient evidence to recommend selenium supplementation during pregnancy in thyroid Ab-positive women (see Recommendation 12).

■ RECOMMENDATION 92

Treatment of euthyroid thyroid Ab-positive pregnant woman with either LT4 or iodine to prevent PPT is ineffective and is not recommended.

Strong recommendation, high-quality evidence.

XIII. SCREENING FOR THYROID DYSFUNCTION BEFORE OR DURING PREGNANCY

Whether to universally screen for thyroid disease either before or during pregnancy remains controversial. For universal screening to be recommended, any index condition must be prevalent, associated with adverse health outcomes, and treatable. Furthermore, effective therapy must exist but also be practical and effectively deliverable. Finally, screening must be cost effective.

Screening for thyroid dysfunction in pregnancy fulfills some of these criteria. Hypothyroidism, hyperthyroidism, and thyroid autoimmunity are common conditions, with prevalence rates of 2%–3% (259,260,289,583), 0.1%–0.4% (397), and up to 17% (137), respectively. Untreated overt hypothyroidism and hyperthyroidism are associated with adverse obstetrical and fetal outcomes. Among overtly hypothyroid, but undiagnosed pregnant women, the majority will remain hypothyroid following pregnancy, with a mean time to diagnosis in one study of 5 years (276). However, overt thyroid dysfunction is less common than subclinical dysfunction and is frequently identified by clinical assessment at the time of presentation. Mild TSH suppression can be seen in normal early pregnancy and, even when TSH is $\leq 2.5\%$ of the gestational reference range in association with normal FT4 (cutoff between 0.008 and 0.668 mU/L, depending on the gestational age), it is not associated with adverse obstetrical outcomes (22). Thus, the most notable impact of a universal screening mandate for thyroid dysfunction would be the identification of the large proportion of patients with subclinical hypothyroidism (mild elevations in serum TSH with normal thyroid hormone levels).

Presently, most studies investigating subclinical hypothyroidism suggest an association with adverse obstetrical outcomes that is linear because greater degrees of TSH elevation are associated with increased risks to the pregnancy. Such adverse outcomes also appear to be influenced by concomitant antithyroid autoimmunity. Exemplifying this, a large prospective study of 3315 women demonstrated the additive

effect of anti-TPO positivity to the degree of TSH elevation. In anti-TPO-positive women the risk of pregnancy loss increased significantly beyond a TSH concentration >2.5 mU/L (OR 4.95 for TSH 2.5–5.2), with an even greater increase when TSH was >5.2 mU/L (OR 9.56 for TSH 5.2–10 mU/L), whereas in anti-TPO-negative women, significant increases in risk of pregnancy loss was identified only when TSH concentrations exceeded 5.2 mU/L (OR 3.4 for TSH 5.2–10) (288).

Thyroid status can be accurately assessed with currently available blood tests, including TSH, TT4/FT4, and TPOAb. These tests are relatively inexpensive and widely available. Thus, the principal complexity surrounding the screening question relates to the evidence for treatment effectiveness, especially in the population of pregnant women with subclinical hypothyroidism. While prior retrospective studies have suggested a benefit of LT4 treatment in this population (276,277), such findings have not been sufficiently replicated in blinded, prospective analyses thus far.

Important to this discussion is the understanding of two different adverse endpoints that must be considered. These endpoints are maternal/pregnancy risk (primarily pregnancy loss), and separately the adverse effects on fetal neurocognitive development as manifested by lower IQ in the offspring. These endpoints do not necessarily cosegregate. Therefore, each must be considered separately, as different interventions and their timing may affect such endpoints discordantly.

Question 109: Does identification and LT4 treatment of women with subclinical hypothyroidism or thyroid autoimmunity reduce pregnancy complications and pregnancy loss?

Retrospective studies have frequently suggested a positive correlation between treatment of mild elevations of maternal serum TSH and decreased risk of pregnancy loss. For example, a retrospective study from Belgium assessed the effect of TSH screening in a small cohort of pregnant women. LT4 treatment of TPOAb-positive women with TSH >1 mU/L led to a reduction of pregnancy loss rates from 16% to 0% (180). A separate historical cohort analysis studied >1000 pregnant women in the United Kingdom treated with long-term LT4. Pregnancy loss rates were significantly higher when TSH was >4.5 mU/L (and especially when >10 mU/L) compared with when TSH was <2.5 mU/L (318). The retrospective design and lack of TPOAb data are two important limitations of this study.

Limited RCT data on the effects of treatment of subclinical hypothyroidism in pregnancy are available to date. Negro and colleagues (28) randomized euthyroid (TSH <4.2 mU/L) TPOAb-positive women to treatment with LT4 or no treatment. This intervention demonstrated a significant reduction in preterm deliveries (from 22% to 7%) and miscarriage rates (from 14% to 3.5%). However, limitations of this study include its small sample size and the fact that average time of LT4 initiation was late in the first trimester (mean 10.3 weeks), at a time when many pregnancy losses had already occurred.

Two more recent prospective RCTs have investigated the benefits of universal screening of pregnant women with subsequent LT4 treatment of women with TSH >2.5 mU/L. Negro and colleagues (584) randomized 4562 women to universal screening versus screening of women at high risk for thyroid disease. Women identified to have TPOAb posi-

tivity and TSH concentrations >2.5 mU/L were treated with LT4. The timing of treatment initiation was 8.8 weeks gestation. Even though this is well into the first trimester, limiting assessment of the outcome, this is the earliest randomized intervention trial published to date. In the primary analysis, there was no significant difference in adverse events detected between the universally screened cohort versus those who underwent high-risk screening. The main limitation of this study was its design given that all high-risk patients were treated in both study arms, limiting the ability to detect a treatment effect. However, in a secondary analysis of the low-risk cohort, a significant decrease in adverse obstetrical and neonatal outcomes was detected when a screening and treatment strategy was applied (OR 0.43 [CI 0.26–0.70]). How to best interpret this finding, however, remains unclear because this study used a composite endpoint for assessing adverse outcomes. For these reasons, this study alone is often viewed as insufficient to warrant adoption of a universal screening mandate.

A second investigation employed cluster randomization at two centers in China, comparing a process of universal screening at one hospital (and treatment when TSH >2.5 mU/L) to a traditional nonscreening approach in the second institution. In the latter group, serum was obtained from women in early pregnancy, but banked and not analyzed until the postnatal period. The authors concluded that identifying and treating elevated TSH values early in pregnancy reduced pregnancy loss rates (OR 0.343 [CI 0.21–0.56]) and macrosomia (OR 0.46 [CI 0.28–0.74]) (585). However, these data should be viewed with caution because the timing of enrollment at the two centers varied substantially, likely influencing the reporting of pregnancy loss rates. Notably, the screening arm enrolled women at week 11 of pregnancy, whereas the control arm enrolled women at week 7. Because most pregnancy losses occur during the first trimester, the earlier time at testing in the control group suggests that the much higher rates of pregnancy loss were simply due to early enrollment bias.

Importantly, two additional prospective, randomized trials investigating obstetric outcomes are currently in progress. The Thyroid Antibodies and Levothyroxine Trial (586) is a large RCT conducted in the United Kingdom, randomizing TPOAb-positive euthyroid women with a history of pregnancy loss or infertility to treatment with 50 μ g LT4 versus placebo. The endpoint is successful delivery of the infant beyond 34 weeks of gestation. Separately, the T4Life trial in the Netherlands is randomizing euthyroid anti-TPOAb-positive women with a history of recurrent pregnancy losses to treatment with LT4 (starting before conception) versus placebo. The primary outcome is live birth after 24 weeks gestation (587).

Question 110: Does identification and LT4 treatment of pregnant women with subclinical hypothyroidism improve neurocognitive outcomes in offspring?

With regards to the endpoint of fetal neurocognitive development, there are mixed findings from currently available studies. Haddow and colleagues (276) first published seminal data from a large case-control study investigating the effects of untreated maternal hypothyroidism on the IQ of offspring tested at ~ 8 years. This study analyzed sera from 25,216

women obtained at an average gestational age of 17 weeks. They identified 62 women with either TSH concentrations ≥ 99.7 th percentile, or women with TSH concentrations between the 98 and 99.6th percentiles who also had low FT4 concentrations. The authors compared the IQ of the offspring (at 7–9 years) with that of 124 matched children born to euthyroid mothers. IQ scores were 7 points lower in children born to untreated hypothyroid mothers in comparison to controls. However, the retrospective nature of this study has led to questions regarding its reproducibility.

The largest prospective randomized trial of LT4 therapy investigating the treatment of maternal hypothyroidism or hypothyroxinemia in early pregnancy is the CATS study. This study randomized 22,000 women with singleton pregnancies at <16 weeks gestation to immediate thyroid function testing and LT4 treatment if serum TSH was >97.5 th percentile and/or if serum FT4 was <2.5 th percentile. The control arm received no treatment, though serum was obtained in early pregnancy and banked until testing was performed postpartum. The primary outcome was offspring IQ at 3 years of age. Ultimately, the trial found no significant difference in IQ between the children of 390 treated mothers compared with those of 404 untreated mothers (306). Limitations of this study include the timing of LT4 administration (median initiation >13 weeks), and the mild degree of TSH elevation (median TSH 3.1–3.8 mU/L) in its subjects. Furthermore, the predictive value of early IQ testing at age 3 years remains uncertain. With these major limitations, the negative primary endpoint nonetheless does not support universal screening for thyroid disease in pregnant women.

Two separate randomized trials evaluating the effect of LT4 treatment upon fetal neurodevelopment are currently in progress. The first is a multicenter, randomized placebo-controlled clinical trial conducted by the National Institutes of Health, which seeks to evaluate the effects of LT4 treatment in women with either subclinical hypothyroidism or isolated hypothyroxinemia on offspring cognition. Preliminary results presented as an oral abstract at the 2016 Annual Meeting of the Society for Maternal Fetal Medicine meeting showed that initiation of LT4 treatment in these women at a mean gestational age of 17 weeks had no effect on child IQ (21). Full results are anticipated in 2017. A separate, large prospective study is also underway in China. The Subclinical Hypothyroid during Early Pregnancy (SHEP) trial will screen 21,500 women planning pregnancy and seek to assess the effects of LT4 treatment initiated before conception in women with iodine deficiency, those with subclinical hypothyroidism, and those positive for TPOAb (588). The results of the study are expected between 2018 and 2020.

Importantly, if a screening paradigm is conclusively proved effective, universal screening of women in the first trimester has been shown to be cost effective (589–592). A study by Dosiou *et al.* (590) established the cost-effectiveness of universal screening of pregnant women with both TSH and TPO antibodies in the first trimester compared with a strategy of high-risk screening. At base-case analysis, this study assumed that treatment of identified women with TPOAb positivity and subclinical hypothyroidism reduced pregnancy loss and preterm delivery rates as per the randomized clinical trial of Negro and colleagues (28), and that screening resulted in more prompt diagnosis of PPT and hypothyroidism later in life, in the women who developed these disorders. The study did not

assume any adverse effects of untreated thyroid disease on child IQ. In sensitivity analyses, even when the benefits of screening were limited to detection and treatment of overt hypothyroidism, screening was shown to be highly cost-effective at $<\$8000$ /quality-adjusted life year (QALY). Assumption of benefit for fetal IQ made screening cost-saving. Another recent study from Spain demonstrated similar findings, with an incremental cost-effectiveness ratio of $\text{€}374.28/\text{QALY}$ (592). The cost effectiveness of screening for thyroid disease in pregnancy compares well with that for other widely used screening practices in pregnant women such as cystic fibrosis screening ($\$8290/\text{QALY}$) (593) and prenatal diabetes screening ($\$16,689\text{--}19,339/\text{QALY}$) (594).

Finally, it is important to place the screening discussion in the context of worldwide opinion, and acknowledge the recommendations of other Societies. The 2011 ATA pregnancy guidelines and the 2012 Endocrine Society pregnancy guidelines significantly expanded the definition of “high risk” women from that used in earlier editions to include women >30 years of age. This expanded definition of high risk encompassed many more women in populations with a high mean maternal pregnancy age. In a 2012 study by Potlukova *et al.* (595), with a mean maternal age at pregnancy of 31 years, the addition of age 30 or older as a risk factor increased the proportion of women correctly identified in a case-finding strategy from 55.3% to 85.6%. However, in a recent Chinese population with mean maternal age at pregnancy of 26.6 years, testing using only high-risk criteria missed 82.4% of women with subclinical hypothyroidism, 28.6% of women with overt hyperthyroidism, and 74.6% of women with antithyroid antibodies (596). Member surveys of professional societies have shown that 42.7% of responders in Latin America and 43% in Europe perform universal screening (597,598), whereas only 21% of the Asia-Oceania Thyroid Association members do so (599), and 74% of ATA members support such an approach (600). In the recent 2014 ETA guidelines, the majority of authors recommended universal screening because of the beneficial effects of treatment for overt hypothyroidism, and the fact that the targeted approach will miss a large percentage of women with subclinical hypothyroidism (601). The Spanish Society of Endocrinology and Nutrition (602) and the Indian Thyroid Society (603) have expressed support for universal screening in early pregnancy or preconception. The Indian National Guidelines recommend testing only high-risk women (604). Finally, the American Society for Reproductive Medicine recommends TSH testing in all infertile women attempting pregnancy and in high-risk women in early pregnancy (605).

Together, these data provide a perplexing dilemma. Studies strongly suggest an increase in pregnancy loss risk associated with elevated maternal TSH concentrations, especially when elevated TPOAb are detected. Similarly, thyroid dysfunction is a prevalent condition that can be diagnosed with readily available and inexpensive tests. However, the effectiveness of LT4 therapy has not yet been conclusively demonstrated. Importantly, many have argued that screening for thyroid dysfunction must occur very early in pregnancy (e.g., 4–7 weeks of gestation) to maximize potential benefits of LT4 treatment upon pregnancy loss rates and possibly neurocognitive development. The largest prospective screening studies thus far have provided data most translatable to typical pregnancy care currently provided

worldwide, with initial evaluation between 10 and 15 weeks of gestation. This is important to consider because the feasibility of any screening earlier in gestation is unclear.

Further uncertainty surrounds the choice of biochemical test to use for any screening mandate. Most studies identify two dichotomous hypothyroid populations detected in early pregnancy—those with elevated maternal TSH concentrations, but separately from those with decreased FT4 concentrations. Such populations rarely overlap. These findings suggest that the most effective screening strategies may require multimodal testing beyond just TSH evaluation. And, as mentioned above, detection of elevated TPOAb concentrations may ultimately prove the strongest risk factor for adverse outcomes, though no RCT to date has used this as the sole initial testing approach. Furthermore, the only study to date to target treatment of pregnant women with low FT4 concentrations demonstrated no benefit (306).

Therefore, while acknowledging an impressive amount of retrospective data associating thyroid dysfunction with pregnancy harm, the above uncertainties preclude the task force from recommending for or against a universal screening mandate. The outcome of future studies will prove critical for expanding our knowledge, and the task force acknowledged the wealth of data which will become available in the years ahead. In coming to this conclusion, the task force noted that the majority of patients identified through any universal screening process have TSH concentrations between 2.5 and 5.0 mU/L—a population in whom a treatment benefit is not well established. Furthermore, such a strategy could have detrimental effects, labeling many patients with a biochemical abnormality, and in many cases leading to initiation of possibly inappropriate long-term treatment.

One task force member (CD) dissented from this recommendation, feeling that universal testing for maternal TSH and anti-TPO antibodies soon after pregnancy confirmation is warranted given the existing support for the obstetrical benefits of treatment, with minimal risk of harm with appropriate monitoring. There might be an additional benefit of treatment on fetal neurodevelopment, as shown in animal studies. Unfortunately, the timing of initiation of LT4 treatment after completion of the first trimester in the RCTs that examine child IQ as the outcome is a very serious design flaw that biases these studies to producing negative results, since they miss the relevant window of opportunity to influence neurodevelopment. Nonetheless, universal screening would allow identification and treatment of all women with overt hypothyroidism, which is prevalent in 0.3%–0.5% of pregnant women (259,260,583) and asymptomatic in about 70% of patients (606). It would also allow identification and selective treatment of women with autoimmune subclinical hypothyroidism, and appropriate close monitoring of euthyroid women with isolated thyroid autoimmunity, some of whom will develop hypothyroidism during gestation or PPT.

The task force uniformly recommends that health care providers identify all newly pregnant women at high risk for thyroid disease. This would include women with a history of thyroid dysfunction, symptoms or signs of thyroid dysfunction, presence of a goiter, and known thyroid antibody positivity. Other risk factors for thyroid disease include age >30 years (145,607), history of diabetes mellitus type 1 (608) or other autoimmune disorders (609), history of pregnancy loss, preterm delivery or infertility (206,208), history of head or

neck radiation (610) or prior thyroid surgery (611,612), family history of autoimmune thyroid disease or thyroid dysfunction (613,614), morbid obesity (615,616), use of amiodarone (617), lithium (392), or recent administration of iodinated radiologic contrast (618), two or more prior pregnancies (619), and residing in area of moderate to severe iodine deficiency (620). In these women, measurement of serum TSH concentration should be performed as soon as pregnancy is confirmed, with reflex anti-TPOAb if TSH is 2.5–10 mU/L (Fig. 1).

While screening women for thyroid disease preconception may also prove beneficial, there are currently no data to support such an approach, and the process of testing such a high volume of women, the majority of whom will not become pregnant, seems impractical. Universal screening for TPOAb in early pregnancy or possibly preconception may also prove an attractive alternative, but it warrants further investigation. The high prevalence of anti-TPO positivity (up to 17% in reproductive age women), the extensive findings demonstrating increased risks in the anti-TPOAb-positive population, and the fact that this test would also identify women at risk for developing hypothyroidism during gestation (20%) (147) and PPT (30–50%) (621), make this test an attractive theoretical consideration. However, no data support this testing algorithm at present.

Question 111: Should women be universally tested for thyroid function before or during pregnancy?

■ **RECOMMENDATION 93**

There is insufficient evidence to recommend for or against universal screening for abnormal TSH concentrations in early pregnancy.

No recommendation, insufficient evidence.

■ **RECOMMENDATION 94**

There is insufficient evidence to recommend for or against universal screening for abnormal TSH concentrations preconception, with the exception of women planning assisted reproduction or those known to have TPOAb positivity.

No recommendation, insufficient evidence.

■ **RECOMMENDATION 95**

Universal screening to detect low FT4 concentrations in pregnant women is not recommended.

Weak recommendation, moderate-quality evidence.

■ **RECOMMENDATION 96**

All pregnant women should be verbally screened at the initial prenatal visit for any history of thyroid dysfunction, and prior or current use of either thyroid hormone (LT4) or antithyroid medications (MMI, CM, or PTU).

Strong recommendation, high-quality evidence.

■ **RECOMMENDATION 97**

All patients seeking pregnancy, or newly pregnant, should undergo clinical evaluation. If any of the following risk factors are identified, testing for serum TSH is recommended:




FIG. 1. Testing for thyroid dysfunction in pregnancy. ULRR, upper limit of the reference range.

1. A history of hypothyroidism/hyperthyroidism or current symptoms/signs of thyroid dysfunction
2. Known thyroid antibody positivity or presence of a goiter
3. History of head or neck radiation or prior thyroid surgery
4. Age >30 years
5. Type 1 diabetes or other autoimmune disorders
6. History of pregnancy loss, preterm delivery, or infertility
7. Multiple prior pregnancies (≥ 2)
8. Family history of autoimmune thyroid disease or thyroid dysfunction
9. Morbid obesity (BMI ≥ 40 kg/m²)
10. Use of amiodarone or lithium, or recent administration of iodinated radiologic contrast
11. Residing in an area of known moderate to severe iodine insufficiency

Strong recommendation, moderate-quality evidence.

XIV. FUTURE RESEARCH DIRECTIONS

In developing the Guidelines, the task force frequently struggled with the paucity of high-quality double-blinded placebo-controlled trials in the field of thyroid and pregnancy. In fact, only a minority of the 97 recommendations (24%) in the present Guidelines were graded at the highest American College of Physicians Grading System level of evidence. The Guidelines task force identified topics for future research that will be critical in resolving many of the

unanswered questions in the field of thyroid and pregnancy. Of concern to the task force is that most of the double-blind placebo-controlled studies either recently completed, or presently underway, began screening and intervention after the first trimester. As such, these studies will not be able to address the impact of LT4 treatment in the first trimester in women with subclinical hypothyroidism, isolated hypothyroxinemia, or thyroid Ab positivity on the mother and developing fetus. A trial that screens women preconception and then randomizes women with subclinical hypothyroidism, isolated hypothyroxinemia, and isolated thyroid autoimmunity (with normal TSH) to either a treatment or no treatment arm is needed. The task force is aware of the difficulties inherent in performing such a trial, and the ethical challenges to be faced. Nevertheless, we believe that such a trial is feasible, can be ethically performed with appropriate study design and safeguards, and will yield invaluable information related to the optimal care of the pregnant women and the developing fetus. Other areas for future research include:

- A study evaluating the impact of iodine supplementation in pregnant women with the mildest form of iodine deficiency (median UICs 100–150 μ g/L).
- A RCT of early LT4 intervention (at 4–8 weeks) in women with either subclinical hypothyroidism or isolated hypothyroxinemia to determine effects on child IQ.
- A study focused on the effects of iodine supplementation during lactation on infant thyroid function and cognition.
- A study to determine safe upper limits for iodine ingestion in pregnancy and lactation.

- A comprehensive study to assess the iodine status of pregnant and lactating women in the United States.
- A trial assessing the optimal targeted FT4 level in pregnant women treated for hyperthyroidism.
- Another well powered, prospective, randomized interventional trial of LT4 in euthyroid patients who are anti-TPOAb positive for the prevention of miscarriage and preterm delivery.
- A study to evaluate the impact of LT4 therapy in euthyroid thyroid Ab-positive women with recurrent pregnancy loss.
- Basic and clinical studies aimed at elucidating the mechanisms underlying thyroid Ab-associated adverse pregnancy outcomes.
- Studies examining the effects of TgAb on pregnancy outcomes.
- A study investigating the best criteria that can be used to predict which patients with hyperthyroidism can safely tapered off antithyroid medication in the first trimester.
- A study evaluating the safest timing of administration of the different ATDs for management of hyperthyroidism in pregnancy.
- Novel ways to differentiate fetal hyperthyroidism from fetal hypothyroidism when a fetal goiter is detected.

ACKNOWLEDGMENTS

The task force wishes to thank Ms. Bobbi Smith, Executive Director, ATA, and the rest of the ATA staff for their constant help and support, as well as Ms. Sheri Slaughter for her substantial assistance in compiling meeting notes, updating drafts, and final manuscript preparation. We thank Dr. Tim I. Korevaar (Erasmus University, the Netherlands) for his help in compiling Tables 4–7. We would like to thank the ATA members who responded to our survey in preparation for this iteration of the guidelines as well as manuscript review prior to journal submission. These guidelines were funded by the ATA without support from any commercial sources. The following groups reviewed and endorsed the final document: American Association of Clinical Endocrinologists (AACE); American Radium Society (ARS); Brazilian Society of Head and Neck Surgery (BSHNS); Graves' Disease and Thyroid Foundation; International Association of Endocrine Surgeons (IAES); Japan Thyroid Association; The Endocrine Society of Australia (ESA); ThyCa: Thyroid Cancer Survivors' Association, Inc.; and the Thyroid Federation International. The Pediatric Endocrine Society affirms the value of these guidelines.

AUTHOR DISCLOSURE STATEMENT

These guidelines were funded by the ATA without support from any commercial sources.

CD, GB, HC, JL, PL, SM, SS, and WG have no significant financial or competing interests to disclose.

EP has been a consultant for the Scientific Consulting Company GmbH.

EA has been a consultant for Veracyte.

RB has received grant and/or research support from Verastis, Lilly, Novo Nordisk and Genentech.

RP has received a speaker honoraria from Genzyme and Ipsen as well as serving on the advisory boards of Genzyme, Veracyte and Eisai. Additionally, as patent holder for a test to predict thyroid dysfunction during pregnancy, he recused himself from review and approval of the Screening for Thyroid Dysfunction Before or During Pregnancy (Section XIII).

REFERENCES

1. Stagnaro-Green A, Abalovich M, Alexander E, Azizi F, Mestman J, Negro R, Nixon A, Pearce EN, Soldin OP, Sullivan S, Wiersinga W, American Thyroid Association Taskforce on Thyroid Disease During Pregnancy and Postpartum 2011 Guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and postpartum. *Thyroid* **21**:1081–1125.
2. Qaseem A, Snow V, Owens DK, Shekelle P 2010 The development of clinical practice guidelines and guidance statements of the American College of Physicians: summary of methods. *Ann Intern Med* **153**:194–199.
3. van Raaij JM, Vermaat-Miedema SH, Schonk CM, Peek ME, Hautvast JG 1987 Energy requirements of pregnancy in The Netherlands. *Lancet* **2**:953–955.
4. Glinoer D 1997 The regulation of thyroid function in pregnancy: pathways of endocrine adaptation from physiology to pathology. *Endocr Rev* **18**:404–433.
5. Weeke J, Dybkjaer L, Granlie K, Eskjaer Jensen S, Kjaerulff E, Laurberg P, Magnusson B 1982 A longitudinal study of serum TSH, and total and free iodothyronines during normal pregnancy. *Acta Endocrinologica* **101**:531.
6. Baloch Z, Carayon P, Conte-Devolx B, Demers LM, Feldt-Rasmussen U, Henry JF, LiVosli VA, Niccoli-Sire P, John R, Ruf J, Smyth PP, Spencer CA, Stockigt JR, Guidelines Committee, National Academy of Clinical Biochemistry 2003 Laboratory medicine practice guidelines. Laboratory support for the diagnosis and monitoring of thyroid disease. *Thyroid* **13**:3–126.
7. Soldin OP, Tractenberg RE, Hollowell JG, Jonklaas J, Janicic N, Soldin SJ 2004 Trimester-specific changes in maternal thyroid hormone, thyrotropin, and thyroglobulin concentrations during gestation: trends and associations across trimesters in iodine sufficiency. *Thyroid* **14**:1084–1090.
8. Kahric-Janicic N, Soldin SJ, Soldin OP, West T, Gu J, Jonklaas J 2007 Tandem mass spectrometry improves the accuracy of free thyroxine measurements during pregnancy. *Thyroid* **17**:303–311.
9. Orito Y, Oku H, Kubota S, Amino N, Shimogaki K, Hata M, Manki K, Tanaka Y, Sugino S, Ueta M, Kawakita K, Nunotani T, Tatsumi N, Ichihara K, Miyauchi A, Miyake M 2009 Thyroid function in early pregnancy in Japanese healthy women: relation to urinary iodine excretion, emesis, and fetal and child development. *J Clin Endocrinol Metab* **94**:1683–1688.
10. Roti E, Gardini E, Minelli R, Bianconi L, Flisi M 1991 Thyroid function evaluation by different commercially available free thyroid hormone measurement kits in term pregnant women and their newborns. *J Endocrinol Invest* **14**:1–9.
11. Sapin R, D'Herbomez M, Schlienger JL 2004 Free thyroxine measured with equilibrium dialysis and nine immunoassays decreases in late pregnancy. *Clin Lab* **50**:581–584.

12. Haddow JE, Knight GJ, Palomaki GE, McClain MR, Pulkkinen AJ 2004 The reference range and within-person variability of thyroid stimulating hormone during the first and second trimesters of pregnancy. *J Med Screen* **11**:170–174.
13. Panesar NS, Li CY, Rogers MS 2001 Reference intervals for thyroid hormones in pregnant Chinese women. *Ann Clin Biochem* **38**:329–332.
14. Lockwood CM, Grenache DG, Gronowski AM 2009 Serum human chorionic gonadotropin concentrations greater than 400,000 IU/L are invariably associated with suppressed serum thyrotropin concentrations. *Thyroid* **19**:863–868.
15. De Groot L, Abalovich M, Alexander EK, Amino N, Barbour L, Cobin RH, Eastman CJ, Lazarus JH, Luton D, Mandel SJ, Mestman J, Rovet J, Sullivan S 2012 Management of thyroid dysfunction during pregnancy and postpartum: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* **97**:2543–2565.
16. Yan YQ, Dong ZL, Dong L, Wang FR, Yang XM, Jin XY, Lin LX, Sun YN, Chen ZP 2011 Trimester- and method-specific reference intervals for thyroid tests in pregnant Chinese women: methodology, euthyroid definition and iodine status can influence the setting of reference intervals. *Clin Endocrinol (Oxf)* **74**:262–269.
17. Li C, Shan Z, Mao J, Wang W, Xie X, Zhou W, Li C, Xu B, Bi L, Meng T, Du J, Zhang S, Gao Z, Zhang X, Yang L, Fan C, Teng W 2014 Assessment of thyroid function during first-trimester pregnancy: what is the rational upper limit of serum TSH during the first trimester in Chinese pregnant women? *J Clin Endocrinol Metab* **99**:73–79.
18. Marwaha RK, Chopra S, Gopalakrishnan S, Sharma B, Kanwar RS, Sastry A, Singh S 2008 Establishment of reference range for thyroid hormones in normal pregnant Indian women. *BJOG* **115**:602–606.
19. Korevaar TI, Schalekamp-Timmermans S, de Rijke YB, Visser WE, Visser W, de Muinck Keizer-Schrama SM, Hofman A, Ross HA, Hooijkaas H, Tiemeier H, Bongers-Schokking JJ, Jaddoe VW, Visser TJ, Steegers EA, Medici M, Peeters RP 2013 Hypothyroxinemia and TPO-antibody positivity are risk factors for premature delivery: the generation R study. *J Clin Endocrinol Metab* **98**:4382–4390.
20. Moon HW, Chung HJ, Park CM, Hur M, Yun YM 2015 Establishment of trimester-specific reference intervals for thyroid hormones in Korean pregnant women. *Ann Lab Med* **35**:198–204.
21. Casey B. Effect of treatment of maternal subclinical hypothyroidism or hypothyroxinemia on IQ in offspring. [Abstract]. Available at: [www.ajog.org/article/S0002-9378\(15\)01319-8/pdf](http://www.ajog.org/article/S0002-9378(15)01319-8/pdf) (accessed February 9, 2017).
22. Casey BM, Dashe JS, Wells CE, McIntire DD, Leveno KJ, Cunningham FG 2006 Subclinical hyperthyroidism and pregnancy outcomes. *Obstet Gynecol* **107**:337–341.
23. Thienpont LM, Van Uytvanghe K, Beastall G, Faix JD, Ieiri T, Miller WG, Nelson JC, Ronin C, Ross HA, Thijssen JH, Toussaint B, IFCC Working Group on Standardization of Thyroid Function Tests 2010 Report of the IFCC Working Group for Standardization of Thyroid Function Tests; Part 1: Thyroid-stimulating hormone. *Clin Chem* **56**:902–911.
24. Bestwick JP, John R, Maina A, Guaraldo V, Joomun M, Wald NJ, Lazarus JH 2014 Thyroid stimulating hormone and free thyroxine in pregnancy: expressing concentrations as multiples of the median (MoMs). *Clin Chim Acta* **430**:33–37.
25. Toft AD, Beckett GJ 2005 Measuring serum thyrotropin and thyroid hormone and assessing thyroid hormone transport. In: Braverman, LE, Utiger, RD (eds) *Werner & Ingbar's The Thyroid: A Fundamental and Clinical Text*. Ninth edition. Lippincott, Williams & Wilkins, Philadelphia, PA, pp 329–344.
26. Anckaert E, Poppe K, Van Uytvanghe K, Schiettecatte J, Foulon W, Thienpont LM 2010 FT4 immunoassays may display a pattern during pregnancy similar to the equilibrium dialysis ID-LC/tandem MS candidate reference measurement procedure in spite of susceptibility towards binding protein alterations. *Clin Chim Acta* **411**:1348–1353.
27. Sapin R, d'Herbomez M 2003 Free thyroxine measured by equilibrium dialysis and nine immunoassays in sera with various serum thyroxine-binding capacities. *Clin Chem* **49**:1531–1535.
28. Negro R, Formoso G, Mangieri T, Pezzarossa A, Dazzi D, Hassan H 2006 Levothyroxine treatment in euthyroid pregnant women with autoimmune thyroid disease: effects on obstetrical complications. *J Clin Endocrinol Metab* **91**:2587–2591.
29. Lee RH, Spencer CA, Mestman JH, Miller EA, Petrovic I, Braverman LE, Goodwin TM 2009 Free T4 immunoassays are flawed during pregnancy. *Am J Obstet Gynecol* **200**:260.e1–260.e6.
30. Stockigt J 2003 Assessment of thyroid function: towards an integrated laboratory–clinical approach. *Clin Biochem Rev* **24**:109–122.
31. Yue B, Rockwood AL, Sandrock T, La'ulu SL, Kushnir MM, Meikle AW 2008 Free thyroid hormones in serum by direct equilibrium dialysis and online solid-phase extraction–liquid chromatography/tandem mass spectrometry. *Clin Chem* **54**:642–651.
32. Thienpont LM, Van Uytvanghe K, Beastall G, Faix JD, Ieiri T, Miller WG, Nelson JC, Ronin C, Ross HA, Thijssen JH, Toussaint B, IFCC Working Group on Standardization of Thyroid Function Tests 2010 Report of the IFCC Working Group for Standardization of Thyroid Function Tests; part 2: free thyroxine and free triiodothyronine. *Clin Chem* **56**:912–920.
33. Glinioer D 2007 The importance of iodine nutrition during pregnancy. *Public Health Nutr* **10**:1542–1546.
34. Liberman CS, Pino SC, Fang SL, Braverman LE, Emerson CH 1998 Circulating iodide concentrations during and after pregnancy. *J Clin Endocrinol Metab* **83**:3545–3549.
35. Brander L, Als C, Buess H, Haldimann F, Harder M, Hanggi W, Herrmann U, Lauber K, Niederer U, Zurcher T, Burgi U, Gerber H 2003 Urinary iodine concentration during pregnancy in an area of unstable dietary iodine intake in Switzerland. *J Endocrinol Invest* **26**:389–396.
36. Azizi F, Smyth P 2009 Breastfeeding and maternal and infant iodine nutrition. *Clin Endocrinol (Oxf)* **70**:803–809.
37. Andersen S, Karmisholt J, Pedersen KM, Laurberg P 2008 Reliability of studies of iodine intake and recommendations for number of samples in groups and in individuals. *Br J Nutr* **99**:813–818.
38. König F, Andersson M, Hotz K, Aeberli I, Zimmermann MB 2011 Ten repeat collections for urinary iodine from spot samples or 24-hour samples are needed to reliably estimate individual iodine status in women. *J Nutr* **141**:2049–2054.
39. Berghout A, Wiersinga W 1998 Thyroid size and thyroid function during pregnancy: an analysis. *Eur J Endocrinol* **138**:536–542.

40. Sahin SB, Ogullar S, Ural UM, Ilkkilic K, Metin Y, Ayaz T 2014 Alterations of thyroid volume and nodular size during and after pregnancy in a severe iodine-deficient area. *Clin Endocrinol (Oxf)* **81**:762–768.
41. Delange FM, Dunn JT 2005 Iodine deficiency. In: Braverman LE, Utiger RD (eds) *Werner and Ingbar's The Thyroid: A Fundamental and Clinical Text*. Ninth edition. Lippincott, Williams and Wilkins, Philadelphia, pp 264–288.
42. World Health Organization/International Council for the Control of the Iodine Deficiency Disorders/United Nations Children's Fund (WHO/ICCIDD/UNICEF) 2007 Assessment of the iodine deficiency disorders and monitoring their elimination. World Health Organization, Geneva.
43. Vermiglio F, Lo Presti VP, Castagna MG, Violi MA, Moleti M, Finocchiaro MD, Mattina F, Artemisia A, Trimarchi F 1999 Increased risk of maternal thyroid failure with pregnancy progression in an iodine deficient area with major iodine deficiency disorders. *Thyroid* **9**:19–24.
44. de Escobar GM, Obregon MJ, del Rey FE 2007 Iodine deficiency and brain development in the first half of pregnancy. *Public Health Nutr* **10**:1554–1570.
45. Costeira MJ, Oliveira P, Santos NC, Ares S, Saenz-Rico B, Morreale de Escobar G, Palha JA 2011 Psychomotor development of children from an iodine-deficient region. *J Pediatr* **159**:447–453.
46. Zimmermann MB 2012 The effects of iodine deficiency in pregnancy and infancy. *Paediatr Perinat Epidemiol* **26**(Suppl 1):108–117.
47. Moreno-Reyes R, Glinoe D, Van Oyen H, Vandevijvere S 2013 High prevalence of thyroid disorders in pregnant women in a mildly iodine-deficient country: a population-based study. *J Clin Endocrinol Metab* **98**:3694–3701.
48. Olivares JL, Olivi GI, Verdasco C, Ortiz VA, Mayer MA, Cresto JC 2012 Low iodine intake during pregnancy: relationship to placental development and head circumference in newborn. *Endocrinol Nutr* **59**:326–330.
49. Lean MI, Lean ME, Yajnik CS, Bhat DS, Joshi SM, Raut DA, Lubree HG, Combet E 2014 Iodine status during pregnancy in India and related neonatal and infant outcomes. *Public Health Nutr* **17**:1353–1362.
50. Vermiglio F, Lo Presti VP, Moleti M, Sidoti M, Tortorella G, Scaffidi G, Castagna MG, Mattina F, Violi MA, Crisa A, Artemisia A, Trimarchi F 2004 Attention deficit and hyperactivity disorders in the offspring of mothers exposed to mild-moderate iodine deficiency: a possible novel iodine deficiency disorder in developed countries. *J Clin Endocrinol Metab* **89**:6054–6060.
51. Bath SC, Steer CD, Golding J, Emmett P, Rayman MP 2013 Effect of inadequate iodine status in UK pregnant women on cognitive outcomes in their children: results from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Lancet* **382**:331–337.
52. Hynes KL, Otahal P, Hay I, Burgess JR 2013 Mild iodine deficiency during pregnancy is associated with reduced educational outcomes in the offspring: 9-year follow-up of the gestational iodine cohort. *J Clin Endocrinol Metab* **98**:1954–1962.
53. van Mil NH, Tiemeier H, Bongers-Schokking JJ, Ghasabian A, Hofman A, Hooijkaas H, Jaddoe VW, de Muinck Keizer-Schrama SM, Steegers EA, Visser TJ, Visser W, Ross HA, Verhulst FC, de Rijke YB, Steegers-Theunissen RP 2012 Low urinary iodine excretion during early pregnancy is associated with alterations in executive functioning in children. *J Nutr* **142**:2167–2174.
54. Santiago P, Velasco I, Muela JA, Sánchez B, Martínez J, Rodríguez A, Berrio M, Gutierrez-Repiso C, Carreira M, Moreno A, García-Fuentes E, Soriguer F 2013 Infant neurocognitive development is independent of the use of iodised salt or iodine supplements given during pregnancy. *Br J Nutr* **110**:831–839.
55. Trumpff C, De Schepper J, Tafforeau J, Van Oyen H, Vanderfaeillie J, Vandevijvere S 2013 Mild iodine deficiency in pregnancy in Europe and its consequences for cognitive and psychomotor development of children: a review. *J Trace Elem Med Biol* **27**:174–183.
56. Soldin OP, Soldin SJ, Pezzullo JC 2003 Urinary iodine percentile ranges in the United States. *Clin Chim Acta* **328**:185–190.
57. Caldwell KL, Jones R, Hollowell JG 2005 Urinary iodine concentration: United States National Health And Nutrition Examination Survey 2001–2002. *Thyroid* **15**:692–699.
58. Caldwell KL, Makhmudov AA, Ely EK, Jarrett JM, Henahan D, Jones RL 2009 Iodine status of the U.S. population, NHANES 2005–2006. *Thyroid* **19**:S26–S27.
59. Caldwell KL, Miller GA, Wang RY, Jain RB, Jones RL 2008 Iodine status of the U.S. population, National Health and Nutrition Examination Survey 2003–2004. *Thyroid* **18**:1207–1214.
60. Hollowell JG, Staehling NW, Hannon WH, Flanders DW, Gunter EW, Maberly GF, Braverman LE, Pino S, Miller DT, Garbe PL, DeLozier DM, Jackson RJ 1998 Iodine nutrition in the United States. Trends and public health implications: iodine excretion data from National Health and Nutrition Examination Surveys I and III (1971–1974 and 1988–1994). *J Clin Endocrinol Metab* **83**:3401–3408.
61. Caldwell KL, Makhmudov A, Ely E, Jones RL, Wang RY 2011 Iodine status of the U.S. population, National Health and Nutrition Examination Survey, 2005–2006 and 2007–2008. *Thyroid* **21**:419–427.
62. Caldwell KL, Pan Y, Mortensen ME, Makhmudov A, Merrill L, Moye J 2013 Iodine status in pregnant women in the National Children's Study and in U.S. women (15–44 years), National Health and Nutrition Examination Survey 2005–2010. *Thyroid* **23**:927–937.
63. Pearce EN, Leung AM, Blount BC, Bazrafshan HR, He X, Pino S, Valentin-Blasini L, Braverman LE 2007 Breast milk iodine and perchlorate concentrations in lactating Boston-area women. *J Clin Endocrinol Metab* **92**:1673–1677.
64. Kirk AB, Martinelango PK, Tian K, Dutta A, Smith EE, Dasgupta PK 2005 Perchlorate and iodide in dairy and breast milk. *Environ Sci Technol* **39**:2011–2017.
65. Gordon JH, Leung AM, Hale AR, Pearce EN, Braverman LE, He X, Belfort MB, Nelson SM, Brown RS 2014 No difference in urinary iodine concentrations between Boston-area breastfed and formula-fed infants. *Thyroid* **24**:1309–1313.
66. Pearce EN, Andersson M, Zimmermann MB 2013 Global iodine nutrition: where do we stand in 2013? *Thyroid* **23**:523–528.
67. Jaiswal N, Melse-Boonstra A, Sharma SK, Srinivasan K, Zimmermann MB 2015 The iodized salt programme in Bangalore, India provides adequate iodine intakes in pregnant women and more-than-adequate iodine intakes in their children. *Public Health Nutr* **18**:403–413.

68. Majumder A, Jaiswal A, Chatterjee 2014 Prevalence of iodine deficiency among pregnant and lactating women: experience in Kolkata. *Indian J Endocrinol Metab* **18**:486–490.
69. Patey-Pirra S, Keriell-Gascou M, Borson-Chazot F 2014 Benefits and risks of iodine supplementation during pregnancy: a review of observational and experimental studies in mild-to-moderate iodine deficiency areas. *Rev Epidemiol Sante Publique* **62**:65–74.
70. Rydbeck F, Bottai M, Tofail F, Persson LÅ, Kippler M 2014 Urinary iodine concentrations of pregnant women in rural Bangladesh: a longitudinal study. *J Expo Sci Environ Epidemiol* **24**:504–509.
71. Elahi S, Nagra SA 2014 Low maternal iodine intake and early pregnancy hypothyroxinemia: possible repercussions for children. *Indian J Endocrinol Metab* **18**:526–530.
72. Wei Z, Wang W, Zhang J, Zhang X, Jin L, Yu X 2015 Urinary iodine level and its determinants in pregnant women of Shanghai, China. *Br J Nutr* **113**:1427–1432.
73. Shi X, Han C, Li C, Mao J, Wang W, Xie X, Li C, Xu B, Meng T, Du J, Zhang S, Gao Z, Zhang X, Fan C, Shan Z, Teng W 2015 Optimal and safe upper limits of iodine intake for early pregnancy in iodine-sufficient regions: a cross-sectional study of 7,190 pregnant women in China. *J Clin Endocrinol Metab* **100**:1630–1638.
74. Cho YY, Kim HJ, Oh SY, Choi SJ, Lee SY, Joung JY, Jeong DJ, Sohn SY, Chung JH, Roh CR, Kim SW 2016 Iodine status in healthy pregnant women in Korea: a first report. *Eur J Nutr* **55**:469–475.
75. Sadou H, Seyfoulaye A, Malam Alma M, Daouda H 2014 Inadequate status of iodine nutrition among pregnant women residing in three districts of Niamey, the Niger Republic's capital. *Matern Child Nutr* **10**:650–656.
76. Kedir H, Berhane Y, Worku A 2014 Subclinical iodine deficiency among pregnant women in Haramaya district, Eastern Ethiopia: a community-based study. *J Nutr Metab* **2014**:878926.
77. Amouzegar A, Khazan M, Hedayati M, Azizi F 2014 An assessment of the iodine status and the correlation between iodine nutrition and thyroid function during pregnancy in an iodine sufficient area. *Eur J Clin Nutr* **68**:397–400.
78. Ferreira SM, Navarro AM, Magalhães PK, Maciel LM 2014 Iodine insufficiency in pregnant women from the State of São Paulo. *Arq Bras Endocrinol Metabol* **58**:282–287.
79. Jiskra J, Fait T, Bílek R, Krátký J, Bartáková J, Lukáš J, Límanová Z, Telička Z, Zamrazil V, Potluková E 2014 Mild iodine deficiency in women after spontaneous abortions living in an iodine-sufficient area of Czech Republic: prevalence and impact on reproductive health. *Clin Endocrinol (Oxf)* **80**:452–458.
80. Andersen SL, Sørensen LK, Krejbjerg A, Møller M, Laurberg P 2013 Iodine deficiency in Danish pregnant women. *Dan Med J* **60**:A4657.
81. Raverot V, Bournaud C, Sassolas G, Orgiazzi J, Claustat F, Gaucherand P, Mellier G, Claustat B, Borson-Chazot F, Zimmermann M 2012 Pregnant French women living in the Lyon area are iodine deficient and have elevated serum thyroglobulin concentrations. *Thyroid* **22**:522–528.
82. Konrade I, Kalere I, Strele I, Makrecka-Kuka M, Jekabsons A, Tetere E, Veisa V, Gavars D, Rezeberga D, Pīrāgs V, Lejnīeks A, Dambrova M 2015 Iodine deficiency during pregnancy: a national cross-sectional survey in Latvia. *Public Health Nutr* **18**:2990–2997.
83. Brantsæter AL, Abel MH, Haugen M, Meltzer HM 2013 Risk of suboptimal iodine intake in pregnant Norwegian women. *Nutrients* **5**:424–440.
84. Aguayo A, Grau G, Vela A, Aniel-Quiroga A, Espada M, Martul P, Castaño L, Rica I 2013 Urinary iodine and thyroid function in a population of healthy pregnant women in the North of Spain. *J Trace Elem Med Biol* **27**:302–306.
85. Bath SC, Sleeth ML, McKenna M, Walter A, Taylor A, Rayman MP 2014 Iodine intake and status of UK women of childbearing age recruited at the University of Surrey in the winter. *Br J Nutr* **112**:1715–1723.
86. Pharoah PO, Butfield IH, Hetzel BS 1971 Neurological damage to the fetus resulting from severe iodine deficiency during pregnancy. *Lancet* **1**:308–310.
87. O'Donnell KJ, Rakeman MA, Zhi-Hong D, Xue-Yi C, Mei ZY, DeLong N, Brenner G, Tai M, Dong W, DeLong GR 2002 Effects of iodine supplementation during pregnancy on child growth and development at school age. *Dev Med Child Neurol* **44**:76–81.
88. Fierro-Benitez R, Cazar R, Stanbury JB, Rodriguez P, Garces F, Fierro-Renoy F, Estrella E 1988 Effects on school children of prophylaxis of mothers with iodized oil in an area of iodine deficiency. *J Endocrinol Invest* **11**:327–335.
89. Cao XY, Jiang XM, Dou ZH, Rakeman MA, Zhang M, O'Donnell K, Ma T, Amette K, DeLong N, DeLong GR 1994 Timing of vulnerability of the brain to iodine deficiency in endemic cretinism. *N Engl J Med* **331**:1739–1744.
90. DeLong GR, Leslie PW, Wang SH, Jiang XM, Zhang ML, Rakeman M, Jiang JY, Ma T, Cao XY 1997 Effect on infant mortality of iodination of irrigation water in a severely iodine-deficient area of China. *Lancet* **350**:771–773.
91. Chaouki ML, Benmiloud M 1994 Prevention of iodine deficiency disorders by oral administration of lipiodol during pregnancy. *Eur J Endocrinol* **130**:547–551.
92. Anees M, Anis RA, Yousaf S, Murtaza I, Sultan A, Arslan M, Shahab M 2015 Effect of maternal iodine supplementation on thyroid function and birth outcome in goiter endemic areas. *Curr Med Res Opin Feb* **13**:1–8.
93. Bouhouch RR, Bouhouch S, Cherkaoui M, Aboussad A, Stinca S, Haldimann M, Andersson M, Zimmermann MB 2014 Direct iodine supplementation of infants versus supplementation of their breastfeeding mothers: a double-blind, randomised, placebo-controlled trial. *Lancet Diabetes Endocrinol* **2**:197–209.
94. Antonangeli L, Maccherini D, Cavaliere R, Di Giulio C, Reinhardt B, Pinchera A, Aghini-Lombardi F 2002 Comparison of two different doses of iodide in the prevention of gestational goiter in marginal iodine deficiency: a longitudinal study. *Eur J Endocrinol* **147**:29–34.
95. Berbel P, Mestre JL, Santamaria A, Palazon I, Franco A, Graells M, Gonzalez-Torga A, de Escobar GM 2009 Delayed neurobehavioral development in children born to pregnant women with mild hypothyroxinemia during the first month of gestation: the importance of early iodine supplementation. *Thyroid* **19**:511–519.
96. Glinoe D, De Nayer P, Delange F, Lemone M, Toppet V, Spehl M, Grun JP, Kinthaert J, Lejeune B 1995 A randomized trial for the treatment of mild iodine deficiency during pregnancy: maternal, neonatal effects. *J Clin Endocrinol Metab* **80**:258–269.

97. Liesenkotter KP, Gopel W, Bogner U, Stach B, Gruters A 1996 Earliest prevention of endemic goiter by iodine supplementation during pregnancy. *Eur J Endocrinol* **134**: 443–448.
98. Nohr SB, Laurberg P 2000 Opposite variations in maternal and neonatal thyroid function induced by iodine supplementation during pregnancy. *J Clin Endocrinol Metab* **85**:623–627.
99. Pedersen KM, Laurberg P, Iversen E, Knudsen PR, Gregersen HE, Rasmussen OS, Larsen KR, Eriksen GM, Johannessen PL 1993 Amelioration of some pregnancy-associated variations in thyroid function by iodine supplementation. *J Clin Endocrinol Metab* **77**:1078–1083.
100. Romano R, Jannini EA, Pepe M, Grimaldi A, Olivieri M, Spennati P, Cappa F, D'Armiento M 1991 The effects of iodophrophylaxis on thyroid size during pregnancy. *Am J Obstet Gynecol* **164**:482–485.
101. Velasco I, Carreira M, Santiago P, Muela JA, Garcia-Fuentes E, Sanchez-Munoz B, Garriga MJ, Gonzalez-Fernandez MC, Rodriguez A, Caballero FF, Machado A, Gonzalez-Romero S, Anarte MT, Soriguer F 2009 Effect of iodine prophylaxis during pregnancy on neurocognitive development of children during the first two years of life. *J Clin Endocrinol Metab* **94**:3234–3241.
102. Sukkhajaiwaratkul D, Mahachoklertwattana P, Poomthavorn P, Panburana P, Chailurkit LO, Khlairit P, Pongratanakul S 2014 Effects of maternal iodine supplementation during pregnancy and lactation on iodine status and neonatal thyroid-stimulating hormone. *J Perinatol* **34**:594–598.
103. Rebagliato M, Murcia M, Alvarez-Pedrerol M, Espada M, Fernández-Somoano A, Lertxundi N, Navarrete-Muñoz EM, Fornis J, Aranbarri A, Llop S, Julvez J, Tardón A, Ballester F 2013 Iodine supplementation during pregnancy and infant neuropsychological development. INMA Mother and Child Cohort Study. *Am J Epidemiol* **177**: 944–953.
104. Vandevijvere S, Amsalkhir S, Mourri AB, Van Oyen H, Moreno-Reyes R 2013 Iodine deficiency among Belgian pregnant women not fully corrected by iodine-containing multivitamins: a national cross-sectional survey. *Br J Nutr* **109**:2276–2284.
105. Taylor PN, Okosieme OE, Dayan CM, Lazarus JH 2013 Therapy of endocrine disease: impact of iodine supplementation in mild-to-moderate iodine deficiency: systematic review and meta-analysis. *Eur J Endocrinol* **170**: R1–R15.
106. Zhou SJ, Anderson AJ, Gibson RA, Makrides M 2013 Effect of iodine supplementation in pregnancy on child development and other clinical outcomes: a systematic review of randomized controlled trials. *Am J Clin Nutr* **98**:1241–1254.
107. Bath SC, Rayman MP 2015 A review of the iodine status of UK pregnant women and its implications for the offspring. *Environ Geochem Health* **37**:619–629.
108. Stagnaro-Green A, Pearce EN 2013 Iodine and pregnancy: a call to action. *Lancet* **382**:292–293.
109. Bougma K, Aboud FE, Harding KB, Marquis GS 2013 Iodine and mental development of children 5 years old and under: a systematic review and meta-analysis. *Nutrients* **5**:1384–1416.
110. Menéndez Torre E, Delgado Alvarez E, Rabal Artal A, Suárez Gutiérrez L, Rodríguez Caballero MG, Ares Blanco J, Díaz Naya L, Fernández Fernández JC 2014 Iodine nutrition in pregnant women from Oviedo area. Is iodine supplementation necessary? *Endocrinol Nutr* **61**:404–409.
111. Trumbo P, Yates AA, Schlicker S, Poos M 2001 Dietary reference intakes: vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. *J Am Diet Assoc* **101**: 294–301.
112. Maalouf J, Barron J, Gunn JP, Yuan K, Perrine CG, Cogswell ME 2015 Iodized salt sales in the United States. *Nutrients* **7**:1691–1695.
113. Galton DM, Petersson LG, Erb HN 1986 Milk iodine residues in herds practicing iodophor premilking teat disinfection. *J Dairy Sci* **69**:267–271.
114. Conrad LM 3rd, Hemken RW 1978 Milk iodine as influenced by an iodophor teat dip. *J Dairy Sci* **61**:776–780.
115. Pearce EN, Pino S, He X, Bazrafshan HR, Lee SL, Braverman LE 2004 Sources of dietary iodine: bread, cows' milk, and infant formula in the Boston area. *J Clin Endocrinol Metab* **89**:3421–3424.
116. Haldimann M, Alt A, Blanc A, Blondeau K 2005 Iodine content of food groups. *J Food Comp Anal* **18**:461–471.
117. Public Health Committee of the American Thyroid Association, Becker DV, Braverman LE, Delange F, Dunn JT, Franklyn JA, Hollowell JG, Lamm SH, Mitchell ML, Pearce E, Robbins J, Rovet JF 2006 Iodine supplementation for pregnancy and lactation—United States and Canada: recommendations of the American Thyroid Association. *Thyroid* **16**:949–951.
118. Obican SG, Jahnke GD, Soldin OP, Scialli AR 2012 Teratology public affairs committee position paper: iodine deficiency in pregnancy. *Birth Defects Res A Clin Mol Teratol* **94**:677–682.
119. Council on Environmental Health, Rogan WJ, Paulson JA, Baum C, Brock-Utne AC, Brumberg HL, Campbell CC, Lanphear BP, Lowry JA, Osterhoudt KC, Sandel MT, Spanier A, Trasande L 2014 Iodine deficiency, pollutant chemicals, and the thyroid: new information on an old problem. *Pediatrics* **133**:1163–1166.
120. Leung AM, Lamar A, He X, Braverman LE, Pearce EN 2011 Iodine status and thyroid function of Boston-area vegetarians and vegans. *J Clin Endocrinol Metab* **96**: E1303–E1307.
121. Gregory CO, Serdula MK, Sullivan KM 2009 Use of supplements with and without iodine in women of child-bearing age in the United States. *Thyroid* **19**:1019–1020.
122. Leung AM, Pearce EN, Braverman LE 2009 Iodine content of prenatal multivitamins in the United States. *N Engl J Med* **360**:939–940.
123. U.S. Salt Institute 2007 Iodized salt. Available at <http://www.saltinstitute.org/Uses-benefits/Salt-in-Food/Essential-nutrient/Iodized-salt> (accessed September 12, 2011).
124. Pennington JA 1990 A review of iodine toxicity reports. *J Am Diet Assoc* **90**:1571–1581.
125. Wolff J, Chaikoff IL 1948 Plasma inorganic iodide as a homeostatic regulator of thyroid function. *J Biol Chem* **174**:555–564.
126. Eng PH, Cardona GR, Fang SL, Previti M, Alex S, Carasco N, Chin WW, Braverman LE 1999 Escape from the acute Wolff-Chaikoff effect is associated with a decrease in thyroid sodium/iodide symporter messenger ribonucleic acid and protein. *Endocrinology* **140**:3404–3410.
127. Fisher DA, Klein AH 1981 Thyroid development and disorders of thyroid function in the newborn. *N Engl J Med* **304**:702–712.

128. Theodoropoulos T, Braverman LE, Vagenakis AG 1979 Iodide-induced hypothyroidism: a potential hazard during perinatal life. *Science* **205**:502–503.
129. Fang MC, Stafford RS, Ruskin JN, Singer DE 2004 National trends in antiarrhythmic and antithrombotic medication use in atrial fibrillation. *Arch Intern Med* **164**:55–60.
130. Aiba M, Ninomiya J, Furuya K, Arai H, Ishikawa H, Asami S, Takagi A, Ohwada S, Morishita Y 1999 Induction of a critical elevation of povidone-iodine absorption in the treatment of a burn patient: report of a case. *Surg Today* **29**:157–159.
131. Leung AM, Avram AM, Brenner AV, Duntas LH, Ehrenkranz J, Hennessey JV, Lee SL, Pearce EN, Roman SA, Stagnaro-Green A, Sturgis EM, Sundaram K, Thomas MJ, Wexler JA 2015 Potential risks of excess iodine ingestion and exposure: statement by the American Thyroid Association public health committee. *Thyroid* **25**:145–146.
132. Aakre I, Bjørø T, Norheim I, Strand TA, Barikmo I, Henjum S 2015 Excessive iodine intake and thyroid dysfunction among lactating Saharawi women. *J Trace Elem Med Biol* **31**:279–284.
133. Sang Z, Wei W, Zhao N, Zhang G, Chen W, Liu H, Shen J, Liu J, Yan Y, Zhang W 2012 Thyroid dysfunction during late gestation is associated with excessive iodine intake in pregnant women. *J Clin Endocrinol Metab* **97**:E1363–E1369.
134. Chen W, Sang Z, Tan L, Zhang S, Dong F, Chu Z, Wei W, Zhao N, Zhang G, Yao Z, Shen J, Zhang W 2015 Neonatal thyroid function born to mothers living with long-term excessive iodine intake from drinking water. *Clin Endocrinol (Oxf)* **83**:399–404.
135. Medici M, Ghassabian A, Visser W, de Muinck Keizer-Schrama SM, Jaddoe VW, Visser WE, Hooijkaas H, Hofman A, Steegers EA, Bongers-Schokking JJ, Ross HA, Tiemeier H, Visser TJ, de Rijke YB, Peeters RP 2014 Women with high early pregnancy urinary iodine levels have an increased risk of hyperthyroid newborns: the population-based Generation R Study. *Clin Endocrinol (Oxf)* **80**:598–606.
136. Thaker VV, Leung AM, Braverman LE, Brown RS, Levine B 2014 Iodine-induced hypothyroidism in full-term infants with congenital heart disease: more common than currently appreciated? *J Clin Endocrinol Metab* **99**:3521–3526.
137. Ashoor G, Maiz N, Rotas M, Jawdat F, Nicolaides KH 2010 Maternal thyroid function at 11 to 13 weeks of gestation and subsequent fetal death. *Thyroid* **20**:989–993.
138. Benhadi N, Wiersinga WM, Reitsma JB, Vrijkotte TG, van der Wal MF, Bonsel GJ 2007 Ethnic differences in TSH but not in free T4 concentrations or TPO antibodies during pregnancy. *Clin Endocrinol (Oxf)* **66**:765–770.
139. La'ulu SL, Roberts WL 2007 Second-trimester reference intervals for thyroid tests: the role of ethnicity. *Clin Chem* **53**:1658–1664.
140. Abbassi-Ghanavati M, Casey BM, Spong CY, McIntire DD, Halvorson LM, Cunningham FG 2010 Pregnancy outcomes in women with thyroid peroxidase antibodies. *Obstet Gynecol* **116**(2 Pt 1):381–386.
141. Wasserman EE, Nelson K, Rose NR, Eaton W, Pillion JP, Seaberg E, Talor MV, Burek L, Duggan A, Yolken RH 2008 Maternal thyroid autoantibodies during the third trimester and hearing deficits in children: an epidemiologic assessment. *Am J Epidemiol* **167**:701–710.
142. Pearce EN, Oken E, Gillman MW, Lee SL, Magnani B, Platek D, Braverman LE 2008 Association of first-trimester thyroid function test values with thyroperoxidase antibody status, smoking, and multivitamin use. *Endocr Pract* **14**:33–39.
143. Moleti M, Lo Presti VP, Mattina F, Mancuso A, De Vivo A, Giorgianni G, Di Bella B, Trimarchi F, Vermiglio F 2009 Gestational thyroid function abnormalities in conditions of mild iodine deficiency: early screening versus continuous monitoring of maternal thyroid status. *Eur J Endocrinol* **160**:611–617.
144. McElduff A, Morris J 2008 Thyroid function tests and thyroid autoantibodies in an unselected population of women undergoing first trimester screening for aneuploidy. *Aust N Z J Obstet Gynaecol* **48**:478–480.
145. Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, Braverman LE 2002 Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab* **87**:489–499.
146. Unuane D, Velkeniers B, Anckaert E, Schiettecatte J, Tournaye H, Haentjens P, Poppe K 2013 Thyroglobulin autoantibodies: is there any added value in the detection of thyroid autoimmunity in women consulting for fertility treatment? *Thyroid* **23**:1022–1028.
147. Glinoe D, Riahi M, Grün JP, Kintthart J 1994 Risk of subclinical hypothyroidism in pregnant women with asymptomatic autoimmune thyroid disorders. *J Clin Endocrinol Metab* **79**:197–204.
148. Seror J, Amand G, Guibourdenche J, Ceccaldi PF, Luton D 2014 Anti-TPO antibodies diffusion through the placental barrier during pregnancy. *PLoS One* **9**(1):e84647.
149. Gartner R, Gasnier BC, Dietrich JW, Krebs B, Angstwurm MW 2002 Selenium supplementation in patients with autoimmune thyroiditis decreases thyroid peroxidase antibodies concentrations. *J Clin Endocrinol Metab* **87**:1687–1691.
150. Duntas LH, Mantzou E, Koutras DA 2003 Effects of a six month treatment with selenomethionine in patients with autoimmune thyroiditis. *Eur J Endocrinol* **148**:389–393.
151. Mazokopakis EE, Papadakis JA, Papadomanolaki MG, Batistakis AG, Giannakopoulos TG, Protopapadakis EE, Ganotakis ES 2007 Effects of 12 months treatment with L-selenomethionine on serum anti-TPO Levels in Patients with Hashimoto's thyroiditis. *Thyroid* **17**:609–612.
152. Fan Y, Xu S, Zhang H, Cao W, Wang K, Chen G, Di H, Cao M, Liu C 2014 Selenium supplementation for autoimmune thyroiditis: a systematic review and meta-analysis. *Int J Endocrinol* **2014**:904573.
153. Karanikas G, Schuetz M, Kontur S, Duan H, Kommata S, Schoen R, Antoni A, Kletter K, Dudczak R, Willheim M 2008 No immunological benefit of selenium in consecutive patients with autoimmune thyroiditis. *Thyroid* **18**:7–12.
154. Negro R, Greco G, Mangieri T, Pezzarossa A, Dazzi D, Hassan H 2007 The influence of selenium supplementation on postpartum thyroid status in pregnant women with thyroid peroxidase autoantibodies. *J Clin Endocrinol Metab* **92**:1263–1268.
155. Mao J, Pop VJ, Bath SC, Vader HL, Redman CW, Rayman MP 2016 Effect of low-dose selenium on thyroid autoimmunity and thyroid function in UK pregnant women with mild-to-moderate iodine deficiency. *Eur J Nutr* **55**:55–61.
156. Stranges S, Marshall JR, Natarajan R, Donahue RP, Trevisan M, Combs GF, Cappuccio FP, Ceriello A, Reid ME

- 2007 Effects of long-term selenium supplementation on the incidence of type 2 diabetes: a randomized trial. *Ann Intern Med* **147**:217–223.
157. Ellish NJ, Saboda K, O'Connor J, Nasca PC, Stanek EJ, Boyle C 1996 A prospective study of early pregnancy loss. *Hum Reprod* **11**:406–412.
158. Wilcox AJ, Weinberg CR, O'Connor JF, Baird DD, Schlatterer JP, Canfield RE, Armstrong EG, Nisula BC 1988 Incidence of early loss of pregnancy. *N Engl J Med* **319**:189–194.
159. Toth B, Jeschke U, Rogenhofer N, Scholz C, Wurfel W, Thaler CJ, Makrigiannakis A 2010 Recurrent miscarriage: current concepts in diagnosis and treatment. *J Reprod Immunol* **85**:25–32.
160. Dudley DJ 2007 Diabetic-associated stillbirth: incidence, pathophysiology, and prevention. *Clin Perinatol* **34**:611–626, vii.
161. De Vivo A, Mancuso A, Giacobbe A, Moleti M, Maggio Savasta L, De Dominicis R, Priolo AM, Vermiglio F 2010 Thyroid function in women found to have early pregnancy loss. *Thyroid* **20**:633–637.
162. Stagnaro-Green A, Roman SH, Cobin RH, el-Harazy E, Alvarez-Marfany M, Davies TF 1990 Detection of at-risk pregnancy by means of highly sensitive assays for thyroid autoantibodies. *JAMA* **264**:1422–1425.
163. Chen L, Hu R 2011 Thyroid autoimmunity and miscarriage: a meta-analysis. *Clin Endocrinol (Oxf)* **74**:513–519.
164. Thangaratinam S, Tan A, Knox E, Kilby MD, Franklyn J, Coomarasamy A 2011 Association between thyroid autoantibodies and miscarriage and preterm birth: meta-analysis of evidence. *BMJ* **342**:d2616.
165. Lee YL, Ng HP, Lau KS, Liu WM, O WS, Yeung WS, Kung AW 2009 Increased fetal abortion rate in autoimmune thyroid disease is related to circulating TPO autoantibodies in an autoimmune thyroiditis animal model. *Fertil Steril* **91**:2104–2109.
166. Imaizumi M, Pritsker A, Kita M, Ahmad L, Unger P, Davies T 2001 Pregnancy and murine thyroiditis: thyroglobulin immunization leads to fetal loss in specific allogeneic pregnancies. *Endocrinology* **142**:823–829.
167. Matalon ST, Blank M, Levy Y, Carp HJ, Arad A, Burek L, Grunebaum E, Sherer Y, Ornoy A, Refetoff S, Weiss RE, Rose NR, Shoenfeld Y 2003 The pathogenic role of anti-thyroglobulin antibody on pregnancy: evidence from an active immunization model in mice. *Hum Reprod* **18**:1094–1099.
168. Twig G, Shina A, Amital H, Shoenfeld Y 2012 Pathogenesis of infertility and recurrent pregnancy loss in thyroid autoimmunity. *J Autoimmun* **38**:J275–J281.
169. Lee RM, Silver RM 2000 Recurrent pregnancy loss: summary and clinical recommendations. *Semin Reprod Med* **18**:433–440.
170. Baek KH, Lee EJ, Kim YS 2007 Recurrent pregnancy loss: the key potential mechanisms. *Trends Mol Med* **13**:310–317.
171. Irvani AT, Saeedi MM, Pakravesh J, Hamidi S, Abbasi M 2008 Thyroid autoimmunity and recurrent spontaneous abortion in Iran: a case-control study. *Endocr Pract* **14**:458–464.
172. Kutteh WH, Yetman DL, Carr AC, Beck LA, Scott RT Jr 1999 Increased prevalence of antithyroid antibodies identified in women with recurrent pregnancy loss but not in women undergoing assisted reproduction. *Fertil Steril* **71**:843–848.
173. Esplin MS, Branch DW, Silver R, Stagnaro-Green A 1998 Thyroid autoantibodies are not associated with recurrent pregnancy loss. *Am J Obstet Gynecol* **179**(6 Pt 1):1583–1586.
174. Pratt DE, Kaberlein G, Dudkiewicz A, Karande V, Gleicher N 1993 The association of antithyroid antibodies in euthyroid nonpregnant women with recurrent first trimester abortions in the next pregnancy. *Fertil Steril* **60**:1001–1005.
175. Rushworth FH, Backos M, Rai R, Chilcott IT, Baxter N, Regan L 2000 Prospective pregnancy outcome in untreated recurrent miscarriers with thyroid autoantibodies. *Hum Reprod* **15**:1637–1639.
176. Lata K, Dutta P, Sridhar S, Rohilla M, Srinivasan A, Prashad GR, Shah VN, Bhansali A 2013 Thyroid autoimmunity and obstetric outcomes in women with recurrent miscarriage: a case-control study. *Endocr Connect* **2**:118–124.
177. van den Boogaard E, Vissenberg R, Land JA, van Wely M, van der Post JA, Goddijn M, Bisschop PH 2011 Significance of (sub)clinical thyroid dysfunction and thyroid autoimmunity before conception and in early pregnancy: a systematic review. *Hum Reprod Update* **17**:605–619.
178. De Carolis C, Greco E, Guarino MD, Perricone C, Dal Lago A, Giacomelli R, Fontana L, Perricone R 2004 Anti-thyroid antibodies and antiphospholipid syndrome: evidence of reduced fecundity and of poor pregnancy outcome in recurrent spontaneous aborters. *Am J Reprod Immunol* **52**:263–266.
179. Kim NY, Cho HJ, Kim HY, Yang KM, Ahn HK, Thornton S, Park JC, Beaman K, Gilman-Sachs A, Kwak-Kim J 2011 Thyroid autoimmunity and its association with cellular and humoral immunity in women with reproductive failures. *Am J Reprod Immunol* **65**:78–87.
180. Lepoutre T, Debiève F, Gruson D, Daumerie C 2012 Reduction of miscarriages through universal screening and treatment of thyroid autoimmune diseases. *Gynecol Obstet Invest* **74**:265–273.
181. Kiproff DD, Nachtigall RD, Weaver RC, Jacobson A, Main EK, Garovoy MR 1996 The use of intravenous immunoglobulin in recurrent pregnancy loss associated with combined alloimmune and autoimmune abnormalities. *Am J Reprod Immunol* **36**:228–234.
182. Stricker RB, Steinleitner A, Bookoff CN, Weckstein LN, Winger EE 2000 Successful treatment of immunologic abortion with low-dose intravenous immunoglobulin. *Fertil Steril* **73**:536–540.
183. Vaquero E, Lazzarin N, De Carolis C, Valensise H, Moretti C, Ramanini C 2000 Mild thyroid abnormalities and recurrent spontaneous abortion: diagnostic and therapeutic approach. *Am J Reprod Immunol* **43**:204–208.
184. Martin JA, Hamilton BE, Osterman MJK, Curtin SC, Mathews T 2015 Births: final data for 2013. *Vital Statistics Reports*. Available at http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_01.pdf (accessed February 28, 2015).
185. Institute of Medicine (US) Committee on Understanding Premature Birth and Assuring Healthy Outcomes 2007 Preterm Birth: Causes, Consequences, and Prevention. Behrman RE, Butler AS (eds). National Academies Press, Washington, DC.
186. Goldenberg RL, Culhane JF, Iams JD, Romero R 2008 Epidemiology and causes of preterm birth. *Lancet* **371**:75–84.
187. Glinioer D, Soto MF, Bourdoux P, Lejeune B, Delange F, Lemone M, Kinthaert J, Robijn C, Grun JP, de Nayer P

- 1991 Pregnancy in patients with mild thyroid abnormalities: maternal and neonatal repercussions. *J Clin Endocrinol Metab* **73**:421–427.
188. Ghafoor F, Mansoor M, Malik T, Malik MS, Khan AU, Edwards R, Akhtar W 2006 Role of thyroid peroxidase antibodies in the outcome of pregnancy. *J Coll Physicians Surg Pak* **16**:468–471.
189. Iijima T, Tada H, Hidaka Y, Mitsuda N, Murata Y, Amino N 1997 Effects of autoantibodies on the course of pregnancy and fetal growth. *Obstet Gynecol* **90**:364–369.
190. Haddow JE, Cleary-Goldman J, McClain MR, Palomaki GE, Neveux LM, Lambert-Messerlian G, Canick JA, Malone FD, Porter TF, Nyberg DA, Bernstein PS, D'Alton ME; First- and Second-Trimester Risk of Aneuploidy (FaSTER) Research Consortium 2010 Thyroperoxidase and thyroglobulin antibodies in early pregnancy and preterm delivery. *Obstet Gynecol* **116**:58–62.
191. Männistö T, Väärasmäki M, Pouta A, Hartikainen AL, Ruokonen A, Surcel HM, Bloigu A, Järvelin MR, Suvanto-Luukkonen E 2009 Perinatal outcome of children born to mothers with thyroid dysfunction or antibodies: a prospective population-based cohort study. *J Clin Endocrinol Metab* **94**:772–779.
192. Karakosta P, Alegakis D, Georgiou V, Roumeliotaki T, Fthenou E, Vassilaki M, Boumpas D, Castanas E, Kogevinas M, Chatzi L 2012 Thyroid dysfunction and autoantibodies in early pregnancy are associated with increased risk of gestational diabetes and adverse birth outcomes. *J Clin Endocrinol Metab* **97**:4464–4472.
193. Kumru P, Erdogdu E, Arisoy R, Demirci O, Ozkoral A, Ardic C, Ertekin AA, Erdogan S, Ozdemir NN 2015 Effect of thyroid dysfunction and autoimmunity on pregnancy outcomes in low risk population. *Arch Gynecol Obstet* **291**:1047–1054.
194. Negro R 2011 Thyroid autoimmunity and pre-term delivery: brief review and meta-analysis. *J Endocrinol Invest* **34**:155–158.
195. He X, Wang P, Wang Z, He X, Xu D, Wang B 2012 Thyroid antibodies and risk of preterm delivery: a meta-analysis of prospective cohort studies. *Eur J Endocrinol* **167**:455–464.
196. Negro R, Schwartz A, Gismondi R, Tinelli A, Mangieri T, Stagnaro-Green A 2011 Thyroid antibody positivity in the first trimester of pregnancy is associated with negative pregnancy outcomes. *J Clin Endocrinol Metab* **96**:E920–E924.
197. Groer MW, Vaughan JH 2013 Positive thyroid peroxidase antibody titer is associated with dysphoric moods during pregnancy and postpartum. *J Obstet Gynecol Neonatal Nurs* **42**:E26–E32.
198. Kuijpers JL, Vader HL, Drexhage HA, Wiersinga WM, van Son MJ, Pop VJ 2001 Thyroid peroxidase antibodies during gestation are a marker for subsequent depression postpartum. *Eur J Endocrinol* **145**:579–584.
199. Li Y, Shan Z, Teng W, Yu X, Li Y, Fan C, Teng X, Guo R, Wang H, Li J, Chen Y, Wang W, Chawinga M, Zhang L, Yang L, Zhao Y, Hua T 2010 Abnormalities of maternal thyroid function during pregnancy affect neuropsychological development of their children at 25–30 months. *Clin Endocrinol (Oxf)* **72**:825–829.
200. Williams FL, Watson J, Ogston SA, Visser TJ, Hume R, Willatts P 2013 Maternal and umbilical cord levels of T4, FT4, TSH, TPOAb, and TgAb in term infants and neurodevelopmental outcome at 5.5 years. *J Clin Endocrinol Metab* **98**:829–838.
201. Wasserman EE, Pillion JP, Duggan A, Nelson K, Rohde C, Seaberg EC, Talor MV, Yolken RH, Rose NR 2012 Childhood IQ, hearing loss, and maternal thyroid autoimmunity in the Baltimore Collaborative Perinatal Project. *Pediatr Res* **72**:525–530.
202. Ghassabian A, Bongers-Schokking JJ, de Rijke YB, van Mil N, Jaddoe VW, de Muinck Keizer-Schrama SM, Hooijkaas H, Hofman A, Visser W, Roman GC, Visser TJ, Verhulst FC, Tiemeier H 2012 Maternal thyroid autoimmunity during pregnancy and the risk of attention deficit/hyperactivity problems in children: the Generation R Study. *Thyroid* **22**:178–186.
203. Brown AS, Surcel HM, Hinkka-Yli-Salomäki S, Cheslack-Postava K, Bao Y, Sourander A 2015 Maternal thyroid autoantibody and elevated risk of autism in a national birth cohort. *Prog Neuropsychopharmacol Biol Psychiatry* **57**:86–92.
204. Zegers-Hochschild F, G. D. Adamson, J. de Mouzon, O. Ishihara, R. Mansour, K. Nygren, E. Sullivan, and S. Vanderpoel, for ICMART and WHO 2009 International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology, 2009. *Fertil Steril* **92**:1520–1524.
205. Chandra A, Martinez GM, Mosher WD, Abma JC, Jones J 2005 Fertility, family planning, and reproductive health of U.S. women: data from the 2002 National Survey of Family Growth. *Vital Health Stat* **23(25)**:1–160.
206. Krassas GE, Poppe K, Glinioer D 2010 Thyroid function and human reproductive health. *Endocr Rev* **31**:702–755.
207. Krassas GE, Pontikides N, Kaltsas T, Papadopoulou P, Batrinos M 1994 Menstrual disturbances in thyrotoxicosis. *Clin Endocrinol (Oxf)* **40**:641–644.
208. Poppe K, Glinioer D, Van Steirteghem A, Tournaye H, Devroey P, Schiettecatte J, Velkeniers B 2002 Thyroid dysfunction and autoimmunity in infertile women. *Thyroid* **12**:997–1001.
209. Krassas GE, Pontikides N, Kaltsas T, Papadopoulou P, Paunkovic J, Paunkovic N, Duntas LH 1999 Disturbances of menstruation in hypothyroidism. *Clin Endocrinol (Oxf)* **50**:655–659.
210. Grassi G, Balsamo A, Ansaldi C, Balbo A, Massobrio M, Benedetto C 2001 Thyroid autoimmunity and infertility. *Gynecol Endocrinol* **15**:389–396.
211. Quintino-Moro A, Zantut-Wittmann DE, Tambascia M, Machado Hda C, Fernandes A 2014 High prevalence of infertility among women with Graves' disease and Hashimoto's thyroiditis. *Int J Endocrinol* **2014**:982705.
212. Lincoln SR, Ke RW, Kutteh WH 1999 Screening for hypothyroidism in infertile women. *J Reprod Med* **44**:455–457.
213. Abalovich M, Mitelberg L, Allami C, Gutierrez S, Alcaraz G, Otero P, Levalle O 2007 Subclinical hypothyroidism and thyroid autoimmunity in women with infertility. *Gynecol Endocrinol* **23**:279–283.
214. Lee YJ, Kim CH, Kwack JY, Ahn JW, Kim SH, Chae HD, Kang BM 2014 Subclinical hypothyroidism diagnosed by thyrotropin-releasing hormone stimulation test in infertile women with basal thyroid-stimulating hormone levels of 2.5 to 5.0 mIU/L. *Obstet Gynecol Sci* **57**:507–512.
215. Arojoki M, Jokimaa V, Juuti A, Koskinen P, Irjala K, Anttila L 2000 Hypothyroidism among infertile women in Finland. *Gynecol Endocrinol* **14**:127–131.
216. Verma I, Sood R, Juneja S, Kaur S 2012 Prevalence of hypothyroidism in infertile women and evaluation of response

- of treatment for hypothyroidism on infertility. *Int J Appl Basic Med Res* **2**:17–19.
217. Yoshioka W, Amino N, Ide A, Kang S, Kudo T, Nishihara E, Ito M, Nakamura H, Miyauchi A 2015 Thyroxine treatment may be useful for subclinical hypothyroidism in patients with female infertility. *Endocr J* **62**:87–92.
 218. Kachuei M, Jafari F, Kachuei A, Keshteli AH 2012 Prevalence of autoimmune thyroiditis in patients with polycystic ovary syndrome. *Arch Gynecol Obstet* **285**:853–856.
 219. Monteleone P, Parrini D, Faviana P, Carletti E, Casarosa E, Uccelli A, Cela V, Genazzani AR, Artini PG 2011 Female infertility related to thyroid autoimmunity: the ovarian follicle hypothesis. *Am J Reprod Immunol* **66**:108–114.
 220. Ott J, Aust S, Kurz C, Nouri K, Wirth S, Huber JC, Mayerhofer K 2010 Elevated antithyroid peroxidase antibodies indicating Hashimoto's thyroiditis are associated with the treatment response in infertile women with polycystic ovary syndrome. *Fertil Steril* **94**:2895–2897.
 221. Jatzko B, Vytiska-Bistorfer E, Pawlik A, Promberger R, Mayerhofer K, Ott J. 2014 The impact of thyroid function on intrauterine insemination outcome—a retrospective analysis. *Reprod Biol Endocrinol* **12**:28.
 222. Karmon AE, Batsis M, Chavarro JE, Souter I 2015 Preconceptional thyroid-stimulating hormone levels and outcomes of intrauterine insemination among euthyroid infertile women. *Fertil Steril* **103**:258–263.
 223. Baker VL, Rone HM, Pasta DJ, Nelson HP, Gvakharia M, Adamson GD 2006 Correlation of thyroid stimulating hormone (TSH) level with pregnancy outcome in women undergoing in vitro fertilization. *Am J Obstet Gynecol* **194**:1668–1674.
 224. Fumarola A, Grani G, Romanzi D, Del Sordo M, Bianchini M, Aragona A, Tranquilli D, Aragona C 2013 Thyroid function in infertile patients undergoing assisted reproduction. *Am J Reprod Immunol* **70**:336–341.
 225. Reh A, Grifo J, Danoff A 2010 What is a normal thyroid-stimulating hormone (TSH) level? Effects of stricter TSH thresholds on pregnancy outcomes after in vitro fertilization. *Fertil Steril* **94**:2920–2922.
 226. Chai J, Yeung WY, Lee CY, Li HW, Ho PC, Ng HY 2014 Live birth rates following in vitro fertilization in women with thyroid autoimmunity and/or subclinical hypothyroidism. *Clin Endocrinol (Oxf)* **80**:122–127.
 227. Weghofer A, Himaya E, Kushnir VA, Barad DH, Gleicher N 2015 The impact of thyroid function and thyroid autoimmunity on embryo quality in women with low functional ovarian reserve: a case-control study. *Reprod Biol Endocrinol* **13**:43.
 228. Aghahosseini M, Asgharifard H, Aleyasin A, Tehrani Banihashemi A 2014 Effects of thyroid stimulating hormone (TSH) level on clinical pregnancy rate via in vitro fertilization (IVF) procedure. *Med J Islam Repub Iran* **28**:46.
 229. Green KA, Werner MD, Franasiak JM, Juneau CR, Hong KH, Scott RT Jr 2015 Investigating the optimal preconception TSH range for patients undergoing IVF when controlling for embryo quality. *J Assist Reprod Genet* **32**:1469–1476.
 230. Abdel Rahman AH, Aly Abbassy H, Abbassy AA 2010 Improved in vitro fertilization outcomes after treatment of subclinical hypothyroidism in infertile women. *Endocr Pract* **16**:792–797.
 231. Kim CH, Ahn JW, Kang SP, Kim SH, Chae HD, Kang BM 2011 Effect of levothyroxine treatment on in vitro fertilization and pregnancy outcome in infertile women with subclinical hypothyroidism undergoing in vitro fertilization/intracytoplasmic sperm injection. *Fertil Steril* **95**:1650–1654.
 232. Negro R, Mangieri T, Coppola L, Presicce G, Casavola EC, Gismondi R, Locorotondo G, Caroli P, Pezzarossa A, Dazzi D, Hassan H 2005 Levothyroxine treatment in thyroid peroxidase antibody-positive women undergoing assisted reproduction technologies: a prospective study. *Hum Reprod* **20**:1529–1533.
 233. Velkeniers B, Van Meerhaeghe A, Poppe K, Unuane D, Tournaye H, Haentjens P 2013 Levothyroxine treatment and pregnancy outcome in women with subclinical hypothyroidism undergoing assisted reproduction technologies: systematic review and meta-analysis of RCTs. *Hum Reprod Update* **19**:251–258.
 234. Karmisholt J, Andersen S, Laurberg P 2011 Variation in thyroid function in subclinical hypothyroidism: importance of clinical follow-up and therapy. *Eur J Endocrinol* **164**:317–323.
 235. Scoccia B, Demir H, Kang Y, Fierro MA, Winston NJ 2012 In vitro fertilization pregnancy rates in levothyroxine-treated women with hypothyroidism compared to women without thyroid dysfunction disorders. *Thyroid* **22**:631–636.
 236. Busnelli A, Somigliana E, Benaglia L, Leonardi M, Ragni G, Fedele L 2013 In vitro fertilization outcomes in treated hypothyroidism. *Thyroid* **23**:1319–1325.
 237. Toulis KA, Goulis DG, Venetis CA, Kolibianakis EM, Negro R, Tarlatzis BC, Papadimas I 2010 Risk of spontaneous miscarriage in euthyroid women with thyroid autoimmunity undergoing IVF: a meta-analysis. *Eur J Endocrinol* **162**:643–652.
 238. Tan S, Dieterle S, Pechlavanis S, Janssen OE, Fuhrer D 2014 Thyroid autoantibodies per se do not impair intracytoplasmic sperm injection outcome in euthyroid healthy women. *Eur J Endocrinol* **170**:495–500.
 239. Łukaszuk K, Kunicki M, Kulwikowska P, Liss J, Pastuszek E, Jaszczolt M, Męczekalski B, Skowroński K 2015 The impact of the presence of antithyroid antibodies on pregnancy outcome following intracytoplasmic sperm injection-ICSI and embryo transfer in women with normal thyrotropine levels. *J Endocrinol Invest* **38**:1335–1343.
 240. Karacan M, Alwaeely F, Cebi Z, Berberoglugil M, Batukan M, Ulug M, Arvas A, Camlibel T 2013 Effect of antithyroid antibodies on ICSI outcome in antiphospholipid antibody-negative euthyroid women. *Reprod Biomed Online* **27**:376–380.
 241. Zhong YP, Ying Y, Wu HT, Zhou CQ, Xu YW, Wang Q, Li J, Shen XT, Li J 2012 Relationship between antithyroid antibody and pregnancy outcome following in vitro fertilization and embryo transfer. *Int J Med Sci* **9**:121–125.
 242. Revelli A, Casano S, Piane LD, Grassi G, Gennarelli G, Guidetti D, Massobrio M 2009 A retrospective study on IVF outcome in euthyroid patients with anti-thyroid antibodies: effects of levothyroxine, acetyl-salicylic acid and prednisolone adjuvant treatments. *Reprod Biol Endocrinol* **7**:137.
 243. Litwicka K, Arrivi C, Varricchio MT, Mencacci C, Greco E 2015 In women with thyroid autoimmunity, does low-dose prednisolone administration, compared with no adjuvant therapy, improve in vitro fertilization clinical results? *J Obstet Gynaecol Res* **41**:722–728.
 244. Turi A, Giannubilo SR, Zanconi S, Mascetti A, Tranquilli AL 2010 Preconception steroid treatment in infertile women

- with antithyroid autoimmunity undergoing ovarian stimulation and intrauterine insemination: a double-blind, randomized, prospective cohort study. *Clin Ther* **32**:2415–2421.
245. Michael AE, Papageorgiou AT 2008 Potential significance of physiological and pharmacological glucocorticoids in early pregnancy. *Hum Reprod Update* **14**:497–517.
 246. Poppe K, Unuane D, D'Haeseleer M, Tournaye H, Schiettecatte J, Haentjens P, Velkeniers B 2011 Thyroid function after controlled ovarian hyperstimulation in women with and without the hyperstimulation syndrome. *Fertil Steril* **96**:241–245.
 247. Mintziori G, Goulis DG, Toulis KA, Venetis CA, Kolibianakis EM, Tarlatzis BC 2011 Thyroid function during ovarian stimulation: a systematic review. *Fertil Steril* **96**:780–785.
 248. Muller AF, Verhoeff A, Mantel MJ, De Jong FH, Berghout A 2000 Decrease of free thyroxine levels after controlled ovarian hyperstimulation. *J Clin Endocrinol Metab* **85**:545–548.
 249. Poppe K, Glinooer D, Tournaye H, Schiettecatte J, Devroey P, van Steirteghem A, Haentjens P, Velkeniers B 2004 Impact of ovarian hyperstimulation on thyroid function in women with and without thyroid autoimmunity. *J Clin Endocrinol Metab* **89**:3808–3812.
 250. Poppe K, Glinooer D, Tournaye H, Schiettecatte J, Haentjens P, Velkeniers B 2005 Thyroid function after assisted reproductive technology in women free of thyroid disease. *Fertil Steril* **83**:1753–1757.
 251. Gracia CR, Morse CB, Chan G, Schilling S, Prewitt M, Sammel MD, Mandel SJ 2012 Thyroid function during controlled ovarian hyperstimulation as part of in vitro fertilization. *Fertil Steril* **97**:585–591.
 252. Reinblatt S, Herrero B, Correa JA, Shalom-Paz E, Ata B, Wiser A, Morris D, Holzer H 2013 Thyroid stimulating hormone levels rise after assisted reproductive technology. *J Assist Reprod Genet* **30**:1347–1352.
 253. Corbett S, Shmorgun D, Claman P, Reproductive Endocrinology Infertility Committee, Healey S, Gysler M 2014 The prevention of ovarian hyperstimulation syndrome. *J Obstet Gynaecol Can* **36**:1024–1036.
 254. Skweres T, Wójcik D, Ciepluch R, Sliwiński W, Czech R, Gruszczyński W, Rogacki M 2014 Thyroid dysfunction during severe ovarian hyperstimulation syndrome. A case report. *Ginekol Pol* **85**:472–475.
 255. Stuckey BG, Yeap D, Turner SR 2010 Thyroxine replacement during super-ovulation for in vitro fertilization: a potential gap in management? *Fertil Steril* **93**:2414.e1–3.
 256. Busnelli A, Somigliana E, Benaglia L, Sarais V, Ragni G, Fedele L 2014 Thyroid axis dysregulation during in vitro fertilization in hypothyroid-treated patients. *Thyroid* **24**:1650–1655.
 257. Busnelli A, Vannucchi G, Paffoni A, Faulisi S, Fugazzola L, Fedele L, Somigliana E 2015 Levothyroxine dose adjustment in hypothyroid women achieving pregnancy through IVF. *Eur J Endocrinol* **173**:417–424.
 258. Davis LB, Lathi RB, Dahan MH 2007 The effect of infertility medication on thyroid function in hypothyroid women who conceive. *Thyroid* **17**:773–777.
 259. Casey BM, Dashe JS, Wells CE, McIntire DD, Byrd W, Leveno KJ, Cunningham FG 2005 Subclinical hypothyroidism and pregnancy outcomes. *Obstet Gynecol* **105**:239–245.
 260. Allan WC, Haddow JE, Palomaki GE, Williams JR, Mitchell ML, Hermos RJ, Faix JD, Klein RZ 2000 Maternal thyroid deficiency and pregnancy complications: implications for population screening. *J Med Screen* **7**:127–130.
 261. Casey BM, Dashe JS, Spong CY, McIntire DD, Leveno KJ, Cunningham GF 2007 Perinatal significance of isolated maternal hypothyroxinemia identified in the first half of pregnancy. *Obstet Gynecol* **109**:1129–1135.
 262. Stricker R, Echenard M, Eberhart R, Chevaillier MC, Perez V, Quinn FA, Stricker R 2007 Evaluation of maternal thyroid function during pregnancy: the importance of using gestational age-specific reference intervals. *Eur J Endocrinol* **157**:509–514.
 263. Soldin OP, Soldin D, Sastoque M 2007 Gestation-specific thyroxine and thyroid stimulating hormone levels in the United States and worldwide. *Ther Drug Monit* **29**:553–559.
 264. Bocos-Terraz JP, Izquierdo-Alvarez S, Bancalero-Flores JL, Alvarez-Lahuerta R, Aznar-Sauca A, Real-Lopez E, Ibanez-Marco R, Bocanegra-Garcia V, Rivera-Sanchez G 2009 Thyroid hormones according to gestational age in pregnant Spanish women. *BMC Res Notes* **2**:237.
 265. La'ulu SL, Roberts WL 2011 Ethnic differences in first trimester thyroid reference intervals. *Clin Chem* **57**:913–915.
 266. Mannisto T, Surcel HM, Ruokonen A, Vaarasmaki M, Pouta A, Bloigu A, Järvelin MR, Hartikainen AL, Suvanto E 2011 Early pregnancy reference intervals of thyroid hormone concentrations in a thyroid antibody-negative pregnant population. *Thyroid* **21**:291–298.
 267. Medici M, de Rijke YB, Peeters RP, Visser W, de Muinck Keizer-Schrama SM, Jaddoe VV, Hofman A, Hooijkaas H, Steegers EA, Tiemeier H, Bongers-Schokking JJ, Visser TJ 2011 Maternal early pregnancy and newborn thyroid hormone parameters: the Generation R study. *J Clin Endocrinol Metab* **97**:646–652.
 268. Springer D, Zima T, Limanova Z 2009 Reference intervals in evaluation of maternal thyroid function during the first trimester of pregnancy. *Eur J Endocrinol* **160**:791–797.
 269. Medici M, Korevaar TI, Visser WE, Visser TJ, Peeters RP 2015 Thyroid function in pregnancy: what is normal? *Clin Chem* **61**:704–713.
 270. Lambert-Messerlian G, McClain M, Haddow JE, Palomaki GE, Canick JA, Cleary-Goldman J, Malone FD, Porter TF, Nyberg DA, Bernstein P, D'Alton ME; FaSTER Research Consortium 2008 First- and second-trimester thyroid hormone reference data in pregnant women: a FaSTER (First- and Second-Trimester Evaluation of Risk for aneuploidy) Research Consortium study. *Am J Obstet Gynecol* **199**:62.e1–62.e6.
 271. Gilbert RM, Hadlow NC, Walsh JP, Fletcher SJ, Brown SJ, Stuckey BG, Lim EM 2008 Assessment of thyroid function during pregnancy: first-trimester (weeks 9–13) reference intervals derived from Western Australian women. *Med J Aust* **189**:250–253.
 272. Quinn FA, Gridasov GN, Vdovenko SA, Krasnova NA, Vodopianova NV, Epiphanova MA, Schulten M 2005 Prevalence of abnormal thyroid stimulating hormone and thyroid peroxidase antibody-positive results in a population of pregnant women in the Samara region of the Russian Federation. *Clin Chem Lab Med* **43**:1223–1226.
 273. Vaidya B, Hubalewska-Dydejczyk A, Laurberg P, Negro R, Vermiglio F, Poppe K 2012 Treatment and screening of hypothyroidism in pregnancy: results of a European survey. *Eur J Endocrinol* **166**:49–54.
 274. Vaidya B, Anthony S, Bilous M, Shields B, Drury J, Hutchison S, Bilous R 2007 Detection of thyroid dysfunction

- in early pregnancy: universal screening or targeted high-risk case findings? *J Clin Endocrinol Metab* **92**:203–207.
275. van den Boogaard E, Vissenberg R, Land JA, van Wely M, van der Post JA, Goddijn M, Bisschop PH 2011 Significance of (sub)clinical thyroid dysfunction and thyroid autoimmunity before conception and in early pregnancy: a systematic review. *Hum Reprod Update* **17**:605–619.
 276. Haddow JE, Palomaki GE, Allan WC, Williams JR, Knight GJ, Gagnon J, O’Heir CE, Mitchell ML, Hermos RJ, Waisbren SE, Faix JD, Klein RZ 1999 Maternal thyroid deficiency during pregnancy and subsequent neuropsychological development of the child. *N Engl J Med* **341**:549–555.
 277. Abalovich M, Gutierrez S, Alcaraz G, Maccallini G, Garcia A, Levalle O 2002 Overt and subclinical hypothyroidism complicating pregnancy. *Thyroid* **12**:63–68.
 278. Leung AS, Millar LK, Koonings PP, Montoro M, Mestman JH 1993 Perinatal outcome in hypothyroid pregnancies. *Obstet Gynecol* **81**:349–353.
 279. Sahu MT, Das V, Mittal S, Agarwal A, Sahu M 2010 Overt and subclinical thyroid dysfunction among Indian pregnant women and its effect on maternal and fetal outcome. *Arch Gynecol Obstet* **281**:215–220.
 280. Ong GS, Hadlow NC, Brown SJ, Lim EM, Walsh JP 2014 Does the thyroid-stimulating hormone measured concurrently with first trimester biochemical screening tests predict adverse pregnancy outcomes occurring after 20 weeks gestation? *J Clin Endocrinol Metab* **99**:E2668–E2672.
 281. Chen LM, Du WJ, Dai J, Zhang Q, Si GX, Yang H, Ye EL, Chen QS, Yu LC, Zhang C, Lu XM 2014 Effects of subclinical hypothyroidism on maternal and perinatal outcomes during pregnancy: a single-center cohort study of a Chinese population. *PLoS One* **9**(10):e109364.
 282. Godoy GA, Korevaar TI, Peeters RP, Hofman A, de Rijke YB, Bongers-Schokking JJ, Tiemeier H, Jaddoe VW, Gaillard R 2014 Maternal thyroid hormones during pregnancy, childhood adiposity and cardiovascular risk factors: the Generation R Study. *Clin Endocrinol (Oxf)* **81**:117–125.
 283. León G, Murcia M, Rebagliato M, Álvarez-Pedrerol M, Castilla AM, Basterrechea M, Iñiguez C, Fernández-Somoano A, Blarduni E, Foradada CM, Tardón A, Vioque J 2015 Maternal thyroid dysfunction during gestation, preterm delivery, and birthweight. *The Infancia y Medio Ambiente Cohort, Spain. Paediatr Perinat Epidemiol* **29**:113–122.
 284. Pääkkilä F, Männistö T, Pouta A, Hartikainen AL, Ruokonen A, Surcel HM, Bloigu A, Väärasmäki M, Järvelin MR, Moilanen I, Suvanto E 2014 The impact of gestational thyroid hormone concentrations on ADHD symptoms of the child. *J Clin Endocrinol Metab* **99**:E1–E8.
 285. Medici M, Korevaar T, Schalekamp-Timmermans S, Gaillard R, de Rijke YB, Visser WE, Visser W, de Muinck Keizer-Schrama SM, Hofman A, Hooijkaas H, Bongers-Schokking JJ, Tiemeier H, Jaddoe VW, Visser TJ, Peeters RP, Steegers EA 2014 Maternal early-pregnancy thyroid function is associated with subsequent hypertensive disorders of pregnancy: the generation R study. *J Clin Endocrinol Metab* **99**:E2591–E2598.
 286. Negro R, Schwartz A, Gismondi R, Tinelli A, Mangieri T, Stagnaro-Green A 2010 Increased pregnancy loss rate in thyroid antibody negative women with TSH levels between 2.5 and 5.0 in the first trimester of pregnancy. *J Clin Endocrinol Metab* **95**:E44–E48.
 287. Benhadi N, Wiersinga WM, Reitsma JB, Vrijkotte TG, Bonsel GJ 2009 Higher maternal TSH levels in pregnancy are associated with increased risk for miscarriage, fetal or neonatal death. *Eur J Endocrinol* **160**:985–991.
 288. Liu H, Shan Z, Li C, Mao J, Xie X, Wang W, Fan C, Wang H, Zhang H, Han C, Wang X, Liu X, Fan Y, Bao S, Teng W 2014 Maternal subclinical hypothyroidism, thyroid autoimmunity, and the risk of miscarriage: a prospective cohort study. *Thyroid* **24**:1642–1649.
 289. Cleary-Goldman J, Malone FD, Lambert-Messerlian G, Sullivan L, Canick J, Porter TF, Luthy D, Gross S, Bianchi DW, D’Alton ME 2008 Maternal thyroid hypofunction and pregnancy outcome. *Obstet Gynecol* **112**:85–92.
 290. Stagnaro-Green A, Chen X, Bogden JD, Davies TF, Scholl TO 2005 The thyroid and pregnancy: a novel risk factor for very preterm delivery. *Thyroid* **15**:351–357.
 291. Su PY, Huang K, Hao JH, Xu YQ, Yan SQ, Li T, Xu YH, Tao FB 2011 Maternal thyroid function in the first twenty weeks of pregnancy and subsequent fetal and infant development: a prospective population-based cohort study in China. *J Clin Endocrinol Metab* **96**:3234–3241.
 292. Schneuer FJ, Nassar N, Tasevski V, Morris JM, Roberts CL 2012 Association and predictive accuracy of high TSH serum levels in first trimester and adverse pregnancy outcomes. *J Clin Endocrinol Metab* **97**:3115–3122.
 293. Wilson KL, Casey BM, McIntire DD, Halvorson LM, Cunningham FG 2012 Subclinical thyroid disease and the incidence of hypertension in pregnancy. *Obstet Gynecol* **119**:315–320.
 294. Mannisto T, Vaarasmäki M, Pouta A, Hartikainen AL, Ruokonen A, Surcel HM, Bloigu A, Järvelin MR, Suvanto E 2010 Thyroid dysfunction and autoantibodies during pregnancy as predictive factors of pregnancy complications and maternal morbidity in later life. *J Clin Endocrinol Metab* **95**:1084–1094.
 295. Mannisto T, Vaarasmäki M, Pouta A, Hartikainen AL, Ruokonen A, Surcel HM, Bloigu A, Järvelin MR, Suvanto-Luukkonen E 2009 Perinatal outcome of children born to mothers with thyroid dysfunction or antibodies: a prospective population-based cohort study. *J Clin Endocrinol Metab* **94**:772–779.
 296. Chan S, Boelaert K 2015 Optimal management of hypothyroidism, hypothyroxinaemia and euthyroid TPO antibody positivity preconception and in pregnancy. *Clin Endocrinol (Oxf)* **82**:313–326.
 297. Henrichs J, Bongers-Schokking JJ, Schenk JJ, Ghassabian A, Schmidt HG, Visser TJ, Hooijkaas H, de Muinck Keizer-Schrama SM, Hofman A, Jaddoe VW, Visser W, Steegers EA, Verhulst FC, de Rijke YB, Tiemeier H 2010 Maternal thyroid function during early pregnancy and cognitive functioning in early childhood: the Generation R study. *J Clin Endocrinol Metab* **95**:4227–4234.
 298. Craig WY, Allan WC, Kloza EM, Pulkkinen AJ, Waisbren S, Spratt DI, Palomaki GE, Neveux LM, Haddow JE 2012 Mid-gestational maternal free thyroxine concentration and offspring neurocognitive development at age two years. *J Clin Endocrinol Metab* **97**:E22–E28.
 299. Román GC, Ghassabian A, Bongers-Schokking JJ, Jaddoe VW, Hofman A, de Rijke YB, Verhulst FC, Tiemeier H 2013 Association of gestational maternal hypothyroxinemia and increased autism risk. *Ann Neurol* **74**:733–742.
 300. Finken MJ, van Eijsden M, Loomans EM, Vrijkotte TG, Rotteveel J 2013 Maternal hypothyroxinemia in early pregnancy predicts reduced performance in reaction time

- tests in 5- to 6-year-old offspring. *J Clin Endocrinol Metab* **98**:1417–1426.
301. Julvez J, Alvarez-Pedrerol M, Rebagliato M, Murcia M, Fornis J, Garcia-Esteban R, Lertxundi N, Espada M, Tardón A, Rialño Galán I, Sunyer J 2013 Thyroxine levels during pregnancy in healthy women and early child neurodevelopment. *Epidemiology* **24**:150–157.
 302. Ghassabian AE, Marroun H, Peeters RP, Jaddoe VW, Hofman A, Verhulst FC, Tiemeier H, White T 2014 Downstream effects of maternal hypothyroxinemia in early pregnancy: nonverbal IQ and brain morphology in school-age children. *J Clin Endocrinol Metab* **99**:2383–2390.
 303. Gyllenberg D, Sourander A, Surcel HM, Hinkka-Yli-Salomäki S, McKeague IW, Brown AS 2016 Hypothyroxinemia during gestation and offspring schizophrenia in a national birth cohort. *Biol Psychiatry* **79**:962–970.
 304. Modesto T, Tiemeier H, Peeters RP, Jaddoe VW, Hofman A, Verhulst FC, Ghassabian A 2015 Maternal mild thyroid hormone insufficiency in early pregnancy and attention-deficit/hyperactivity disorder symptoms in children. *JAMA Pediatr* **169**:838–845.
 305. Pop VJ, de Vries E, van Baar AL, Waelkens JJ, de Rooy HA, Horsten M, Donkers MM, Komprou IH, van Son MM, Vader HL 1995 Maternal thyroid peroxidase antibodies during pregnancy: a marker of impaired child development? *J Clin Endocrinol Metab* **80**:3561–3566.
 306. Lazarus JH, Bestwick JP, Channon S, Paradise R, Maina A, Rees R, Chiusano E, John R, Guaraldo V, George LM, Perona M, Dall'Amico D, Parkes AB, Joomun M, Wald NJ 2012 Antenatal thyroid screening and childhood cognitive function. *N Engl J Med* **366**:493–501.
 307. Liu H, Momotani N, Noh JY, Ishikawa N, Takebe K, Ito K 1994 Maternal hypothyroidism during early pregnancy and intellectual development of the progeny. *Arch Intern Med* **154**:785–787.
 308. Momotani N, Iwama S, Momotani K 2012 Neurodevelopment in children born to hypothyroid mothers restored to normal thyroxine (T4) concentration by late pregnancy in Japan: no apparent influence of maternal T4 deficiency. *J Clin Endocrinol Metab* **97**:1104–1108.
 309. Downing S, Halpern L, Carswell J, Brown RS 2012 Severe maternal hypothyroidism corrected prior to the third trimester is associated with normal cognitive outcome in the offspring. *Thyroid* **22**:625–630.
 310. Ausó E, Lavado-Autric R, Cuevas E, Del Rey FE, Morreale De Escobar G, Berbel P 2004 A moderate and transient deficiency of maternal thyroid function at the beginning of fetal neocortico-genesis alters neuronal migration. *Endocrinol* **145**:4037–4047.
 311. Wang S, Teng W, Gao Y, Fan C, Zhang H, Shan Z 2012 Early levothyroxine treatment on maternal subclinical hypothyroidism improves spatial learning of offspring in rats. *J Neuroendocrinol* **24**:841–848.
 312. Pop VJ, Brouwers EP, Vader HL, Vulsma T, van Baar AL, de Vijlder JJ 2003 Maternal hypothyroxinaemia during early pregnancy and subsequent child development: a 3-year follow-up study. *Clin Endocrinol (Oxf)* **59**:282–288.
 313. Henrichs J, Ghassabian A, Peeters RP, Tiemeier H 2013 Maternal hypothyroxinemia and effects on cognitive functioning in childhood: how and why? *Clin Endocrinol (Oxf)* **79**:152–162.
 314. Medici M, Timmermans S, Visser W, de Muinck Keizer-Schrama SM, Jaddoe VW, Hofman A, Hooijkaas H, de Rijke YB, Tiemeier H, Bongers-Schokking JJ, Visser TJ, Peeters RP, Steegers EA 2013 Maternal thyroid hormone parameters during early pregnancy and birth weight: the Generation R Study. *J Clin Endocrinol Metab* **98**:59–66.
 315. León G, Murcia M, Rebagliato M, Álvarez-Pedrerol M, Castilla AM, Basterrechea M, Iñiguez C, Fernández-Somoano A, Blarduni E, Foradada CM, Tardón A, Vioque J 2015 Maternal thyroid dysfunction during gestation, preterm delivery, and birthweight. The Infancia y Medio Ambiente Cohort, Spain. *Paediatr Perinat Epidemiol* **29**:113–122.
 316. Haddow JE, Craig WY, Neveux LM, Haddow HR, Palomaki GE, Lambert-Messerlian G, Malone FD, D'Alton ME, First and Second Trimester Risk of Aneuploidy (FaSTER) Research Consortium 2014 Implications of high free thyroxine (FT4) concentrations in euthyroid pregnancies: the FaSTER trial. *J Clin Endocrinol Metab* **99**:2038–2044.
 317. Korevaar TI, Muetzel R, Medici M, Chaker L, Jaddoe VW, de Rijke YB, Steegers EA, Visser TJ, White T, Tiemeier H, Peeters RP 2016 Association of maternal thyroid function during early pregnancy with offspring IQ and brain morphology in childhood: a population-based prospective cohort study. *Lancet Diabetes Endocrinol* **4**:35–43.
 318. Taylor PN, Minassian C, Rehman A, Iqbal A, Draman MS, Hamilton W, Dunlop D, Robinson A, Vaidya B, Lazarus JH, Thomas S, Dayan CM, Okosieme OE 2014 TSH levels and risk of miscarriage in women on long-term levothyroxine: a community-based study. *J Clin Endocrinol Metab* **99**:3895–3902.
 319. Lev-Ran A 1983 Part-of-the-day hypertriiodothyroninemia caused by desiccated thyroid. *JAMA* **250**:2790–2791.
 320. Wiersinga W, Duntas L, Fadéyev V, Nygaard B, Vanderpump M 2012 2012 ETA guidelines: the use of L-T4 + L-T3 in the treatment of hypothyroidism. *Eur Thyroid J* **1**:55–71.
 321. Alexander EK, Marqusee E, Lawrence J, Jarolim P, Fischer GA, Larsen PR 2004 Timing and magnitude of increases in levothyroxine requirements during pregnancy in women with hypothyroidism. *N Engl J Med* **351**:241–249.
 322. Yassa L, Marqusee E, Fawcett R, Alexander EK 2010 Thyroid hormone early adjustment in pregnancy (the THERAPY) trial. *J Clin Endocrinol Metab* **95**:3234–3241.
 323. Mandel SJ, Larsen PR, Seely EW, Brent GA 1990 Increased need for thyroxine during pregnancy in women with primary hypothyroidism. *N Engl J Med* **323**:91–96.
 324. Kaplan MM 1992 Monitoring thyroxine treatment during pregnancy. *Thyroid* **2**:147–152.
 325. Loh JA, Wartofsky L, Jonklaas J, Burman KD 2009 The magnitude of increased levothyroxine requirements in hypothyroid pregnant women depends upon the etiology of the hypothyroidism. *Thyroid* **19**:269–275.
 326. Abalovich M, Alcaraz G, Kleiman-Rubinshtein J, Pavlove MM, Cornelio C, Levalle O, Gutierrez S 2010 The relationship of preconception thyrotropin levels to requirements for increasing the levothyroxine dose during pregnancy in women with primary hypothyroidism. *Thyroid* **20**:1175–1178.
 327. Galofre JC, Haber RS, Mitchell AA, Pessah R, Davies TF. 2010 Increased postpartum thyroxine replacement in Hashimoto's thyroiditis. *Thyroid* **20**:901–908.
 328. Cooper DS, Laurberg P 2013 Hyperthyroidism in pregnancy. *Lancet Diabetes Endocrinol* **1**:238–249.

329. Bolz M, Korber S, Schober HC 2013 TSH secreting adenoma of pituitary gland (TSHom)—rare cause of hyperthyroidism in pregnancy. *Dtsch Med Wochenschr* **138**:362–366.
330. Merza Z, White D, Khanem N 2015 Struma ovarii in pregnancy: an uncommon cause of hyperthyroidism. *Clin Nucl Med* **40**:687–688.
331. Paschke R, Niedziela M, Vaidya B, Persani L, Rapoport B, Leclere J 2012 2012 European Thyroid Association guidelines for the management of familial and persistent sporadic non-autoimmune hyperthyroidism caused by thyroid-stimulating hormone receptor germline mutations. *Eur Thyroid J* **1**:142–147.
332. Goodwin TM, Montoro M, Mestman JH 1992 Transient hyperthyroidism and hyperemesis gravidarum: clinical aspects. *Am J Obstet Gynecol* **167**:648–652.
333. Tan JY, Loh KC, Yeo GS, Chee YC 2002 Transient hyperthyroidism of hyperemesis gravidarum. *BJOG* **109**:683–688.
334. Niebyl JR 2010 Clinical practice. Nausea and vomiting in pregnancy. *N Engl J Med* **363**:1544–1550.
335. Verberg MF, Gillott DJ, Al-Fardan N, Grudzinskas JG 2005 Hyperemesis gravidarum, a literature review. *Hum Reprod Update* **11**:527–539.
336. Hershman JM 1999 Human chorionic gonadotropin and the thyroid: hyperemesis gravidarum and trophoblastic tumors. *Thyroid* **9**:653–657.
337. Grun JP, Meuris S, De Nayer P, Glinoe D 1997 The thyrotrophic role of human chorionic gonadotrophin (hCG) in the early stages of twin (versus single) pregnancies. *Clin Endocrinol (Oxf)* **46**:719–725.
338. Rodien P, Bremont C, Sanson ML, Parma J, Van Sande J, Costagliola S, Luton JP, Vassart G, Duprez L 1998 Familial gestational hyperthyroidism caused by a mutant thyrotropin receptor hypersensitive to human chorionic gonadotropin. *N Engl J Med* **339**:1823–1826.
339. Glinoe D, Spencer CA 2010 Serum TSH determinations in pregnancy: how, when and why? *Nat Rev Endocrinol* **6**:526–529.
340. Yoshihara A, Noh JY, Mukasa K, Suzuki M, Ohye H, Matsumoto M, Kunii Y, Watanabe N, Suzuki N, Kameda T, Sugino K, Ito K 2015 Serum human chorionic gonadotropin levels and thyroid hormone levels in gestational transient thyrotoxicosis: is the serum hCG level useful for differentiating between active Graves' disease and GTT? *Endocr J* **62**:557–560.
341. Carle A, Knudsen N, Pedersen IB, Perrild H, Ovesen L, Rasmussen LB, Laurberg P 2013 Determinants of serum T4 and T3 at the time of diagnosis in nosological types of thyrotoxicosis: a population-based study. *Eur J Endocrinol* **169**:537–545.
342. Bouillon R, Naesens M, Van Assche FA, De Keyser L, De Moor P, Renaer M, De Vos P, De Roo M 1982 Thyroid function in patients with hyperemesis gravidarum. *Am J Obstet Gynecol* **143**:922–926.
343. Ross DS, Burch HB, Cooper DS, Greenlee MC, Laurberg P, Maia AL, Rivkees SA, Samuels M, Sosa JA, Stan MN, Walter MA 2016 2016 American Thyroid Association guidelines for diagnosis and management of hyperthyroidism and other causes of thyrotoxicosis. *Thyroid* **26**:1343–1421.
344. Laurberg P, Bornaud C, Karmisholt J, Orgiazzi J 2009 Management of Graves' hyperthyroidism in pregnancy: focus on both maternal and foetal thyroid function, and caution against surgical thyroidectomy in pregnancy. *Eur J Endocrinol* **160**:1–8.
345. Alexander EK, Larsen PR 2002 High dose of (131)I therapy for the treatment of hyperthyroidism caused by Graves' disease. *J Clin Endocrinol Metab* **87**:1073–1077.
346. Moura-Neto A, Mosci C, Santos AO, Amorim BJ, de Lima MC, Etchebehere EC, Tambascia MA, Ramos CD, Zantut-Wittmann DE 2012 Predictive factors of failure in a fixed 15 mCi ¹³¹I-iodide therapy for Graves' disease. *Clin Nucl Med* **37**:550–554.
347. de Jong JA, Verkooijen HM, Valk GD, Zelissen PM, de Keizer B 2013 High failure rates after (131)I therapy in Graves hyperthyroidism patients with large thyroid volumes, high iodine uptake, and high iodine turnover. *Clin Nucl Med* **38**:401–406.
348. Schneider DF, Sonderman PE, Jones MF, Ojomo KA, Chen H, Jaume JC, Elson DF, Perlman SB, Sippel RS 2014 Failure of radioactive iodine in the treatment of hyperthyroidism. *Ann Surg Oncol* **21**:4174–4180.
349. Davis LE, Lucas MJ, Hankins GD, Roark ML, Cunningham FG 1989 Thyrotoxicosis complicating pregnancy. *Am J Obstet Gynecol* **160**:63–70.
350. Millar LK, Wing DA, Leung AS, Koonings PP, Montoro MN, Mestman JH 1994 Low birth weight and pre-eclampsia in pregnancies complicated by hyperthyroidism. *Obstet Gynecol* **84**:946–949.
351. Andersen SL, Olsen J, Laurberg P 2015 Foetal programming by maternal thyroid disease. *Clin Endocrinol (Oxf)* **83**:751–758.
352. Mandel SJ, Cooper DS 2001 The use of antithyroid drugs in pregnancy and lactation. *J Clin Endocrinol Metab* **86**:2354–2359.
353. Azizi F 2006 The safety and efficacy of antithyroid drugs. *Expert Opin Drug Saf* **5**:107–116.
354. Nicholas WC, Fischer RG, Stevenson RA, Bass JD 1995 Single daily dose of methimazole compared to every 8 hours propylthiouracil in the treatment of hyperthyroidism. *South Med J* **88**:973–976.
355. Kallner G, Vitols S, Ljunggren JG 1996 Comparison of standardized initial doses of two antithyroid drugs in the treatment of Graves' disease. *J Intern Med* **239**:525–529.
356. Nakamura H, Noh JY, Itoh K, Fukata S, Miyauchi A, Hamada N 2007 Comparison of methimazole and propylthiouracil in patients with hyperthyroidism caused by Graves' disease. *J Clin Endocrinol Metab* **92**:2157–2162.
357. Homsanit M, Sriussadaporn S, Vannasaeng S, Peerapatdit T, Nitiyanant W, Vichayanrat A 2001 Efficacy of single daily dosage of methimazole vs. propylthiouracil in the induction of euthyroidism. *Clin Endocrinol (Oxf)* **54**:385–390.
358. He CT, Hsieh AT, Pei D, Hung YJ, Wu LY, Yang TC, Lian WC, Huang WS, Kuo SW 2004 Comparison of single daily dose of methimazole and propylthiouracil in the treatment of Graves' hyperthyroidism. *Clin Endocrinol (Oxf)* **60**:676–681.
359. Nakamura H, Miyauchi A, Miyawaki N, Imagawa J 2013 Analysis of 754 cases of antithyroid drug-induced agranulocytosis over 30 years in Japan. *J Clin Endocrinol Metab* **98**:4776–4783.
360. Watanabe N, Narimatsu H, Noh JY, Yamaguchi T, Kobayashi K, Kami M, Kunii Y, Mukasa K, Ito K, Ito K 2012 Antithyroid drug-induced hematopoietic damage: a retrospective cohort study of agranulocytosis and pancytopenia involving 50,385 patients with Graves' disease. *J Clin Endocrinol Metab* **97**:E49–E53.

361. Kobayashi S, Noh JY, Mukasa K, Kunii Y, Watanabe N, Matsumoto M, Ohye H, Suzuki M, Yoshihara A, Iwaku K, Sugino K, Ito K 2014 Characteristics of agranulocytosis as an adverse effect of antithyroid drugs in the second or later course of treatment. *Thyroid* **24**:796–801.
362. Rivkees SA, Mattison DR 2009 Propylthiouracil (PTU) hepatotoxicity in children and recommendations for discontinuation of use. *Int J Pediatr Endocrinol* **2009**:132041.
363. Russo MW, Galanko JA, Shrestha R, Fried MW, Watkins P 2004 Liver transplantation for acute liver failure from drug induced liver injury in the United States. *Liver Transpl* **10**:1018–1023.
364. Bahn RS, Burch HS, Cooper DS, Garber JR, Greenlee CM, Klein IL, Laurberg P, McDougall IR, Rivkees SA, Ross D, Sosa JA, Stan MN 2009 The role of propylthiouracil in the management of Graves' disease in adults: report of a meeting jointly sponsored by the American Thyroid Association and the Food and Drug Administration. *Thyroid* **19**:673–674.
365. Anderson SL, Olsen J, Laurberg P 2016 Antithyroid drug side effects in the population and in pregnancy. *J Clin Endocrinol Metab* **101**:1606–1614.
366. Milham SJ, Elledge W 1972 Maternal methimazole and congenital defects in children. *Teratology* **5**:125–126.
367. Clementi M, Di Gianantonio E, Cassina M, Leoncini E, Botto LD, Mastroiaco P, SAFE-Med Study Group 2010 Treatment of hyperthyroidism in pregnancy and birth defects. *J Clin Endocrinol Metab* **95**:E337–E341.
368. Foulds N, Walpole I, Elmslie F, Mansour S 2005 Carbimazole embryopathy: an emerging phenotype. *Am J Med Genet A* **132A**(2):130–135.
369. Yoshihara A, Noh J, Yamaguchi T, Ohye H, Sato S, Sekiya K, Kosuga Y, Suzuki M, Matsumoto M, Kunii Y, Watanabe N, Mukasa K, Ito K, Ito K 2012. Treatment of Graves' disease with antithyroid drugs in the first trimester of pregnancy and the prevalence of congenital malformation. *J Clin Endocrinol Metab* **97**:2396–2403.
370. Andersen SL, Olsen J, Wu CS, Laurberg P 2013 Birth defects after early pregnancy use of antithyroid drugs: a Danish nationwide study. *J Clin Endocrinol Metab* **98**:4373–4381.
371. Andersen SL, Laurberg P 2014 Antithyroid drugs and congenital heart defects: ventricular septal defect is part of the methimazole/carbimazole embryopathy. *Eur J Endocrinol* **171**(5):C1–C3.
372. Laurberg P, Andersen SL 2014 Therapy of endocrine disease: antithyroid drug use in early pregnancy and birth defects: time windows of relative safety and high risk? *Eur J Endocrinol* **171**:R13–R20.
373. Andersen SL, Olsen J, Wu CS, Laurberg P 2014 Severity of birth defects after propylthiouracil exposure in early pregnancy. *Thyroid* **24**:1533–1540.
374. Rubin PC 1981 Current concepts: beta-blockers in pregnancy. *N Engl J Med* **305**:1323–1326.
375. Sherif IH, Oyan WT, Bosairi S, Carrascal SM 1991 Treatment of hyperthyroidism in pregnancy. *Acta Obstet Gynecol Scand* **70**:461–463.
376. Nedrebo BG, Holm PI, Uhlving S, Sorheim JI, Skeie S, Eide GE, Husebye ES, Lien EA, Aanderud S 2002 Predictors of outcome and comparison of different drug regimens for the prevention of relapse in patients with Graves' disease. *Eur J Endocrinol* **147**:583–589.
377. Moore KL, Persaud TVN, Torchia MG 2013 Human birth defects. In: *The Developing Human: Clinically Oriented Embryology*. 9th edition. Saunders/Elsevier, Philadelphia, PA, pp 471–501.
378. Laurberg P, Krejbjerg A, Andersen SL 2014 Relapse following antithyroid drug therapy for Graves' hyperthyroidism. *Curr Opin Endocrinol Diabetes Obes* **21**:415–421.
379. Plummer HS 1923 Results of administering iodine to patients having exophthalmic goiter. *JAMA* **1923**;80:155–156.
380. Means JH 1948 The response to iodine and to antithyroid drugs in Graves' disease. In: *The Thyroid and its Diseases*. 2nd edition. Lippincott Company, Philadelphia, PA, pp 341–374.
381. Momotani N, Hisaoka T, Noh J, Ishikawa N, Ito K 1992 Effects of iodine on thyroid status of fetus versus mother in treatment of Graves' disease complicated by pregnancy. *J Clin Endocrinol Metab* **75**:738–744.
382. Yoshihara A, Noh JY, Watanabe N, Mukasa K, Ohye H, Suzuki M, Matsumoto M, Kunii Y, Suzuki N, Kameda T, Iwaku K, Kobayashi S, Sugino K, Ito K 2015 Substituting potassium iodide for methimazole as the treatment for Graves' disease during the first trimester may reduce the incidence of congenital anomalies: a retrospective study at a single medical institution in Japan. *Thyroid* **25**:1155–1161.
383. Connelly KJ, Boston BA, Pearce EN, Sesser D, Snyder D, Braverman LE, Pino S, LaFranchi SH 2012 Congenital hypothyroidism caused by excess prenatal maternal iodine ingestion. *J Pediatr* **161**:760–762.
384. Chanoine JP, Pardou A, Bourdoux P, Delange F 1988 Withdrawal of iodinated disinfectants at delivery decreases the recall rate at neonatal screening for congenital hypothyroidism. *Arch Dis Child* **63**:1297–1298.
385. Solomon BL, Wartofsky L, Burman KD 1993 Adjunctive cholestyramine therapy for thyrotoxicosis. *Clin Endocrinol (Oxf)* **38**:39–43.
386. Mercado M, Mendoza-Zubieta V, Bautista-Osorio R, Espinoza-de los Monteros AL 1996 Treatment of hyperthyroidism with a combination of methimazole and cholestyramine. *J Clin Endocrinol Metab* **81**:3191–3193.
387. Tsai WC, Pei D, Wang TF, Wu DA, Li JC, Wei CL, Lee CH, Chen SP, Kuo SW 2005 The effect of combination therapy with propylthiouracil and cholestyramine in the treatment of Graves' hyperthyroidism. *Clin Endocrinol (Oxf)* **62**:521–524.
388. Gurung V, Middleton P, Milan SJ, Hague W, Thornton JG 2013 Interventions for treating cholestasis in pregnancy. *Cochrane Database Syst Rev* **6**:CD000493.
389. Leung AM, Pearce EN, Braverman LE 2010 Perchlorate, iodine and the thyroid. *Best Pract Res Clin Endocrinol Metab* **24**:133–141.
390. Wenzel KW, Lente JR 1984 Similar effects of thionamide drugs and perchlorate on thyroid-stimulating immunoglobulins in Graves' disease: evidence against an immunosuppressive action of thionamide drugs. *J Clin Endocrinol Metab* **58**:62–69.
391. EFSA CONTAM Panel (EFSA Panel on Contaminants in the Food Chain) 2014 Scientific Opinion on the risks to public health related to the presence of perchlorate in food, in particular fruits and vegetables. *EFSA Journal* **12**(10):3869. Available at <https://www.efsa.europa.eu/en/efsajournal/pub/3869>
392. Lazarus JH 2009 Lithium and thyroid. *Best Pract Res Clin Endocrinol Metab* **23**:723–733.

393. Diav-Citrin O, Shechtman S, Tahover E, Finkel-Pekarsky V, Arnon J, Kennedy D, Erebara A, Einarson A, Ornoy A 2014 Pregnancy outcome following in utero exposure to lithium: a prospective, comparative, observational study. *Am J Psychiatry* **171**:785–794.
394. Donnelly MA, Wood C, Casey B, Hobbins J, Barbour LA 2015 Early severe fetal Graves disease in a mother after thyroid ablation and thyroidectomy. *Obstet Gynecol* **125**:1059–1062.
395. Momotani N, Noh JY, Ishikawa N, Ito K 1997 Effects of propylthiouracil and methimazole on fetal thyroid status in mothers with Graves' hyperthyroidism. *J Clin Endocrinol Metab* **82**:3633–3636.
396. Patil-Sisodia K, Mestman JH 2010 Graves hyperthyroidism and pregnancy: a clinical update. *Endocr Pract* **16**:118–129.
397. Glinoe D 1998 Thyroid hyperfunction during pregnancy. *Thyroid* **8**:859–864.
398. Momotani N, Noh J, Oyanagi H, Ishikawa N, Ito K 1986 Antithyroid drug therapy for Graves' disease during pregnancy. Optimal regimen for fetal thyroid status. *N Engl J Med* **315**:24–28.
399. Bliddal S, Rasmussen AK, Sundberg K, Brocks V, Feldt-Rasmussen U 2011 Antithyroid drug-induced fetal goitrous hypothyroidism. *Nat Rev Endocrinol* **7**:396–406.
400. Amino N, Tanizawa O, Mori H, Iwatani Y, Yamada T, Kurachi K, Kumahara Y, Miyai K 1982 Aggravation of thyrotoxicosis in early pregnancy and after delivery in Graves' disease. *J Clin Endocrinol Metab* **55**:108–112.
401. Andersen SL, Olsen J, Carle A, Laurberg P 2015 Hyperthyroidism incidence fluctuates widely in and around pregnancy and is at variance with some other autoimmune diseases: a Danish population-based study. *J Clin Endocrinol Metab* **100**:1164–1171.
402. Hamburger JI 1992 Diagnosis and management of Graves' disease in pregnancy. *Thyroid* **2**:219–224.
403. Laurberg P, Bornaud C, Karmisholt J, Orgiazzi J 2009 Management of Graves' hyperthyroidism in pregnancy: focus on both maternal and foetal thyroid function, and caution against surgical thyroidectomy in pregnancy. *Eur J Endocrinol* **160**:1–8.
404. Rotondi M, Cappelli C, Pirali B, Pirola I, Magri F, Fonte R, Castellano M, Rosei EA, Chiovato L 2008 The effect of pregnancy on subsequent relapse from Graves' disease after a successful course of antithyroid drug therapy. *J Clin Endocrinol Metab* **93**:3985–3988.
405. Nakagawa Y, Mori K, Hoshikawa S, Yamamoto M, Ito S, Yoshida K 2002 Postpartum recurrence of Graves' hyperthyroidism can be prevented by the continuation of antithyroid drugs during pregnancy. *Clin Endocrinol (Oxf)* **57**:467–471.
406. Abraham P, Avenell A, McGeoch SC, Clark LF, Bevan JS 2010 Antithyroid drug regimen for treating Graves' hyperthyroidism. *Cochrane Database Syst Rev* **20(1)**:CD003420.
407. McNab T, Ginsberg J 2005 Use of anti-thyroid drugs in euthyroid pregnant women with previous Graves' disease. *Clin Invest Med* **28**:127–131.
408. The American College of Obstetricians and Gynecologists Committee Opinion on Nonobstetric Surgery During Pregnancy. Available at <http://www.acog.org/Resources-And-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Nonobstetric-Surgery-During-Pregnancy> (accessed May 29, 2016).
409. Kempers MJ, van Tijn DA, van Trotsenburg AS, de Vijlder JJ, Wiedijk BM, Vulsma T 2003 Central congenital hypothyroidism due to gestational hyperthyroidism: detection where prevention failed. *J Clin Endocrinol Metab* **88**:5851–5857.
410. Zwaveling-Soonawala N, van Trotsenburg P, Vulsma T 2009 Central hypothyroidism in an infant born to an adequately treated mother with Graves' disease: an effect of maternally derived thyrotrophin receptor antibodies? *Thyroid* **19**:661–662.
411. Luto D, Le Gac I, Vuillard E, Castanet M, Guibourdenche J, Noel M, Toubert ME, Leger J, Boissinot C, Schlageter MH, Garel C, Tebeka B, Oury JF, Czernichow P, Polak M 2005 Management of Graves' disease during pregnancy: the key role of fetal thyroid gland monitoring. *J Clin Endocrinol Metab* **90**:6093–6098.
412. McKenzie JM, Zakarija M 1992 Fetal and neonatal hyperthyroidism and hypothyroidism due to maternal TSH receptor antibodies. *Thyroid* **2**:155–159.
413. Mitsuda N, Tamaki H, Amino N, Hosono T, Miyai K, Tanizawa O 1992 Risk factors for developmental disorders in infants born to women with Graves disease. *Obstet Gynecol* **80**:359–364.
414. Peleg D, Cada S, Peleg A, Ben-Ami M 2002 The relationship between maternal serum thyroid-stimulating immunoglobulin and fetal and neonatal thyrotoxicosis. *Obstet Gynecol* **99**:1040–1043.
415. Laurberg P, Wallin G, Tallstedt L, Abraham-Nordling M, Lundell G, Tørring O 2008 TSH-receptor autoimmunity in Graves' disease after therapy with anti-thyroid drugs, surgery, or radioiodine: a 5-year prospective randomized study. *Eur J Endocrinol* **158**:69–75.
416. Laurberg P, Nygaard B, Glinoe D, Grussendorf M, Orgiazzi J 1998 Guidelines for TSH-receptor antibody measurements in pregnancy: results of an evidence-based symposium organized by the European Thyroid Association. *Eur J Endocrinol* **139**:584–586.
417. Zimmerman D 1999 Fetal and neonatal hyperthyroidism. *Thyroid* **9**:727–733.
418. Abeillon-du Payrat J, Chikh K, Bossard N, Bretones P, Gaucherand P, Claris O, Charrié A, Raverot V, Orgiazzi J, Borson-Chazot F, Bornaud C 2014 Predictive value of maternal second-generation thyroid-binding inhibitory immunoglobulin assay for neonatal autoimmune hyperthyroidism. *Eur J Endocrinol* **171**:451–460.
419. Besancon A, Beltrand J, Le Gac I, Luto D, Polak M 2014 Management of neonates born to women with Graves' disease: a cohort study. *Eur J Endocrinol* **170**:855–862.
420. Phoojaroenchanachai M, Sriussadaporn S, Peerapatdit T, Vannasaeng S, Nitiyanant W, Boonnamsiri V, Vichayanrat A 2001 Effect of maternal hyperthyroidism during late pregnancy on the risk of neonatal low birth weight. *Clin Endocrinol (Oxf)* **54**:365–370.
421. Polak M, Le Gac I, Vuillard E, Guibourdenche J, Leger J, Toubert ME, Madec AM, Oury JF, Czernichow P, Luto D 2004 Fetal and neonatal thyroid function in relation to maternal Graves' disease. *Best Pract Res Clin Endocrinol Metab* **18**:289–302.
422. Cohen O, Pinhas-Hamiel O, Sivan E, Dolitski M, Lipitz S, Achiron R 2003 Serial in utero ultrasonographic measurements of the fetal thyroid: a new complementary tool in the management of maternal hyperthyroidism in pregnancy. *Prenat Diagn* **23**:740–742.

423. Huel C, Guibourdenche J, Vuillard E, Ouahba J, Piketty M, Oury JF, Luton D 2009 Use of ultrasound to distinguish between fetal hyperthyroidism and hypothyroidism on discovery of a goiter. *Ultrasound Obstet Gynecol* **33**:412–420.
424. Porreco RP, Bloch CA 1990 Fetal blood sampling in the management of intrauterine thyrotoxicosis. *Obstet Gynecol* **76**:509–512.
425. Daffos F, Capella-Pavlovsky M, Forestier F 1985 Fetal blood sampling during pregnancy with use of a needle guided by ultrasound: a study of 606 consecutive cases. *Am J Obstet Gynecol* **153**:655–660.
426. Nachum Z, Rakover Y, Weiner E, Shalev E 2003 Graves' disease in pregnancy: prospective evaluation of a selective invasive treatment protocol. *Am J Obstet Gynecol* **189**:159–165.
427. Kilpatrick S 2003 Umbilical blood sampling in women with thyroid disease in pregnancy: is it necessary? *Am J Obstet Gynecol* **189**:1–2.
428. Kahaly GJ 2015 Bioassays for TSH receptor antibodies: quo vadis? *Eur Thyroid J* **4**:3–5.
429. Laurberg P, Pedersen KM, Vestergaard H, Sigurdsson G 1991 High incidence of multinodular toxic goitre in the elderly population in a low iodine intake area vs. high incidence of Graves' disease in the young in a high iodine intake area: comparative surveys of thyrotoxicosis epidemiology in East-Jutland Denmark and Iceland. *J Intern Med* **229**:415–420.
430. Carle A, Pedersen IB, Knudsen N, Perrild H, Ovesen L, Rasmussen LB, Laurberg P 2011 Epidemiology of subtypes of hyperthyroidism in Denmark: a population-based study. *Eur J Endocrinol* **164**:801–809.
431. Stagnaro-Green A 2012 Approach to the patient with postpartum thyroiditis. *J Clin Endocrinol Metab* **97**:334–342.
432. Lazarus JH, Parkes AB, Premawardhana LD 2002 Postpartum thyroiditis. *Autoimmunity* **35**:169–173.
433. Stagnaro-Green A 2004 Postpartum thyroiditis. *Best Pract Res Clin Endocrinol Metab* **18**:303–316.
434. Azizi F 2006 Treatment of post-partum thyrotoxicosis. *J Endocrinol Invest* **29**:244–247.
435. Struve CW, Haupt S, Ohlen S 1993 Influence of frequency of previous pregnancies on the prevalence of thyroid nodules in women without clinical evidence of thyroid disease. *Thyroid* **3**:7–9.
436. Kung AW, Chau MT, Lao TT, Tam SC, Low LC 2002 The effect of pregnancy on thyroid nodule formation. *J Clin Endocrinol Metab* **87**:1010–1014.
437. Tan GH, Gharib H, Goellner JR, van Heerden JA, Bahn RS 1996 Management of thyroid nodules in pregnancy. *Arch Intern Med* **156**:2317–2320.
438. Marley EF, Oertel YC 1997 Fine-needle aspiration of thyroid lesions in 57 pregnant and postpartum women. *Diagn Cytopathol* **16**:122–125.
439. Rosen IB, Walfish PG, Nikore V 1985 Pregnancy and surgical thyroid disease. *Surgery* **98**:1135–1140.
440. Smith LH, Danielsen B, Allen ME, Cress R 2003 Cancer associated with obstetric delivery: results of linkage with the California cancer registry. *Am J Obstet Gynecol* **189**:1128–1135.
441. Loh KC 1997 Familial nonmedullary thyroid carcinoma: a meta-review of case series. *Thyroid* **7**:107–113.
442. Hegedus L 2004 Clinical practice. The thyroid nodule. *N Engl J Med* **351**:1764–1771.
443. Haugen BR, Alexander EK, Bible KC, Doherty GM, Mandel SJ, Nikiforov YE, Pacini F, Randolph GW, Sawka AM, Schlumberger M, Schuff KG, Sherman SI, Sosa JA, Steward DL, Tuttle RM, Wartofsky L 2016 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: the American Thyroid Association Guidelines Task Force on thyroid nodules and differentiated thyroid cancer. *Thyroid* **26**:1–133.
444. Tucker MA, Jones PH, Boice JD Jr, Robison LL, Stone BJ, Stovall M, Jenkin RD, Lubin JH, Baum ES, Siegel SE 1991 Therapeutic radiation at a young age is linked to secondary thyroid cancer. The Late Effects Study Group. *Cancer Res* **51**:2885–2888.
445. Pacini F, Vorontsova T, Demidchik E, Molinaro E, Agate L, Romei C, Shavrova E, Cherstvoy ED, Ivashkevitch Y, Kuchinskaya E, Schlumberger M, Ronga G, Filesi M, Pinchera A 1997 Post-Chernobyl thyroid carcinoma in Belarus children and adolescents: comparison with naturally occurring thyroid carcinoma in Italy and France. *J Clin Endocrinol Metab* **81**:3563–3569.
446. Tan GH, Gharib H, Reading CC 1995 Solitary thyroid nodule. Comparison between palpation and ultrasonography. *Arch Intern Med* **155**:2418–2423.
447. Horvath E, Majlis S, Rossi R, Franco C, Niedmann JP, Castro A, Dominguez M 2009 An ultrasonogram reporting system for thyroid nodules stratifying cancer risk for clinical management. *J Clin Endocrinol Metab* **94**:1748–1751.
448. Russ G, Royer B, Bigorgne C, Rouxel A, Bienvenu-Perrard M, Leenhardt L 2013 Prospective evaluation of thyroid imaging reporting and data system on 4550 nodules with and without elastography. *Eur J Endocrinol* **168**:649–655.
449. Bennedbaek FN, Perrild H, Hegedus L 1999 Diagnosis and treatment of the solitary thyroid nodule. Results of a European survey. *Clin Endocrinol (Oxf)* **50**:357–363.
450. Bennedbaek FN, Hegedus L 2000 Management of the solitary thyroid nodule: results of a North American survey. *J Clin Endocrinol Metab* **85**:2493–2498.
451. Hegedus L, Bonnema SJ, Bennedbaek FN 2003 Management of simple nodular goiter: current status and future perspectives. *Endocr Rev* **24**:102–132.
452. Costante G, Meringolo D, Durante C, Bianchi D, Nocera M, Tumino S, Crocetti U, Attard M, Maranghi M, Tortolano M, Filetti S 2007 Predictive value of serum calcitonin levels for preoperative diagnosis of medullary thyroid carcinoma in a cohort of 5817 consecutive patients with thyroid nodules. *J Clin Endocrinol Metab* **92**:450–455.
453. Pacini F, Pinchera A, Giani C, Grasso L, Baschieri L 1980 Serum thyroglobulin concentrations and ¹³¹I whole body scans in the diagnosis of metastases from differentiated thyroid carcinoma (after thyroidectomy). *Clin Endocrinol (Oxf)* **13**:107–110.
454. Papini E, Guglielmi R, Bianchini A, Crescenzi A, Taccogna S, Nardi F, Panunzi C, Rinaldi R, Toscano V, Pacella CM 2002 Risk of malignancy in nonpalpable thyroid nodules: predictive value of ultrasound and color-Doppler features. *J Clin Endocrinol Metab* **87**:1941–1946.
455. Belfiore A, La Rosa GL 2001 Fine-needle aspiration biopsy of the thyroid. *Endocrinol Metab Clin North Am* **30**:361–400.
456. Goellner JR, Gharib H, Grant CS, Johnson DA 1987 Fine needle aspiration cytology of the thyroid, 1980 to 1986. *Acta Cytol* **31**:587–590.

457. Atkinson BF 1993 Fine needle aspiration of the thyroid. *Monogr Pathol* **(35)**:166–199.
458. Solomon D 1993 Fine needle aspiration of the thyroid: an update. *Thyroid Today* **16**:1–9.
459. Oertel YC 1996 Fine-needle aspiration and the diagnosis of thyroid cancer. *Endocrinol Metab Clin North Am* **25**:69–91.
460. Singer PA 1996 Evaluation and management of the solitary thyroid nodule. *Otolaryngol Clin North Am* **29**:577–591.
461. Powers CN, Frable WJ 1996 *Fine Needle Aspiration Biopsy of the Head and Neck*. Butterworth-Heinemann, Boston, MA.
462. Choe W, McDougall IR 1994 Thyroid cancer in pregnant women: diagnostic and therapeutic management. *Thyroid* **4**:433–435.
463. Hamburger JI 1992 Thyroid nodules in pregnancy. *Thyroid* **2**:165–168.
464. Herzon FS, Morris DM, Segal MN, Rauch G, Parnell T 1994 Coexistent thyroid cancer and pregnancy. *Arch Otolaryngol Head Neck Surg* **120**:1191–1193.
465. Moosa M, Mazzaferri EL 1997 Outcome of differentiated thyroid cancer diagnosed in pregnant women. *J Clin Endocrinol Metab* **82**:2862–2866.
466. Yasmeeen S, Cress R, Romano PS, Xing G, Berger-Chen S, Danielsen B, Smith LH 2005 Thyroid cancer in pregnancy. *Int J Gynaecol Obstet* **91**:15–20.
467. Gorman CA 1999 Radioiodine and pregnancy. *Thyroid* **9**:721–726.
468. IAEA Radiation Protection of Patients [International Atomic Energy Agency] 2015 Pregnancy and Radiation Protection in Nuclear Medicine. Available at https://rpop.iaea.org/RPOP/RPoP/Content/SpecialGroups/1_PregnantWomen/PregnancyNuclearMedicine.htm (accessed August 7, 2015).
469. Pauwels EK, Thomson WH, Blokland JA, Schmidt ME, Bourguignon M, El-Maghraby TA, Broerse JJ, Harding LK 1999 Aspects of fetal thyroid dose following iodine-131 administration during early stages of pregnancy in patients suffering from benign thyroid disorders. *Eur J Nucl Med* **26**:1453–1457.
470. Berg GE, Nystrom EH, Jacobsson L, Lindberg S, Lindstedt RG, Mattsson S, Niklasson CA, Noren AH, Westphal OG 1998 Radioiodine treatment of hyperthyroidism in a pregnant women. *J Nucl Med* **39**:357–361.
471. Zanzonico PB 1997 Radiation dose to patients and relatives incident to ¹³¹I therapy. *Thyroid* **7**:199–204.
472. Vannucchi G, Perrino M, Rossi S, Colombo C, Vicentini L, Dazzi D, Beck-Peccoz P, Fugazzola L 2010 Clinical and molecular features of differentiated thyroid cancer diagnosed during pregnancy. *Eur J Endocrinol* **162**:145–151.
473. Lee JC, Zhao JT, Clifton-Bligh RJ, Gill AJ, Gundara JS, Ip J, Sywak MS, Delbridge LW, Robinson BG, Sidhu SB 2012 Papillary thyroid carcinoma in pregnancy: a variant of the disease? *Ann Surg Oncol* **19**:4210–4216.
474. Messuti I, Corvisieri S, Bardesono F, Rapa I, Giorcelli J, Pellerito R, Volante M, Orlandi F 2014 Impact of pregnancy on prognosis of differentiated thyroid cancer: clinical and molecular features. *Eur J Endocrinol* **170**:659–666.
475. Gharib H, Papini E, Valcavi R, Baskin HJ, Crescenzi A, Dottorini ME, Duick DS, Guglielmi R, Hamilton CR Jr, Zeiger MA, Zini M, AACE/AME Task Force on Thyroid Nodules 2006 American Association of Clinical Endocrinologists and Associazione Medici Endocrinologi medical guidelines for clinical practice for the diagnosis and management of thyroid nodules. *Endocr Pract* **12**:63–102.
476. Rosen IB, Walfish PG 1986 Pregnancy as a predisposing factor in thyroid neoplasia. *Arch Surg* **121**:1287–1290.
477. Doherty CM, Shindo ML, Rice DH, Montero M, Mestman JH 1995 Management of thyroid nodules during pregnancy. *Laryngoscope* **105**:251–255.
478. Chong KM, Tsai YL, Chuang J, Hwang JL, Chen KT 2007 Thyroid cancer in pregnancy: a report of 3 cases. *J Reprod Med* **52**:416–418.
479. Kuy S, Roman SA, Desai R, Sosa JA 2009 Outcomes following thyroid and parathyroid surgery in pregnant women. *Arch Surg* **144**:399–406.
480. Uruno T, Shibuya H, Kitagawa W, Nagahama M, Suino K Ito K 2014 Optimal timing of surgery for differentiated thyroid cancer in pregnant women. *World J Surg* **38**:704–708.
481. Mestman JH, Goodwin TM, Montoro MM 1995 Thyroid disorders of pregnancy. *Endocrinol Metab Clin North Am* **24**:41–71.
482. American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer, Cooper DS, Doherty GM, Haugen BR, Kloos RT, Lee SL, Mandel SJ, Mazzaferri EL, McIver B, Pacini F, Schlumberger M, Sherman SI, Steward DL, Tuttle RM 2009 Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* **19**:1167–1214.
483. Wang CC, Friedman L, Kennedy GC, Wang H, Kebebew E, Steward DL, Zeiger MA, Westra WH, Wang Y, Khanafshar E, Fellegara G, Rosai J, Livolsi V, Lanman RB 2011 A large multicenter correlation study of thyroid nodule cytopathology and histopathology. *Thyroid* **21**:243–251.
484. McLeod DS, Watters KF, Carpenter AD, Ladenson PW, Cooper DS, Ding EL 2012 Thyrotropin and thyroid cancer diagnosis: a systematic review and dose-response meta-analysis. *J Clin Endocrinol Metab* **97**:2682–2692.
485. Pacini F, Schlumberger M, Dralle H, Elisei R, Smit JW, Wiersinga W, European Thyroid Cancer Taskforce 2006 European consensus for the management of patients with differentiated thyroid carcinoma of the follicular epithelium. *Eur J Endocrinol* **154**:787–803.
486. Sawka AM, Lakra DC, Lea J, Alshehri B, Tsang RW, Brierley JD, Straus S, Thabane L, Gafni A, Ezzat S, George SR, Goldstein DP 2008 A systematic review examining the effects of therapeutic radioactive iodine on ovarian function and future pregnancy in female thyroid cancer survivors. *Clin Endocrinol (Oxf)* **69**:479–490.
487. Garsi JP, Schlumberger M, Rubino C, Ricard M, Labbe M, Ceccarelli C, Schvartz C, Henri-Amar M, Bardet S, de Vathaire F 2008 Therapeutic administration of ¹³¹I for differentiated thyroid cancer: radiation dose to ovaries and outcome of pregnancies. *J Nucl Med* **49**:845–852.
488. Pacini F, Gasperi M, Fugazzola L, Ceccarelli C, Lippi F, Centoni R, Martino E, Pinchera A 1994 Testicular function in patients with differentiated thyroid carcinoma treated with radioiodine. *J Nucl Med* **35**:1418–1422.
489. FDA. Drug Approvals and Databases. Available at: www.fda.gov/Drugs/InformationonDrugs/default.htm (accessed February 11, 2017).
490. Rosvoll RV, Winship T 1965 Thyroid carcinoma and pregnancy. *Surg Gynecol Obstet* **121**:1039–1042.
491. Hill CS, Jr, Clark RL, Wolf M 1966 The effect of subsequent pregnancy on patients with thyroid carcinoma. *Surg Gynecol Obstet* **122**:1219–1222.
492. Leboeuf R, Emerick LE, Martorella AJ, Tuttle RM 2007 Impact of pregnancy on serum thyroglobulin and detection

- of recurrent disease shortly after delivery in thyroid cancer survivors. *Thyroid* **17**:543–547.
493. Rosario PW, Barroso AL, Purisch S 2007 The effect of subsequent pregnancy on patients with thyroid carcinoma apparently free of the disease. *Thyroid* **17**:1175–1176.
 494. Hirsch D, Levy S, Tsvetov G, Weinstein R, Lifshitz A, Singer J, Shraga-Slutsky I, Grozinski-Glasberg S, Shimon I, Benbassat C 2010 Impact of pregnancy on outcome and prognosis of survivors of papillary thyroid cancer. *Thyroid* **20**:1179–1185.
 495. Ito Y, Miyauchi A, Kihara M, Higashiyama T, Kobayashi K, Miya A 2014 Patient age is significantly related to the progression of papillary microcarcinoma of the thyroid under observation. *Thyroid* **24**:27–34.
 496. Shindo H, Amino N, Ito Y, Kihara M, Kobayashi K, Miya A, Hirokawa M, Miyauchi A 2014 Papillary thyroid microcarcinoma might progress during pregnancy. *Thyroid* **24**:840–844.
 497. Wells SA Jr, Asa SL, Dralle H, Elisei R, Evans DB, Gagel RF, Lee N, Machens A, Moley JF, Pacini F, Raue F, Frank-Raue K, Robinson B, Rosenthal MS, Santoro M, Schlumberger M, Shah M, Waguespack SG, American Thyroid Association Guidelines Task Force on Medullary Thyroid Carcinoma 2015 Revised American Thyroid Association guidelines for the management of medullary thyroid carcinoma. *Thyroid* **25**:567–610.
 498. Bernal J, Pekonen F 1984 Ontogenesis of the nuclear 3,5,3'-triiodothyronine receptor in the human fetal brain. *Endocrinology* **114**:677–679.
 499. Thorpe-Beeston JG, Nicolaides KH, Felton CV, Butler J, McGregor AM 1991 Maturation of the secretion of thyroid hormone and thyroid-stimulating hormone in the fetus. *N Engl J Med* **324**:532–536.
 500. Thorpe-Beeston JG, Nicolaides KH, AM 1992 Fetal thyroid function. *Thyroid* **2**:207–217.
 501. Burrow GN, Fisher DA, Larsen PR 1994 Maternal and fetal thyroid function. *N Engl J Med* **331**:1073–1078.
 502. Brown RS 2004 Minireview: developmental regulation of thyrotropin receptor gene expression in the fetal and newborn thyroid. *Endocrinology* **145**:4058–4061.
 503. Ferreira B, Bernal J, Goodyer CG, Branchard CL 1988 Estimation of nuclear thyroid hormone receptor saturation in human fetal brain and lung during early gestation. *J Clin Endocrinol Metab* **67**:853–856.
 504. Kester MH, Martinez de Mena R, Obregon MJ, Marinkovic D, Howatson A, Visser TJ, Hume R, Morreale de Escobar G 2004 Iodothyronine levels in the human developing brain: major regulatory roles of iodothyronine deiodinases in different areas. *J Clin Endocrinol Metab* **89**:3117–3128.
 505. Wu SY, Green WL, Huang WS, Hays MT, Chopra IJ 2005 Alternate pathways of thyroid hormone metabolism. *Thyroid* **15**:943–958.
 506. Ballabio M, Nicolini U, Jowett T, Ruiz de Elvira MC, Ekins RP, Rodeck CH 1989 Maturation of thyroid function in normal human fetuses. *Clin Endocrinol (Oxf)* **31**:565–571.
 507. Patel J, Landers K, Li H, Mortimer RH, Richard K 2011 Delivery of maternal thyroid hormones to the fetus. *Trends Endocrinol Metab* **22**:164–170.
 508. Vulsma T, Gons MH, de Vijlder JJ 1989 Maternal-fetal transfer of thyroxine in congenital hypothyroidism due to a total organification defect or thyroid agenesis. *N Engl J Med* **321**:13–16.
 509. Li H, Patel J, Mortimer RH, Richard K 2012 Ontogenic changes in human placental sodium iodide symporter expression. *Placenta* **33**:946–948.
 510. Mathieson L, Nielsen LK, Andersen JT, Grevys A, Sandlie I, Michaelsen TE, Hedegaard M, Knudsen LE, Dziegiel MH 2013 Maternofetal transplacental transport of recombinant IgG antibodies lacking effector functions. *Blood* **122**:1174–1181.
 511. Michalek K, Morshed SA, Latif R, Davies TF 2009 TSH receptor autoantibodies. *Autoimmun Rev* **9**:113–116.
 512. Benvenga S, Ordookhani A, Pearce EN, Tonacchera M, Azizi F, Braverman LE 2008 Detection of circulating autoantibodies against thyroid hormones in an infant with permanent congenital hypothyroidism and her twin with transient congenital hypothyroidism: possible contribution of thyroid hormone autoantibodies to neonatal and infant hypothyroidism. *J Pediatr Endocrinol Metab* **21**:1011–1020.
 513. Zakarija M, McKenzie JM 1983 Pregnancy-associated changes in the thyroid-stimulating antibody of Graves' disease and the relationship to neonatal hyperthyroidism. *J Clin Endocrinol Metab* **57**:1036–1040.
 514. Skuza KA, Sills IN, Stene M, Rapaport R 1996 Prediction of neonatal hyperthyroidism in infants born to mothers with Graves disease. *J Pediatr* **128**:264–268.
 515. Brown RS, Bellisario RL, Botero D, Fournier L, Abrams CA, Cowger ML, David R, Fort P, Richman RA 1996 Incidence of transient congenital hypothyroidism due to maternal thyrotropin receptor-blocking antibodies in over one million babies. *J Clin Endocrinol Metab* **81**:1147–1151.
 516. Fort P, Lifshitz F, Pugliese M, Klein I 1988 Neonatal thyroid disease: differential expression in three successive offspring. *J Clin Endocrinol Metab* **66**:645–647.
 517. Zakarija M, McKenzie JM, Munro DS 1983 Immunoglobulin G inhibitor of thyroid-stimulating antibody is a cause of delay in the onset of neonatal Graves' disease. *J Clin Invest* **72**:1352–1356.
 518. Kohn LD, Suzuki K, Hoffman WH, Tombaccini D, Marcocci C, Shimojo N, Watanabe Y, Amino N, Cho BY, Kohno Y, Hirai A, Tahara K 1997 Characterization of monoclonal thyroid-stimulating and thyrotropin binding-inhibiting autoantibodies from a Hashimoto's patient whose children had intrauterine and neonatal thyroid disease. *J Clin Endocrinol Metab* **82**:3998–4009.
 519. Matsuura N, Yamada Y, Nohara Y, Konishi J, Kasagi K, Endo K, Kojima H, Wataya K 1980 Familial neonatal transient hypothyroidism due to maternal TSH-binding inhibitor immunoglobulins. *N Engl J Med* **303**:738–741.
 520. Matsuura N, Konishi J 1990 Transient hypothyroidism in infants born to mothers with chronic thyroiditis—a nationwide study of twenty-three cases. The Transient Hypothyroidism Study Group. *Endocrinol Jpn* **37**:369–379.
 521. Ford G, LaFranchi SH 2014 Screening for congenital hypothyroidism: a worldwide view of strategies. *Best Pract Res Clin Endocrinol Metab* **28**:175–187.
 522. Rose SR, Brown RS, Foley T, Kaplowitz PB, Kaye CI, Sundararajan S, Varma SK 2006 Update of newborn screening and therapy for congenital hypothyroidism. *Pediatrics* **117**:2290–2303.
 523. Leger J, Olivieri A, Donaldson M, Torresani T, Krude H, van Vliet G, Polak M, Butler G, ESPE-PES-SLEP-JSPE-APEG-APPES-ISPAE, Congenital Hypothyroidism Consensus Conference Group 2014 European Society for Paediatric Endocrinology consensus guidelines on

- screening, diagnosis, and management of congenital hypothyroidism. *J Clin Endocrinol Metab* **99**:363–384.
524. Carswell JM, Gordon JH, Popovsky E, Hale A, Brown RS 2013 Generic and brand-name L-thyroxine are not bioequivalent for children with severe congenital hypothyroidism. *J Clin Endocrinol Metab* **98**:610–617.
 525. LaFranchi SH 2011 Approach to the diagnosis and treatment of neonatal hypothyroidism. *J Clin Endocrinol Metab* **96**:2959–2967.
 526. Brown RS. Disorders of the thyroid gland in infants, children, and adolescents. In DeGroot LJ, ed. *Thyroid Disease Manager*. Available at <http://www.thyroidmanager.org>.
 527. Buckshee K, Kriplani A, Kapil A, Bhargava VL, Takkar D 1992 Hypothyroidism complicating pregnancy. *Aust N Z J Obstet Gynaecol* **32**:240–242.
 528. Stein MT, Kessler DB, Hubbard E 2004 Failure to thrive in a four-month-old nursing infant. *J Dev Behav Pediatr* **25**:S69–S73.
 529. Hapon MB, Varas SM, Giménez MS, Jahn GA 2007 Reduction of mammary and liver lipogenesis and alteration of milk composition during lactation in rats by hypothyroidism. *Thyroid* **17**:11–18.
 530. Thrift TA, Bernal A, Lewis AW, Neuendorff DA, Willard CC, Randel RD 1999 Effects of induced hypothyroidism on weight gains, lactation, and reproductive performance of primiparous Brahman cows. *J Anim Sci* **77**:1844–1850.
 531. Rosato RR, Gimenez MS, Jahn GA 1992 Effects of chronic thyroid hormone administration on pregnancy, lactogenesis and lactation in the rat. *Acta Endocrinol (Copenh)* **127**:547–554.
 532. Varas SM, Muñoz EM, Hapon MB, Aguilera Merlo CI, Giménez MS, Jahn GA 2002 Hyperthyroidism and production of precocious involution in the mammary glands of lactating rats. *Reproduction* **124**:691–702.
 533. van Wassenae AG, Stulp MR, Valianpour F, Tamminga P, Ris Stalpers C, de Randamie JS, van Beusekom C, de Vijlder JJ 2002 The quantity of thyroid hormone in human milk is too low to influence plasma thyroid hormone levels in the very preterm infant. *Clin Endocrinol (Oxf)* **56**:621–627.
 534. Akasha MA, Anderson RR, Ellersieck M, Nixon DA 1987 Concentration of thyroid hormones and prolactin in dairy cattle serum and milk at three stages of lactation. *J Dairy Sci* **70**:271–276.
 535. Akasha M, Anderson RR 1984 Thyroxine and triiodothyronine in milk of cows, goats, sheep, and guinea pigs. *Proc Soc Exp Biol Med* **177**:360–371.
 536. Todini L, Salimei E, Malfatti A, Ferraro S, Fantuz F 2012 Thyroid hormones in milk and blood of lactating donkeys as affected by stage of lactation and dietary supplementation with trace elements. *J Dairy Res* **79**:232–237.
 537. Howe DB, Beardsley M, Bakhsh S 2008 Appendix U. Model procedure for release of patients or human research subjects administered radioactive materials. In Consolidated Guidance About Materials Licenses. Program-Specific Guidance About Medical Use Licenses. Final Report. U.S. Nuclear Regulatory Commission Office of Nuclear Material Safety and Safeguards. NUREG-1556, vol. 9, rev. 2. Available at <http://www.nrc.gov/reading-rm/doc-collections/nuregs/staff/sr1556/v9/r2/>.
 538. Kampmann JP, Johansen K, Hansen JM, Helweg J 1980 Propylthiouracil in human milk. Revision of a dogma. *Lancet* **1**:736–737.
 539. Momotani N, Yamashita R, Yoshimoto M, Noh J, Ishikawa N, Ito K 1989 Recovery from foetal hypothyroidism: evidence for the safety of breast-feeding while taking propylthiouracil. *Clin Endocrinol (Oxf)* **31**:591–595.
 540. Momotani N, Yamashita R, Makino F, Noh JY, Ishikawa N, Ito K 2000 Thyroid function in wholly breast-feeding infants whose mothers take high doses of propylthiouracil. *Clin Endocrinol (Oxf)* **53**:177–181.
 541. Johansen K, Andersen AN, Kampmann JP, Mølholm Hansen JM, Mortensen HB 1982 Excretion of methimazole in human milk. *Eur J Clin Pharmacol* **23**:339–341.
 542. Cooper DS, Bode HH, Nath B, Saxe V, Maloof F, Ridgway EC 1984 Methimazole pharmacology in man: studies using a newly developed radioimmunoassay for methimazole. *J Clin Endocrinol Metab* **58**:473–479.
 543. Azizi F, Khoshniat M, Bahrainian M, Hedayati M 2000 Thyroid function and intellectual development of infants nursed by mothers taking methimazole. *J Clin Endocrinol Metab* **85**:3233–3238.
 544. Lambert BA, Ikonen E, Osterlund K, Teramo K, Pekonen F, Peltola J, Välimäki M 1984 Antithyroid treatment of maternal hyperthyroidism during lactation. *Clin Endocrinol (Oxf)* **21**:81–87.
 545. Azizi F 1996 Effect of methimazole treatment of maternal thyrotoxicosis on thyroid function in breast-feeding infants. *J Pediatr* **128**:855–858.
 546. Institute of Medicine, Food and Nutrition Board 2001 Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. National Academy Press, Washington, DC.
 547. World Health Organization, UNICEF, ICCIDD 2007 Assessment of Iodine Deficiency Disorders and Monitoring Their Elimination. A Guide for Programme Managers. 3rd edition. Geneva, Switzerland.
 548. Amino N, Mori H, Iwatani Y, Tanizawa O, Kawashima M, Tsuge I, Ibaragi K, Kumahara Y, Miyai K 1982 High prevalence of transient post-partum thyrotoxicosis and hypothyroidism. *N Engl J Med* **306**:849–852.
 549. Stagnaro-Green A, Schwartz A, Gismondi R, Tinelli A, Mangieri T, Negro R 2011 High rate of persistent hypothyroidism in a large-scale prospective study of postpartum thyroiditis in southern Italy. *J Clin Endocrinol Metab* **96**:652–657.
 550. Muller AF, Drexhage HA, Berghout A 2001 Postpartum thyroiditis and autoimmune thyroiditis in women of childbearing age: recent insights and consequences for antenatal and postnatal care. *Endocr Rev* **22**:605–630.
 551. Kuijpers JL, De Hann-Meulman M, Vader HL, Pop VJ, Wiersinga WM, Drexhage HA 1998 Cell-mediated immunity and postpartum thyroid dysfunction: a possibility for the prediction of disease? *J Clin Endocrinol Metab* **83**:1959–1966.
 552. Stagnaro-Green A, Roman SH, Cobin RH, el-Harazy E, Wallenstein S, Davies TF 1992 A prospective study of lymphocyte-initiated immunosuppression in normal pregnancy: evidence of a T-cell etiology for postpartum thyroid dysfunction. *J Clin Endocrinol Metab* **74**:645–653.
 553. Smallridge RC 2000 Postpartum thyroid disease: a model of immunologic dysfunction. *Clin Appl Immunol Rev* **1**:89–103.
 554. Ide A, Amino N, Kang S, Yoshioka W, Kudo T, Nishihara E, Ito M, Nakamura H, Miyauchi A 2014 Differentiation of postpartum Graves' thyrotoxicosis from postpartum

- destructive thyrotoxicosis using antithyrotropin receptor antibodies and thyroid blood flow. *Thyroid* **24**:1027–1031.
555. Nicholson WK, Robinson KA, Smallridge RC, Ladenson PW, Powe NR 2006 Prevalence of postpartum thyroid dysfunction: a quantitative review. *Thyroid* **16**:573–582.
 556. Gerstein HC 1993 Incidence of postpartum thyroid dysfunction in patients with type I diabetes mellitus. *Ann Intern Med* **118**:419–423.
 557. Alvarez-Marfany M, Roman SH, Drexler AJ, Robertson C, Stagnaro-Green A 1994 Long-term prospective study of postpartum thyroid dysfunction in women with insulin dependent diabetes mellitus. *J Clin Endocrinol Metab* **79**:10–16.
 558. Elefsiniotis IS, Vezali E, Pantazis KD, Saroglou G 2008 Post-partum thyroiditis in women with chronic viral hepatitis. *J Clin Virol* **41**:318–319.
 559. Stagnaro-Green A, Akhter E, Yim C, Davies TF, Magder L, Petri M 2011 Thyroid disease in pregnant women with systemic lupus erythematosus: increased preterm delivery. *Lupus* **20**:690–699.
 560. Tagami T, Hagiwara H, Kimura T, Usui T, Shimatsu A, Naruse M 2007 The incidence of gestational hyperthyroidism and postpartum thyroiditis in treated patients with Graves' disease. *Thyroid* **17**:767–772.
 561. Manetti L, Parkes AB, Lupi I, Di Cianni G, Bogazzi F, Albertini S, Morselli LL, Raffaelli V, Russo D, Rossi G, Gasperi M, Lazarus JH, Martino E 2008 Serum pituitary antibodies in normal pregnancy and in patients with postpartum thyroiditis: a nested case-control study. *Eur J Endocrinol* **159**:805–809.
 562. Lazarus JH, Ammari F, Oretti R, Parkes AB, Richards CJ, Harris B 1997 Clinical aspects of recurrent postpartum thyroiditis. *Br J Gen Pract* **47**:305–308.
 563. Caixas A, Albareda M, Garcia-Patterson A, Rodriguez-Espinosa J, de Leiva A, Corcoy R 1999 Postpartum thyroiditis in women with hypothyroidism antedating pregnancy? *J Clin Endocrinol Metab* **84**:4000–4005.
 564. Marqusee E, Hill JA, Mandel SJ 1997 Thyroiditis after pregnancy loss. *J Clin Endocrinol Metab* **82**:2455–2457.
 565. Walfish PG, Meyerson J, Provias JP, Vargas MT, Papsin FR 1992 Prevalence and characteristics of post-partum thyroid dysfunction: results of a survey from Toronto, Canada. *J Endocrinol Invest* **15**:265–272.
 566. Hayslip CC, Fein HG, O'Donnell VM, Friedman DS, Klein TA, Smallridge RC 1988 The value of serum anti-microsomal antibody testing in screening for symptomatic postpartum thyroid dysfunction. *Am J Obstet Gynecol* **159**:203–209.
 567. Lazarus JH 1999 Clinical manifestations of postpartum thyroid disease. *Thyroid* **9**:685–689.
 568. Kuijpers JL, Vader HL, Drexhage HA, Wiersinga WM, van Son MJ, Pop VJ 2001 Thyroid peroxidase antibodies during gestation are a marker for subsequent depression postpartum. *Eur J Endocrinol* **145**:579–584.
 569. Pop VJ, de Rooy HA, Vader HL, van der Heide D, van Son M, Komproe IH, Essed GG, de Geus CA 1991 Postpartum thyroid dysfunction and depression in an unselected population. *N Engl J Med* **324**:1815–1816.
 570. Hayslip CC, Fein HG, O'Donnell VM, Friedman DS, Klein TA, Smallridge RC 1988 The value of serum anti-microsomal antibody testing in screening for symptomatic postpartum thyroid dysfunction. *Am J Obstet Gynecol* **159**:203–209.
 571. Lucas A, Pizarro E, Granada ML, Salinas I, Sanmarti A 2001 Postpartum thyroid dysfunction and postpartum depression: are they two linked disorders? *Clin Endocrinol (Oxf)* **55**:809–814.
 572. Lazarus JH, Hall R, Othman S, Parkes AB, Richards CJ, McCulloch B, Harris B 1996 The clinical spectrum of postpartum thyroid disease. *QJM* **89**:429–435.
 573. Le Donne M, Settineri S, Benvenga S 2012 Early postpartum alexithymia and risk for depression: relationship with serum thyrotropin, free thyroid hormones and thyroid autoantibodies. *Psychoneuroendocrinology* **37**:519–533.
 574. Harris B, Othman S, Davies JA, Weppner GJ, Richards CJ, Newcombe RG, Lazarus JH, Parkes AB, Hall R, Phillips DI 1992 Association between postpartum thyroid dysfunction and thyroid antibodies and depression. *BMJ* **305**:152–156.
 575. Pop VJ, de Rooy HA, Vader HL, van der Heide D, van Son MM, Komproe IH 1993 Microsomal antibodies during gestation in relation to postpartum thyroid dysfunction and depression. *Acta Endocrinol (Copenh)* **129**:26–30.
 576. Harris B, Oretti R, Lazarus J, Parkes A, John R, Richards C, Newcombe R, Hall R 2002 Randomised trial of thyroxine to prevent postnatal depression in thyroid-antibody-positive women. *Br J Psychiatry* **180**:327–330.
 577. Fung HY, Kologlu M, Collison K, John R, Richards CJ, Hall R, McGregor AM 1988 Postpartum thyroid dysfunction in Mid Glamorgan. *Br Med J (Clin Res Ed)* **296**:241–244.
 578. Vargas MT, Briones-Urbina R, Gladman D, Papsin FR, Walfish PG 1988 Antithyroid microsomal autoantibodies and HLA-DR5 are associated with postpartum thyroid dysfunction: evidence supporting an autoimmune pathogenesis. *J Clin Endocrinol Metab* **67**:327–333.
 579. Rasmussen NG, Hornnes PJ, Hoier-Madsen M, Feldt-Rasmussen U, Hegedus L 1990 Thyroid size and function in healthy pregnant women with thyroid autoantibodies. Relation to development of postpartum thyroiditis. *Acta Endocrinol (Copenh)* **123**:395–401.
 580. Azizi F 2005 The occurrence of permanent thyroid failure in patients with subclinical postpartum thyroiditis. *Eur J Endocrinol* **153**:367–371.
 581. Nohr SB, Jorgensen A, Pedersen KM, Laurberg P 2000 Postpartum thyroid dysfunction in pregnant thyroid peroxidase antibody-positive women living in an area with mild to moderate iodine deficiency: is iodine supplementation safe? *J Clin Endocrinol Metab* **85**:3191–3198.
 582. Kampe O, Jansson R, Karlsson FA 1990 Effects of L-thyroxine and iodide on the development of autoimmune postpartum thyroiditis. *J Clin Endocrinol Metab* **70**:1014–1018.
 583. Klein RZ, Haddow JE, Faix JD, Brown RS, Hermos RJ, Pulkkinen A, Mitchell ML 1991 Prevalence of thyroid deficiency in pregnant women. *Clin Endocrinol (Oxf)* **35**:41–46.
 584. Negro R, Schwartz A, Gismondi R, Tinelli A, Mangieri T, Stagnaro-Green A 2010 Universal screening versus case finding for detection and treatment of thyroid hormonal dysfunction during pregnancy. *J Clin Endocrinol Metab* **95**:1699–1707.
 585. Ma L, Qi H, Chai X, Jiang F, Mao S, Liu J, Zhang S, Lian X, Sun X, Wang D, Ren, Yan Q 2016 The effects of screening and intervention of subclinical hypothyroidism on pregnancy outcomes: a prospective multicenter single-blind, randomized, controlled study of thyroid function

- screening test during pregnancy. *J Matern Fetal Neonatal Med* **29**:1391–1394.
586. TABLET trial: Randomised Controlled Trial of the Efficacy and Mechanism of Levothyroxine Treatment on Pregnancy and Neonatal Outcomes in Women with Thyroid Antibodies (TABLET). Available at <http://www.birmingham.ac.uk/research/activity/mds/trials/bctu/trials/womens/tablet/index.aspx>.
 587. Netherlands TRIAL Register. T4Life trial. <http://www.trialregister.nl/trialreg/admin/rctview.asp?TC=3364>.
 588. Teng W, Shan Z 2011 Pregnancy and thyroid diseases in China. *Thyroid* **21**:1053–1055.
 589. Dosiou C, Sanders GD, Araki SS, Crapo LM 2008 Screening pregnant women for autoimmune thyroid disease: a cost-effectiveness analysis. *Eur J Endocrinol* **158**:841–851.
 590. Dosiou C, Barnes J, Schwartz A, Negro R, Crapo L, Stagnaro-Green A 2012 Cost-effectiveness of universal and risk-based screening for autoimmune thyroid disease in pregnant women. *J Clin Endocrinol Metab* **97**:1536–1546.
 591. Thung SF, Funai EF, Grobman WA 2009 The cost-effectiveness of universal screening in pregnancy for subclinical hypothyroidism. *Am J Obstet Gynecol* **200**:267.e1–267.e7.
 592. Donnay Candil S, Balsa Barro JA, Álvarez Hernández J, Crespo Palomo C, Pérez-Alcántara F, Polanco Sánchez C 2015 Cost-effectiveness analysis of universal screening for thyroid disease in pregnant women in Spain. *Endocrinol Nutrition* **62**:322–330.
 593. Rowley PT, Loader S, Kaplan RM 1998 Prenatal screening for cystic fibrosis carriers: an economic evaluation. *Am J Hum Genet* **63**:1160–1174.
 594. Werner EF, Pettker CM, Zuckerwise L, Reel M, Funai EF, Henderson J, Thung SF 2012 Screening for gestational diabetes mellitus: are the criteria proposed by the International Association of the Diabetes and Pregnancy Study Groups cost-effective? *Diabetes Care* **35**:529–535.
 595. Potlukova E, Potluka O, Jiskra J, Limanova Z, Telicka Z, Bartakova J, Springer D 2012 Is age a risk factor for hypothyroidism in pregnancy? An analysis of 5223 pregnant women. *J Clin Endocrinol Metab* **97**:1945–1952.
 596. Yang H, Shao M, Chen L, Chen Q, Yu L, Cai L, Lin Z, Zhang C, Lu X 2014 Screening strategies for thyroid disorders in the first and second trimester of pregnancy in China. *PLoS ONE* **9**(6):e99611.
 597. Medeiros MF, Cerqueira TL, Silva Junior JC, Amaral MT, Vaidya B, Poppe KG, Carvalho GA, Gutierrez S, Alcaraz G, Abalovich M, Ramos HE, Latin American Thyroid Society 2014 An international survey of screening and management of hypothyroidism during pregnancy in Latin America. *Arq Bras Endocrinol Metab* **58**:906–911.
 598. Vaidya B, Hubalewska-Dydejczyk A, Laurberg P, Negro R, Vermiglio F, Poppe K 2012 Treatment and screening of hypothyroidism in pregnancy: results of a European survey. *Eur J Endocrinol* **166**:49–54.
 599. Azizi F, Amouzegar A, Mehran L, Alamdari S, Subekti I, Vaidya B, Poppe K, San Luis T Jr, Akamizu T 2014 Screening and management of hypothyroidism in pregnancy: results of an Asian Survey. *Endocr J* **61**:697–704.
 600. Srimatkandada P, Stagnaro-Green A, Pearce EN 2015 Attitudes of ATA survey respondents toward screening and treatment of hypothyroidism in pregnancy. *Thyroid* **25**:368–369.
 601. Lazarus J, Brown RS, Daumerie C, Hubalewska-Dydejczyk A, Negro R, Vaidya B 2014 2014 European Thyroid Association Guidelines for the management of subclinical hypothyroidism in pregnancy and in children. *Eur Thyroid J* **3**:76–94.
 602. Vila L, Velasco I, González S, Morales F, Sánchez E, Lailla JM, Martínez-Astorquiza T, Puig-Domingo M, Grupo de Trabajo de Trastornos por Deficiencia de Yodo y Disfunción Tiroidea de la Sociedad Española de Endocrinología y Nutrición 2012 Detection of thyroid dysfunction in pregnant women: universal screening is justified. *Endocrinol Nutr* **59**:547–60.
 603. 2012 Indian Thyroid Society Guidelines for Management of Thyroid Dysfunction During Pregnancy. Clinical Practice Guidelines. Elsevier. New Delhi.
 604. Maternal Health Division, Ministry of Health and Family Welfare, Government of India 2014 National guidelines for screening of hypothyroidism during pregnancy. Available at http://nrhm.gov.in/images/pdf/programmes/maternal-health/guidelines/National_Guidelines_for_Screening_of_Hypothyroidism_during_Pregnancy.pdf
 605. Practice Committee of the American Society for Reproductive Medicine 2015 Subclinical hypothyroidism in the infertile female population: a guideline. *Fertil and Steril* **104**:545–553.
 606. Canaris GJ, Steiner JF, Ridgway EC 1997 Do traditional symptoms of hypothyroidism correlate with biochemical disease? *J Gen Intern Med* **12**:544–550.
 607. Canaris GJ, Manowitz NR, Mayor G, Ridgway EC 2000 The Colorado thyroid disease prevalence study. *Arch Intern Med* **160**:526–534.
 608. Jovanovic-Peterson L, Peterson CM 1988 De novo clinical hypothyroidism in pregnancies complicated by type I diabetes, subclinical hypothyroidism, and proteinuria: a new syndrome. *Am J Obstet Gynecol* **159**:442–446.
 609. Ladenson PW 2005 Diagnosis of hypothyroidism. In: Braverman LE, Utiger RD (eds) *Werner & Ingbar's The Thyroid: A Fundamental and Clinical Text*. Ninth edition. Lippincott, Williams & Wilkins, Philadelphia, PA, pp 857–863.
 610. Mercado G, Adelstein DJ, Saxton JP, Secic M, Larto MA, Lavertu P 2001 Hypothyroidism: a frequent event after radiotherapy and after radiotherapy with chemotherapy for patients with head and neck carcinoma. *Cancer* **92**:2892–2897.
 611. De Carlucci D Jr, Tavares MR, Obara MT, Martins LA, Hojaij FC, Cernea CR 2008 Thyroid function after unilateral total lobectomy: risk factors for postoperative hypothyroidism. *Arch Otolaryngol Head Neck Surg* **134**:1076–1079.
 612. Verloop H, Louwerens M, Schoones JW, Kievit J, Smit JW, Dekkers OM 2012 Risk of hypothyroidism following hemithyroidectomy: systematic review and meta-analysis of prognostic studies. *J Clin Endocrinol Metab* **97**:2243–2255.
 613. Strieder TG, Prummel MF, Tijssen JG, Endert E, Wiersinga WM 2003 Risk factors for and prevalence of thyroid disorders in a cross-sectional study among healthy female relatives of patients with autoimmune thyroid disease. *Clin Endocrinol (Oxf)* **59**:396–401.
 614. Manji N, Carr-Smith JD, Boelaert K, Allahabadia A, Armitage M, Chatterjee VK, Lazarus JH, Pearce SH, Vaidya B, Gough SC, Franklyn JA 2006 Influences of age, gender, smoking, and family history on autoimmune thyroid disease phenotype. *J Clin Endocrinol Metab* **91**:4873–4880.

615. Michalaki MA, Vagenakis AG, Leonardou AS, Argentou MN, Habeos IG, Makri MG, Psyrogiannis AI, Kalfarentzos FE, Kyriazopoulou VE 2006 Thyroid function in humans with morbid obesity. *Thyroid* **16**:73–78.
616. Rotondi M, Leporati P, La Manna A, Pirali B, Mondello T, Fonte R, Magri F, Chiovato L 2009 Raised serum TSH levels in patients with morbid obesity: is it enough to diagnose subclinical hypothyroidism? *Eur J Endocrinol* **160**:403–408.
617. Eskes SA, Wiersinga WM 2009 Amiodarone and thyroid. *Best Pract Res Clin Endocrinol Metab* **23**:735–751.
618. Vagenakis AG, Braverman LE 1975 Adverse effects of iodides on thyroid function. *Med Clin North Am* **59**: 1075–1088.
619. Carle A, Pedersen IB, Knudsen N, Perrild H, Ovesen L, Rasmussen LB, Laurberg P 2014 Development of autoimmune overt hypothyroidism is highly associated with live births and induced abortions but only in premenopausal women. *J Clin Endocrinol Metab* **99**:2241–2249.
620. Zimmermann MB, Boelaert K 2015 Iodine deficiency and thyroid disorders. *Lancet Diabetes Endocrinol* **3**:286–295.
621. Premawardhana LD, Parkes AB, John R, Harris B, Lazarus JH 2004 Thyroid peroxidase antibodies in early pregnancy: utility for prediction of postpartum thyroid dysfunction and implications for screening. *Thyroid* **14**:610–615.

Address correspondence to:

Elizabeth N. Pearce, MD, MSc

Section of Endocrinology, Diabetes, and Nutrition

Boston University School of Medicine

88 East Newton St., H3600

Boston, MA 02118

E-mail: elizabeth.pearce@bmc.org

This article has been cited by:

1. Federica D'Aurizio, Jürgen Kratzsch, Damien Gruson, Petra Petranović Ovčariček, Luca Giovanella. 2023. Free thyroxine measurement in clinical practice: how to optimize indications, analytical procedures, and interpretation criteria while waiting for global standardization. *Critical Reviews in Clinical Laboratory Sciences* **60**:2, 101-140. [[Crossref](#)]
2. Lennart Blomqvist, Helena F. Nyström, Margareta Hellgren, Annika Strandell. 2023. Preconceptual thyroid peroxidase antibody positivity in women with recurrent pregnancy losses may contribute to an increased risk for another miscarriage. *Clinical Endocrinology* **98**:2, 259-269. [[Crossref](#)]
3. Aida Orois, Eva López-Quesada, Gloria Jalencas, María-José Barahona, Verónica Perea. 2023. Impact of thyroid universal screening on pregnancy outcomes in real-world clinical practice. *International Journal of Gynecology & Obstetrics* **160**:2, 712-714. [[Crossref](#)]
4. Magdalena Zgliczynska, Magdalena Ostrowska, Iwona Szymusik, Michal Ciebia, Katarzyna Kosinska-Kaczynska. 2023. Maternal thyroid function in multiple pregnancies – a systematic review. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
5. Yuanzhi Chen, Zhong-Cheng Luo, Ting Zhang, Pianpian Fan, Rui Ma, Jun Zhang, Fengxiu Ouyang. 2023. Maternal Thyroid Dysfunction and Neuropsychological Development in Children. *The Journal of Clinical Endocrinology & Metabolism* **108**:2, 339-350. [[Crossref](#)]
6. Stephanie Smooke Praw, Gregory A Brent. 2023. Approach to the Patient With a Suppressed TSH. *The Journal of Clinical Endocrinology & Metabolism* **108**:2, 472-482. [[Crossref](#)]
7. Darazel B. Perez, Pranav Jani, Rajesh Maheshwari, Dharmesh Shah. 2023. Thyroid function in infants born to women with hypothyroidism: An observational study at an Australian tertiary perinatal centre. *Clinical Endocrinology* . [[Crossref](#)]
8. Bogumila Urgatz, Salman Razvi. 2023. Subclinical hypothyroidism, outcomes and management guidelines: a narrative review and update of recent literature. *Current Medical Research and Opinion* 1-47. [[Crossref](#)]
9. Kamila Tańska, Małgorzata Gietka-Czernel, Piotr Glinicki, Jarosław Kozakowski. 2023. Thyroid autoimmunity and its negative impact on female fertility and maternal pregnancy outcomes. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
10. Mariacarla Moleti, Angela Alibrandi, Maria Di Mauro, Giuseppe Paola, Laura Giovanna Perdichizzi, Roberta Granese, Annamaria Giacobbe, Angela Scilipoti, Marta Ragonese, Alfredo Ercoli, Salvatore Benvenaga, Francesco Vermiglio. 2023. Preconception Thyrotropin Levels and Thyroid Function at Early Gestation in Women With Hashimoto Thyroiditis. *The Journal of Clinical Endocrinology & Metabolism* **18**. . [[Crossref](#)]
11. Tiancheng Xie, Mingchuan Su, Jie Feng, Xiaoying Pan, Chuan Wang, Tian Tang. 2023. The reference intervals for thyroid hormones: A four year investigation in Chinese population. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
12. Cari M Kitahara, Dagrun Slettebø Daltveit, Anders Ekblom, Anders Engeland, Mika Gissler, Ingrid Glimelius, Tom Grotmol, Ylva Trolle Lagerros, Laura Madanat-Harjuoja, Tuija Männistö, Henrik Toft Sørensen, Rebecca Troisi, Tone Bjørge. 2023. Maternal Health, Pregnancy and Offspring Factors, and Maternal Thyroid Cancer Risk: A Nordic Population-Based Registry Study. *American Journal of Epidemiology* **192**:1, 70-83. [[Crossref](#)]
13. Nuha A. ElSayed, Grazia Aleppo, Vanita R. Aroda, Raveendhara R. Bannuru, Florence M. Brown, Dennis Bruemmer, Billy S. Collins, Marisa E. Hilliard, Diana Isaacs, Eric L. Johnson, Scott Kahan, Kamlesh Khunti, Jose Leon, Sarah K. Lyons, Mary Lou Perry, Priya Prahalad, Richard E. Pratley, Jane Jeffrie Seley, Robert C. Stanton, Robert A. Gabbay. 2023. 15. Management of Diabetes in Pregnancy: Standards of Care in Diabetes—2023. *Diabetes Care* **46**:Supplement_1, S254-S266. [[Crossref](#)]
14. Susan R. Rose, Ari J. Wassner, Kupper A. Wintergerst, Nana-Hawa Yayah-Jones, Robert J. Hopkin, Janet Chuang, Jessica R. Smith, Katherine Abell, Stephen H. LaFranchi, Susan R. Rose, Ari J. Wassner, Kupper A. Wintergerst, Nana-Hawa Yayah Jones, Robert J. Hopkin, Janet Chuang, Jessica R. Smith, Katherine Abell, Stephen H. LaFranchi, Kupper A. Wintergerst, Kathleen E. Bethin, Jill L. Brodsky, David H. Jelley, Bess A. Marshall, Lucy D. Mastrandrea, Jane L. Lynch, Laura Laskosz, Leah W. Burke, Timothy A. Geleske, Ingrid A. Holm, Wendy J. Introne, Kelly Jones, Michael J. Lyons, Danielle C. Monteil, Amanda B. Pritchard, Pamela Lyn Smith Trapane, Samantha A. Vergano, Kathryn Weaver, Aimee A. Alexander, Christopher Cuniff, Mary E. Null, Melissa A. Parisi, Steven J Ralson, Joan Scott, Paul Spire. 2023. Congenital Hypothyroidism: Screening and Management. *Pediatrics* **151**:1. . [[Crossref](#)]
15. Susan R. Rose, Ari J. Wassner, Kupper A. Wintergerst, Nana-Hawa Yayah-Jones, Robert J. Hopkin, Janet Chuang, Jessica R. Smith, Katherine Abell, Stephen H. LaFranchi, Susan R. Rose, Ari J. Wassner, Kupper A. Wintergerst, Nana-Hawa Yayah Jones, Robert J. Hopkin, Janet Chuang, Jessica R. Smith, Katherine Abell, Stephen H. LaFranchi, Kupper A. Wintergerst, Kathleen E. Bethin, Brittany Bruggeman, Jill L. Brodsky, David H. Jelley, Bess A. Marshall, Lucy D. Mastrandrea, Jane L. Lynch, Laura Laskosz, Leah W. Burke, Timothy A. Geleske, Ingrid A. Holm, Wendy J. Introne, Kelly Jones, Michael J. Lyons, Danielle C. Monteil, Amanda B. Pritchard, Pamela Lyn Smith Trapane, Samantha A. Vergano, Kathryn Weaver, Aimee A. Alexander,

- Christopher Cunniff, Mary E. Null, Melissa A. Parisi, Steven J Ralson, Joan Scott. 2023. Congenital Hypothyroidism: Screening and Management. *Pediatrics* **151**:1. . [\[Crossref\]](#)
16. Jörg Bojunga, Oya Güngöze, Ludwig Hofbauer, Elisa Müller. 2023. Schilddrüsenerkrankungen in der Schwangerschaft. *DMW - Deutsche Medizinische Wochenschrift* **148**:01/02, 17-26. [\[Crossref\]](#)
 17. Shivangi Nikhil Dwivedi, Tejas Kalaria, Harit Buch. 2023. Thyroid autoantibodies. *Journal of Clinical Pathology* **76**:1, 19-28. [\[Crossref\]](#)
 18. Francilene Maria Azevedo, Almeida Abudo Leite Machamba, Aline Carare Candido, Carina Aparecida Pinto, Sílvia Oliveira Lopes, Mariana de Souza Macedo, Sarah Aparecida Vieira Ribeiro, Sílvia Eloiza Priore, Sylvia do Carmo Castro Franceschini. 2023. Correlation Between Drinking Water and Iodine Status: a Systematic Review and Meta-analysis. *Biological Trace Element Research* **201**:1, 129-138. [\[Crossref\]](#)
 19. Rishalaiti Tayier, Chenchen Wang, Pinjiang Ma, Yimu Yuan, Yuxia Zhang, Shunhua Wu, Ling Zhang. 2023. Iodine Nutritional Status of Pregnant Women After 14 Years of Lipiodol Supplementation: a Cross-Sectional Study in Historically Iodine-Deficient Areas of China. *Biological Trace Element Research* **201**:1, 14-22. [\[Crossref\]](#)
 20. Jiang-Nan Wu, Faustino R. Pérez-López, Ting Peng, Feng Xie, Ming-Qing Li. 2023. Association of Early Pregnancy Free and Total Triiodothyronine With the Subsequent Risk of Gestational Diabetes Mellitus. *Endocrine Practice* **29**:1, 40-47. [\[Crossref\]](#)
 21. Roxana-Elena Bohîlţea, Bianca-Margareta Mihai, Elena Szini, Ileana-Alina Şucaliuc, Corin Badiu. 2023. Diagnosis and Management of Fetal and Neonatal Thyrotoxicosis. *Medicina* **59**:1, 36. [\[Crossref\]](#)
 22. Yan Zhao, Yongbo Zhao, Yijun Zhang, Mengyuan Li, Xin Su, Yicheng Zhou, Ziyi Zhang, Liping Jin. 2023. Association of COVID-19 vaccination before conception with maternal thyroid function during early pregnancy: A single-center study in China. *Journal of Medical Virology* **95**:1. . [\[Crossref\]](#)
 23. R. Mazzilli, S. Medenica, A. M. Di Tommaso, G. Fabozzi, V. Zamponi, D. Cimadomo, L. Rienzi, F. M. Ubaldi, M. Watanabe, A. Faggiano, S. La Vignera, G. Defeudis. 2023. The role of thyroid function in female and male infertility: a narrative review. *Journal of Endocrinological Investigation* **46**:1, 15-26. [\[Crossref\]](#)
 24. Hyeong-Moo Shin, Jiwon Oh, Rebecca J. Schmidt, Elizabeth N. Pearce. 2022. Prenatal Exposure to Per- and Polyfluoroalkyl Substances, Maternal Thyroid Dysfunction, and Child Autism Spectrum Disorder. *Endocrinology and Metabolism* **37**:6, 819-829. [\[Crossref\]](#)
 25. Mehmet ERDOĞAN, Günel BAYRAMOVA, Hatice ÖZİŞİK, Aslı SUNER. 2022. Investigation on Demographic Characteristics of Pregnant Patients with Thyroid Dysfunction- Ege University Sample. *Medical Journal of Western Black Sea* **6**:3, 345-351. [\[Crossref\]](#)
 26. F. Magri, C. Bellingeri, I. De Maggio, L. Croce, F. Coperchini, M. Rotondi, L. Chiovato, A. Spinillo, F. Beneventi. 2022. A first-trimester serum TSH in the 4–10 mIU/L range is associated with obstetric complications in thyroid peroxidase antibody-negative women. *Journal of Endocrinological Investigation* **27**. . [\[Crossref\]](#)
 27. Piotr Kocelak, Aleksander J. Owczarek, Agnieszka Wikarek, Natalia Ogarek, Paulina Oboza, Małgorzata Sieja, Anna Szyszka, Izabela Rozmus-RogóŹ, Monika Puzianowska-KuŹnicka, Magdalena Olszanecka-Glinianowicz, Jerzy Chudek. 2022. Anti-thyroid antibodies in the relation to TSH levels and family history of thyroid diseases in young Caucasian women. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)
 28. Susmita Reddy Karri, Priya Susan Roy, Nirjhar Nandi, Vasant Shenoy, David Watson. 2022. Management of fetal hyperthyroidism caused by persistent autoimmune antibodies in a case of previously treated maternal Graves' disease. *Obstetric Medicine* **82**, 1753495X2211463. [\[Crossref\]](#)
 29. Sofia Gill, Versha Cheed, Victoria A H Morton, Dayna Gill, Kristien Boelaert, Shiao Chan, Arri Coomarasamy, Rima K Dhillon-Smith. 2022. Evaluating the Progression to Hypothyroidism in Preconception Euthyroid Thyroid Peroxidase Antibody-Positive Women. *The Journal of Clinical Endocrinology & Metabolism* **108**:1, 124-134. [\[Crossref\]](#)
 30. . Medications and Breastfeeding . [\[Crossref\]](#)
 31. Sibel Ozler, Efser Oztas, Aysegul Kebapcilar, Ali Turhan Caglar. 2022. The role of thyroid-stimulating hormone and thyroglobulin antibody in abnormally invasive placenta. *The Journal of Maternal-Fetal & Neonatal Medicine* **35**:25, 5108-5116. [\[Crossref\]](#)
 32. Mohamad Mustafa, Elsheikh Ali, Anne McGowan, Laura McCabe, Laszlo Hegedüs, Roberto Attanasio, Endre V. Nagy, Enrico Papini, Petros Perros, Carla Moran. 2022. Use of Thyroid Hormones in Hypothyroid and Euthyroid Patients: A THESIS questionnaire survey of members of the Irish Endocrine Society. *Irish Journal of Medical Science (1971 -)* **99**. . [\[Crossref\]](#)
 33. Costanzo Moretti, Natalia Lazzarin, Elena Vaquero, Alessandro Dal Lago, Luisa Campagnolo, Herbert Valensise. 2022. A practical approach to the management of thyroid dysfunction during pregnancy. *Gynecological Endocrinology* **19**, 1-7. [\[Crossref\]](#)

34. Louise Brough. 2022. Improving Iodine Status in Lactating Women: What Works?. *Current Nutrition Reports* 11:4, 592-599. [[Crossref](#)]
35. Samuel Chigbo Obiegbusi, Xiao Jing Dong, Mingyu Deng, Chidera Nneji Obiegbusi, Yin Yang, Xinru Deng. 2022. Assessing the Outcome of the Management of Thyroid Dysfunction in Pregnancy. *SN Comprehensive Clinical Medicine* 4:1. . [[Crossref](#)]
36. Seyit Ahmet Erol, Ali Turhan Caglar, Yaprak Engin Ustun, A. Seval Ozgu Erdinc. 2022. Evaluation of Perinatal and Neonatal Outcomes in Pregnant Women with Thyroid Autoantibody Positivity (Anti-thyroglobulin and Anti-thyroid Peroxidase) Due to Thyroiditis. *SN Comprehensive Clinical Medicine* 4:1. . [[Crossref](#)]
37. Delphine Drui, Claire Briet, Carole Guerin, Alexandre Lugat, Francoise Borson-Chazot, Solange Grunenwald. 2022. SFE-AFCE-SFMN 2022 Consensus on the management of thyroid nodules : Thyroid nodules and pregnancy. *Annales d'Endocrinologie* 83:6, 435-439. [[Crossref](#)]
38. Sarika Tyagi. 2022. Assessment of iodine deficiency disorders among pregnant women residing in an urban slum of west Delhi. *Human Nutrition & Metabolism* 30, 200172. [[Crossref](#)]
39. Sima Nazarpour, Fahimeh Ramezani Tehrani, Mina Amiri, Maryam Rahmati, Fereidoun Azizi. 2022. Maternal subclinical hyperthyroidism and adverse pregnancy outcomes considering the iodine status: Tehran thyroid and pregnancy study. *Journal of Trace Elements in Medicine and Biology* 74, 127063. [[Crossref](#)]
40. Wen Chen, Wei Wang, Min Gao, Yanting Chen, Wenxing Guo, Shuyao Dong, Hao Sun, Ziyun Pan, Elizabeth N. Pearce, Long Tan, Jun Shen, Wanqi Zhang. 2022. Iodine Intakes of 550 $\mu\text{g}/\text{day}$ are Not Recommended during Pregnancy: A Balance Study. *The Journal of Nutrition* 144. . [[Crossref](#)]
41. Layal Chaker, Salman Razvi, Isabela M. Bensenor, Fereidoun Azizi, Elizabeth N. Pearce, Robin P. Peeters. 2022. Hypothyroidism. *Nature Reviews Disease Primers* 8:1. . [[Crossref](#)]
42. Charles Bitamazire Businge, Benjamin Longo-Mbenza, Andre Pascal Kengne. 2022. Iodine deficiency in pregnancy along a concentration gradient is associated with increased severity of preeclampsia in rural Eastern Cape, South Africa. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
43. Júlia Siscart, Míriam Orós, M. Catalina Serna, Dani Perejón, Leonardo Galván, Marta Ortega. 2022. Adherence to treatment for hypothyroidism in pregnancy and relationship with thyrotropin control: a retrospective observational cohort study. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
44. Ping Zhou, Qiuping Yao, Qiaohang Zhao, Lihua Yang, Ya Yu, Jilai Xie, Chun Feng, Liming Zhou, Min Jin. 2022. IVF/ICSI outcomes of euthyroid infertile women with thyroid autoimmunity: does treatment with aspirin plus prednisone matter?. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
45. Lingna Han, Xin Yang, Wen Wang, Xueliang Yang, Lina Dong, Shumei Lin, Jianguo Li, Xiaojing Liu. 2022. Cord blood metabolomics reveals gestational metabolic disorder associated with anti-thyroid peroxidase antibodies positivity. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
46. Rana Turkal, Cem Armağan Turan, Onur Elbasan, Serenay Aytan, Burcu Çakmak, Büşra Gözaydinoğlu, Duygu Ceyda Takır, Ozan Ünlü, Günel Bahramzada, Ahmet Faruk Tekin, Tülay Çevlik, Esra Esim Büyükbayrak, Önder Şirikçi, Hülya Gözü, Goncagül Haklar. 2022. Accurate interpretation of thyroid dysfunction during pregnancy: should we continue to use published guidelines instead of population-based gestation-specific reference intervals for the thyroid-stimulating hormone (TSH)?. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
47. Cheng Peng, Weijie Sun, Lixin Fan, Li Li, Xiaojiao Zhang, Ying Gao, Xinlin Hou. 2022. A hypothyroid mother after subtotal thyroidectomy delivered a newborn with hyperthyroidism from fetal stage: a case report. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
48. Jiang-Nan Wu, Ting Peng, Feng Xie, Ming-Qing Li. 2022. Association of thyroid dysfunction and autoantibody positivity with the risk of preterm birth: a hospital-based cohort study. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
49. Ying Wang, Sha Lu, Xianrong Xu, Lijun Zhang, Jun Yang, Wensheng Hu. 2022. The interactive effects of pre-pregnancy body mass index, thyroid function, and blood lipid levels on the risk of gestational diabetes mellitus: a crossover analysis. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
50. Min Wu, Cheng Chi, Yuxi Yang, Shan Guo, Tianhe Li, Muqing Gu, Tingting Zhang, Huimin Gao, Ruixia Liu, Chenghong Yin. 2022. Dynamics of gut microbiota during pregnancy in women with TPOAb-positive subclinical hypothyroidism: a prospective cohort study. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]
51. Yuxi Yang, Huabing Yuan, Xueran Wang, Zheng Zhang, Ruixia Liu, Chenghong Yin. 2022. Significance of levothyroxine treatment on serum lipid in pregnant women with subclinical hypothyroidism. *BMC Pregnancy and Childbirth* 22:1. . [[Crossref](#)]

52. Ping Li, Jinhui Cui, Ling Li, Xinjuan Chen, Liping Ouyang, Jianhui Fan, Shuo Lin. 2022. Association between isolated maternal hypothyroxinemia during the first trimester and adverse pregnancy outcomes in Southern Chinese women: a retrospective study of 7051 cases. *BMC Pregnancy and Childbirth* 22:1. . [\[Crossref\]](#)
53. Ying Zhao, Wanxian Zhang, Xiuying Tian. 2022. Analysis of risk factors of early intraventricular hemorrhage in very-low-birth-weight premature infants: a single center retrospective study. *BMC Pregnancy and Childbirth* 22:1. . [\[Crossref\]](#)
54. Katia Vella, Sandro Vella, C. Savona-Ventura, J. Vassallo. 2022. Thyroid dysfunction in pregnancy - a retrospective observational analysis of a Maltese cohort. *BMC Pregnancy and Childbirth* 22:1. . [\[Crossref\]](#)
55. Geng-dong Chen, Xiao-Yan Gou, Ting-ting Pang, Peng-sheng Li, Zi-xing Zhou, Dong-xin Lin, Da-zhi Fan, Xiao-ling Guo, Li-juan Wang, Zheng-ping Liu. 2022. Associations between thyroid function and gestational diabetes mellitus in Chinese pregnant women: a retrospective cohort study. *BMC Endocrine Disorders* 22:1. . [\[Crossref\]](#)
56. Aimei Dong, Youyuan Huang, Yucheng Huang, Bing Jia. 2022. Effects of calorie intake and sampling time on thyroid stimulating hormone concentration. *BMC Endocrine Disorders* 22:1. . [\[Crossref\]](#)
57. Jan Jiskra, Jan Paleček, Roberto Attanasio, Laszlo Hegedüs, Endre V. Nagy, Enrico Papini, Petros Perros, Roberto Negro, Michal Kršek. 2022. Use of thyroid hormones in hypothyroid and euthyroid patients: a 2020 THESIS questionnaire survey of members of the Czech Society of Endocrinology. *BMC Endocrine Disorders* 22:1. . [\[Crossref\]](#)
58. Tianqi Liu, Xiuying Zhang, Li Long, Lingli Zhou, Jing Chen, Meng Li, Ying Gao, Xianghai Zhou, Xueyao Han, Linong Ji. 2022. Clinical evaluation of an automated TSI bridge immunoassay in the diagnosis of Graves' disease and its relationship to the degree of hyperthyroidism. *BMC Endocrine Disorders* 22:1. . [\[Crossref\]](#)
59. Mahboubeh Hajifoghaha, Saeed Hosseini Teshnizi, Sedighe Forouhari, Mohammad Hossein Dabbaghmanesh. 2022. Association of thyroid function test abnormalities with preeclampsia: a systematic review and meta-analysis. *BMC Endocrine Disorders* 22:1. . [\[Crossref\]](#)
60. Grace Mengqin Ge, Edmund C. L. Cheung, Kenneth K. C. Man, Patrick Ip, Wing Cheong Leung, Gloria H. Y. Li, Annie W. C. Kung, Ching-Lung Cheung, Ian C. K. Wong. 2022. Association of maternal levothyroxine use during pregnancy with offspring birth and neurodevelopmental outcomes: a population-based cohort study. *BMC Medicine* 20:1. . [\[Crossref\]](#)
61. Xiao Chen, Chunfeng Wu, Zhengyuan Wang, Chunxiang Wu, Yan Guo, Xiaoxia Zhu, Yan Ping Hu, Zehuan Shi, Qi Song, Xueying Cui, Jin Su, Jiajie Zang. 2022. Iodine nutrition status and thyroid autoimmunity during pregnancy: a cross-sectional study of 4635 pregnant women. *Nutrition Journal* 21:1. . [\[Crossref\]](#)
62. Yunyi Xu, Yiqi Zhao, Xiaoqin Xu, Qiqi Yan, Liwei Yang. 2022. Serum lipid profile in relation to free thyroxine and the effect of levothyroxine treatment on lipids in patients with isolated hypothyroxinemia during pregnancy: a single-center retrospective study. *Lipids in Health and Disease* 21:1. . [\[Crossref\]](#)
63. Shaoyuan Xu, Ying Zhang, Cancan Qiang, Changjun Zhang. 2022. Effect of TSH on oocyte maturation of PCOS patients with normal thyroid function in IVF. *Reproductive Biology and Endocrinology* 20:1. . [\[Crossref\]](#)
64. Massimo Giusti, Miranda Mittica. 2022. Evaluation of anti-Müllerian hormone in pre-menopausal women stratified according to thyroid function, autoimmunity and age. *Thyroid Research* 15:1. . [\[Crossref\]](#)
65. Stig Andersen, Jesper Karmisholt, Niels Henrik Bruun, Johannes Riis, Paneeraq Noahsen, Louise Westergaard, Stine Linding Andersen. 2022. Interpretation of TSH and T4 for diagnosing minor alterations in thyroid function: a comparative analysis of two separate longitudinal cohorts. *Thyroid Research* 15:1. . [\[Crossref\]](#)
66. Zareen Kiran, Wardah Khalid, Aisha Sheikh, Najmul Islam. 2022. Levothyroxine dosages during pregnancy among hypothyroid women. An experience from a tertiary care center of Karachi, Pakistan, based on data from Maternal Hypothyroidism on Pregnancy Outcomes Study (MHPO-5). *BMC Research Notes* 15:1. . [\[Crossref\]](#)
67. Stine Linding Andersen, Niels Henrik Bruun, Peter Astrup Christensen, Simon Lykkeboe, Aase Handberg, Annebirthe Bo Hansen, Maja Hjelm Lundgaard, Louise Knøsgaard, Nanna Maria Uldall Torp, Allan Carlé, Jesper Karmisholt, Inge Bülow Pedersen, Peter Vestergaard, Stig Andersen. 2022. Cut-offs for thyroid peroxidase and thyroglobulin antibodies in early pregnancy. *European Thyroid Journal* 11:6. . [\[Crossref\]](#)
68. Simon Shenhav, Carlos Benbassat, Dov Gefel, Shmuel Zangen, Shani R. Rosen, Yael Avrahami-Benyounes, Shlomo Almashanu, Ludmila Groisman, Efrat Rorman, Shlomo Fytlovich, Eyal Y. Anteby, Yaniv S. Ovadia. 2022. Can Mild-to-Moderate Iodine Deficiency during Pregnancy Alter Thyroid Function? Lessons from a Mother-Newborn Cohort. *Nutrients* 14:24, 5336. [\[Crossref\]](#)
69. Clelia CIPOLLA, Marta TEDESCO. 2022. Post-partum thyroid disease: differential diagnosis and management. *Minerva Endocrinology* 47:4. . [\[Crossref\]](#)
70. Liujiao Cao, Liang Yao, Wenbo He, Liangying Hou, Zhe Yin, Dan Wang, Ka Li. 2022. Methodological quality in guidelines for enhanced recovery after surgery was suboptimal. *Journal of Clinical Epidemiology* 152, 151-163. [\[Crossref\]](#)

71. Louise Knøsgaard, Stig Andersen, Annebirthe Bo Hansen, Peter Vestergaard, Stine Linding Andersen. 2022. Maternal hypothyroidism and adverse outcomes of pregnancy. *Clinical Endocrinology* **34**. . [\[Crossref\]](#)
72. Constance Nontsikelelo Gubu-Ntaba, Vulikhaya Mpumlwana, Nandipha Mizpa Sotobe-Mbana, Martha Mayer, Chukwuma Ogbonna Ekpebegh, Charles Bitamazire Businge. The Impact of Hyperthyroidism on Fertility, Maternal, Foetal and Perinatal Outcomes in the Era of Iodine Fortification . [\[Crossref\]](#)
73. Divya M Mathews, Jane M Peart, Robert G Sim, Neil P Johnson, Susannah O'Sullivan, José G B Derraik, Paul L Hofman. 2022. The SELFI Study: Iodine Excess and Thyroid Dysfunction in Women Undergoing Oil-Soluble Contrast Hysterosalpingography. *The Journal of Clinical Endocrinology & Metabolism* **107**:12, 3252-3260. [\[Crossref\]](#)
74. Scott A Rivkees. 2022. Approach to the Patient: Management and the Long-term Consequences of Graves' Disease in Children. *The Journal of Clinical Endocrinology & Metabolism* **107**:12, 3408-3417. [\[Crossref\]](#)
75. L. Zhuo, Z. Wang, Y. Yang, Z. Liu, S. Wang, Y. Song. 2022. Obstetric and offspring outcomes in isolated maternal hypothyroxinaemia: a systematic review and meta-analysis. *Journal of Endocrinological Investigation* **105**. . [\[Crossref\]](#)
76. Aysegul Atalay, Berhan Besimoglu, Selcan Sinaci, Ecem Kaya, Onur Ozkavak, Fatma Doga Ocal, Ayse Seval Ozgu-Erdinc, Dilek Sahin. 2022. The impact of covid-19 on thyroid function tests in pregnancy. *Endocrine* **181**. . [\[Crossref\]](#)
77. Katherina Vicky Gantenbein, Christina Kanaka-Gantenbein. 2022. Highlighting the trajectory from intrauterine growth restriction to future obesity. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)
78. Stavroula A. Paschou, Evanthia Bletsas, Maria Papazisi, Nikoletta Mili, Fotini Kanouta, Georgia N. Kassi, Theodora Psaltopoulou, Dimitrios G. Goulis, Irene Lambrinoudaki. 2022. Screening and management of major endocrinopathies during pregnancy: an update. *Endocrine* **22**. . [\[Crossref\]](#)
79. Mandy Gfeller, Gentiane Colque, Peter A. Kopp. 2022. Iodine content of frequently used prenatal and adult multivitamins in Switzerland. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)
80. Natalia Genere. 2022. Who Is the Ideal Pregnant Patient with Graves' Disease to Discontinue Antithyroid Drugs?. *Clinical Thyroidology* **34**:11, 479-482. [\[Citation\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
81. Małgorzata Karbownik-Lewińska, Jan Stępnia, Paulina Iwan, Andrzej Lewiński. 2022. Iodine as a potential endocrine disruptor —a role of oxidative stress. *Endocrine* **78**:2, 219-240. [\[Crossref\]](#)
82. Lluís García González, Luis García Pascual. 2022. Clinical usefulness of thyroid ultrasonography in patients with primary hypothyroidism. *Endocrinología, Diabetes y Nutrición (English ed.)* **69**:9, 686-693. [\[Crossref\]](#)
83. Inés Velasco, Lluís Vila, Maria Goya, Amelia Oleaga, Maria de la Calle, Fco. Javier Santamaria. 2022. Executive summary of the SEEN (Sociedad Española de Endocrinología y Nutrición[Spanish Society of Endocrinology and Nutrition])-SEGO (Sociedad Española de Ginecología y Obstetricia [Spanish Society of Gynaecology and Obstetrics]) consensusdocument on the management of thyroid dysfunction during pregnancy. *Endocrinología, Diabetes y Nutrición (English ed.)* **27**. . [\[Crossref\]](#)
84. Lluís García González, Luis García Pascual. 2022. Utilidad clínica de la ecografía tiroidea en los pacientes con hipotiroidismo primario. *Endocrinología, Diabetes y Nutrición* **69**:9, 686-693. [\[Crossref\]](#)
85. Mariza Kampouri, Klara Gustin, Mia Stråvik, Malin Barman, Michael Levi, Vasiliki Daraki, Bo Jacobsson, Anna Sandin, Ann-Sofie Sandberg, Agnes E. Wold, Marie Vahter, Maria Kippler. 2022. Association of maternal urinary fluoride concentrations during pregnancy with size at birth and the potential mediation effect by maternal thyroid hormones: The Swedish NICE birth cohort. *Environmental Research* **214**, 114129. [\[Crossref\]](#)
86. P. Deruelle, L. Sentilhes, L. Ghesquière, R. Desbrière, G. Ducarme, L. Attali, A. Jarnoux, F. Artzner, A. Tranchant, T. Schmitz, M.-V. Sénat. 2022. Consensus formalisé d'experts du Collège national des gynécologues et obstétriciens français : prise en charge des nausées et vomissements gravidiques et de l'hyperémèse gravidique. *Gynécologie Obstétrique Fertilité & Sénologie* **50**:11, 700-711. [\[Crossref\]](#)
87. Jia Liu, Xiaona Chang, Qiu Wang, Xiaoyu Ding, Tao Jiang, Guang Wang. 2022. Magnetic resonance T1-mapping quantitatively assesses the severity of thyroid destruction in patients with autoimmune thyroiditis. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)
88. . Thyroid Gland 99-187. [\[Crossref\]](#)
89. Yanmin Zhong, Sujian Peng, Qiuju Chen, Dongyin Huang, Guanglan Zhang, Zehong Zhou. 2022. Preconceptional thyroid stimulating hormone level and fecundity: a community-based cohort study of time to pregnancy. *Fertility and Sterility* **32**. . [\[Crossref\]](#)
90. Rodrigo Ayala-Yáñez, Andres Villapando-Basaldúa, Alejandra Halfter-Mijares, Dan Morgenstern-Kaplan, Ixel Lorena Ruiz-Cabrera, Guadalupe Herrera-Fernández. 2022. Pregnancy hypothyroidism incidence and complications using the 2011 and 2017 ATA cutoff values. Experience at a maternity hospital in a densely populated area in Mexico City. *Journal of Perinatal Medicine* **50**:8, 1135-1141. [\[Crossref\]](#)

91. Naziye GÜRKAN, Halime ÇALI ÖZTÜRK. 2022. The findings about relationship between autoimmune thyroid disease and first-trimester aneuploidy results. *Journal of Health Sciences and Medicine* 5:6, 1661-1665. [[Crossref](#)]
92. Jennifer K. Litton. Cancer and Pregnancy 1-11. [[Crossref](#)]
93. Mariano Mascarenhas, Yadava Jeve, Lukasz Polanski, Abigail Sharpe, Ephra Yasmin, Harish M. Bhandari. 2022. Management of recurrent implantation failure: British Fertility Society policy and practice guideline. *Human Fertility* 25:5, 813-837. [[Crossref](#)]
94. Laura Croce, Luca Chiovato, Massimo Tonacchera, Elena Petrosino, Maria Laura Tanda, Mariacarla Moleti, Flavia Magri, Antonella Olivieri, Elizabeth N. Pearce, Mario Rotondi. 2022. Iodine status and supplementation in pregnancy: an overview of the evidence provided by meta-analyses. *Reviews in Endocrine and Metabolic Disorders* 22. . [[Crossref](#)]
95. Viviane Rosado Negreiros d'Assunção, Erik Montagna, Luis Eduardo Negreiros d'Assunção, Maria Madalena Pessoa Caldas, Denise Maria Christofolini, Caio Parente Barbosa, Ricardo Andre Medeiros Negreiros, Antonio Simone Laganà, Renato de Oliveira, Bianca Bianco. 2022. Effect of thyroid function on assisted reproduction outcomes in euthyroid infertile women: A single center retrospective data analysis and a systematic review and meta-analysis. *Frontiers in Endocrinology* 13. . [[Crossref](#)]
96. K. Aaron Geno, Robert D. Nerenz. 2022. Evaluating thyroid function in pregnant women. *Critical Reviews in Clinical Laboratory Sciences* 59:7, 460-479. [[Crossref](#)]
97. Victor J. Pop, Lianne P. Hulsbosch, Myrthe G.B.M. Boekhorst, Maarten A.C. Broeren, Johannes G. Krabbe, Wilmar M. Wiersinga. 2022. Hypothyroid Symptoms Throughout Pregnancy Are Predominantly Associated with Thyroxine and Not with Thyrotropin Concentrations. *Thyroid* 32:10, 1249-1258. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
98. Tatiana Fedorova, Lilah F. Morris-Wiseman. 2022. Delayed (>3 Months) Postoperative Radioactive Iodine Ablation Does Not Impact Clinical Response or Survival in Differentiated Thyroid Cancers. *Clinical Thyroidology* 34:10, 456-459. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
99. Xueying Cui, Huiting Yu, Zhengyuan Wang, Hai Wang, Zehuan Shi, Wei Jin, Qi Song, Changyi Guo, Hongmei Tang, Jiajie Zang. 2022. No Association Was Found Between Mild Iodine Deficiency During Pregnancy and Pregnancy Outcomes: a Follow-up Study Based on a Birth Registry. *Biological Trace Element Research* 200:10, 4267-4277. [[Crossref](#)]
100. C. Matta Coelho, J. Guimarães, I. Bracchi, N. Xavier Moreira, C. Pinheiro, P. Ferreira, D. Pestana, I. Barreiros Mota, A. Cortez, C. Prucha, C. Martins, E. Pinto, A. Almeida, C. Delerue-Matos, C. C. Dias, A. Moreira-Rosário, L. F. Ribeiro de Azevedo, V. Cruz Fernandes, C. Ramalho, C. Calhau, A.-L. Brantsæter, J. Costa Leite, E. Keating. 2022. Noncompliance to iodine supplementation recommendation is a risk factor for iodine insufficiency in Portuguese pregnant women: results from the IoMum cohort. *Journal of Endocrinological Investigation* 45:10, 1865-1874. [[Crossref](#)]
101. Alberto López García-Franco, José Antonio Baeyens Fernández, M. José Iglesias Piñeiro, Pablo Alonso Coello, Cristina Ruiz Cabello, Ana Pereira Iglesias, Jacinta Landa Goñi. 2022. Actividades preventivas en la mujer. Actualización PAPPS 2022. *Atención Primaria* 54, 102471. [[Crossref](#)]
102. Sophie Jansen, Genevieve Leduc-Robert, Faten F. AbdelHafez, Arianne Albert, Ulrike Mayer, Mohamed A. Bedaiwy. 2022. Temporal trends in thyroid-stimulating hormone and live birth rate in subclinical hypothyroid patients in a recurrent pregnancy loss population. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 277, 21-26. [[Crossref](#)]
103. Natália Campos Gonçalves Scherr, Anelise Impelizeri Nogueira, Kamilla Maria Araújo Brandão Rajão, Henrique Vitor Leite. 2022. Nutritional Status of Iodine in a Group of Pregnant Women from the State of Minas Gerais Correlated with Neonatal Thyroid Function. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics* 44:10, 909-914. [[Crossref](#)]
104. Sara Toassa Gomes Solha, Rosiane Mattar, Patrícia de Fátima dos Santos Teixeira, Maria Izabel Chiamolera, Carlos Alberto Maganha, Alberto Carlos Moreno Zaconeta, Renato Teixeira Souza. 2022. Screening, diagnosis and management of hypothyroidism in pregnancy. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics* 44:10, 999-1010. [[Crossref](#)]
105. Carol Chiung-Hui Peng, Spyridoula Maraka, Elizabeth N. Pearce. 2022. Under and overtreatment with thyroid hormone replacement during pregnancy. *Current Opinion in Endocrinology, Diabetes & Obesity* 29:5, 474-482. [[Crossref](#)]
106. Suvi Turunen, Marja Väärasmäki, Riitta Marttila, Maarit K. Leinonen, Mika Gissler, Tuija Männistö, Eila Suvanto. 2022. Indications for intensive care unit treatment among neonates born to mothers with thyroid disease: A population-based cohort study. *Acta Obstetrica et Gynecologica Scandinavica* 101:10, 1093-1101. [[Crossref](#)]
107. Dongmei Sun, Janine R. Hutson, Facundo Garcia-Bournissen. 2022. Drug therapy during pregnancy. *British Journal of Clinical Pharmacology* 88:10, 4247-4249. [[Crossref](#)]
108. Mark R Chester, Anushka Tirlapur, Kanna Jayaprakasan. 2022. Current management of recurrent pregnancy loss. *The Obstetrician & Gynaecologist* 24:4, 260-271. [[Crossref](#)]

109. Leqi He, Xiaoying Li, Zaoping Chen, Wei Wang, Kai Wang, Xinmei Huang, Qian Yang, Wencai Ke, Jun Liu, Bingbing Zha. 2022. High concentration of estradiol has a negative correlation with free thyroxine during the second trimester of pregnancy. *Endocrine Connections* **11**:10. . [[Crossref](#)]
110. Joris A J Osinga, Arash Derakhshan, Glenn E Palomaki, Ghalia Ashoor, Tuija Männistö, Spyridoula Maraka, Liangmiao Chen, Sofie Bliddal, Xuemian Lu, Peter N Taylor, Tanja G M Vrijkotte, Fang-Biao Tao, Suzanne J Brown, Farkhanda Ghafoor, Kris Poppe, Flora Veltri, Lida Chatzi, Bijay Vaidya, Maarten A C Broeren, Beverley M Shields, Sachiko Itoh, Lorena Mosso, Polina V Popova, Anna D Anopova, Reiko Kishi, Ashraf Aminorroaya, Maryam Kianpour, Abel López-Bermejo, Emily Oken, Amna Pirzada, Marina Vafeiadi, Wichor M Bramer, Eila Suvanto, Jun Yoshinaga, Kun Huang, Judit Bassols, Laura Boucai, Ulla Feldt-Rasmussen, Elena N Grineva, Elizabeth N Pearce, Erik K Alexander, Victor J M Pop, Scott M Nelson, John P Walsh, Robin P Peeters, Layal Chaker, Kypros H Nicolaides, Mary E D'Alton, Tim I M Korevaar. 2022. TSH and FT4 Reference Intervals in Pregnancy: A Systematic Review and Individual Participant Data Meta-Analysis. *The Journal of Clinical Endocrinology & Metabolism* **107**:10, 2925-2933. [[Crossref](#)]
111. Andreea Borlea, Laura Cotoi, Corina Paul, Felix Bende, Dana Stoian. Elastography Methods in the Prediction of Malignancy in Thyroid Nodules . [[Crossref](#)]
112. Grigorios Panagiotou, Peter N Taylor, D Aled Rees, Onyebuchi E Okosieme. 2022. Late offspring effects of antenatal thyroid screening. *British Medical Bulletin* **143**:1, 16-29. [[Crossref](#)]
113. Na Zhou, Feng Zhang, Cyril Kuriakose, Dwayne Gordon, Amay Parikh. 2022. A Case of Transient Gestational Hyperthyroidism Complicated With Acute Liver Injury Successfully Treated With Plasma Exchange. *Cureus* **97**. . [[Crossref](#)]
114. Nian Guo, Meng Xue, Zhen Liang. 2022. Advances in the differential diagnosis of transient hyperthyroidism in pregnancy and Graves' disease. *Archives of Gynecology and Obstetrics* **305**. . [[Crossref](#)]
115. Francesca Orsolini, Elena Gianetti, Chiara Terrenzio, Lucia Montanelli, Elena Benelli, Brunella Bagattini, Emilio Fiore, Massimo Tonacchera. 2022. Thyroid Function Rather Than Thyroid Antibodies Affects Pregnancy and Perinatal Outcomes: Results of a Prospective Study. *The Journal of Clinical Endocrinology & Metabolism* **48**. . [[Crossref](#)]
116. Benjamin J. Gigliotti. 2022. Are High-Normal TSH Values Associated with Diminished Ovarian Reserve and Poor Response to Assisted Reproductive Technology?. *Clinical Thyroidology* **34**:9, 392-395. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
117. Josie Rossitto, Ruba Riachy, Nisarg Shah, Meagan Benson, Shannon Clark. 2022. Fertility and Radioactive Iodine: A Case of an Unforeseen Complication. *Clinical Thyroidology* **34**:9, 420-425. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
118. M. Amiri, S. Nazarpour, F. Ramezani Tehrani, A. Sheidaei, F. Azizi. 2022. The targeted high-risk case-finding approach versus universal screening for thyroid dysfunction during pregnancy: thyroid-stimulating hormone (TSH) and/or thyroid peroxidase antibody (TPOAb) test?. *Journal of Endocrinological Investigation* **45**:9, 1641-1651. [[Crossref](#)]
119. Yuchao Zhang, Jia Peng, Yanli Liu, Wenbin Wu, Xingling Wang, Liting Jia, Yichun Guan. 2022. The Impact of High-Normal TSH Levels on Reproductive Outcomes in Women Undergoing ART Treatment: a Systematic Review and Meta-analysis. *Reproductive Sciences* **29**:9, 2440-2451. [[Crossref](#)]
120. Richard Christian Jensen, Dorte Glintborg, Clara Amalie Gade Timmermann, Flemming Nielsen, Henriette Boye, Jeppe Buur Madsen, Niels Bilenberg, Philippe Grandjean, Tina Kold Jensen, Marianne S. Andersen. 2022. Higher free thyroxine associated with PFAS exposure in first trimester. The Odense Child Cohort. *Environmental Research* **212**, 113492. [[Crossref](#)]
121. Raviv Allon, Tal Schiller, Yuval Ziv, Yonatan Lahav, Oded Cohen, Taiba Zornitzki. 2022. Posthemithyroidectomy Pregnancy Thyroid Function Surveillance: Frequency, Adherence, and Guideline Impact. *Endocrine Practice* **28**:9, 847-852. [[Crossref](#)]
122. Odessa P. Hamidi, Linda A. Barbour. 2022. Endocrine Emergencies During Pregnancy. *Obstetrics and Gynecology Clinics of North America* **49**:3, 473-489. [[Crossref](#)]
123. Zhuqing Li, Shiru Xu, Wei Luo, Jingmei Hu, Tao Zhang, Xue Jiao, Yingying Qin. 2022. Association between thyroid autoimmunity and the decline of ovarian reserve in euthyroid women. *Reproductive BioMedicine Online* **45**:3, 615-622. [[Crossref](#)]
124. Priyanka Jha, Liina Pöder, Phyllis Glanc, Krupa Patel-Lippmann, Melissa McGettigan, Mariam Moshiri, Stephanie Nougaret, Margarita V. Revzin, Marcia C. Javitt. 2022. Imaging Cancer in Pregnancy. *RadioGraphics* **42**:5, 1494-1513. [[Crossref](#)]
125. Dauren Adilbay, Avery Yuan, Paul B. Romesser, Richard J. Wong, Jatin P. Shah, Ashok R. Shaha, Michael R. Tuttle, Snehal Patel, Nancy Y. Lee, Ian Ganly. 2022. Well-Differentiated Thyroid Cancer: Who Should Get Postoperative Radiation?. *Annals of Surgical Oncology* **29**:9, 5582-5590. [[Crossref](#)]
126. Hiromi Murata, Susumu Tanaka, Hidetaka Okada. 2022. The Regulators of Human Endometrial Stromal Cell Decidualization. *Biomolecules* **12**:9, 1275. [[Crossref](#)]
127. Alma Linkeviciute, Rita Canario, Fedro Alessandro Peccatori, Kris Dierickx. 2022. Guidelines for Cancer Treatment during Pregnancy: Ethics-Related Content Evolution and Implications for Clinicians. *Cancers* **14**:17, 4325. [[Crossref](#)]

128. Hui Wang, Hai-Jun Wang, Mingyuan Jiao, Na Han, Jinhui Xu, Heling Bao, Zheng Liu, Yuelong Ji. 2022. Associations between Dynamic Vitamin D Level and Thyroid Function during Pregnancy. *Nutrients* 14:18, 3780. [[Crossref](#)]
129. Menglong Geng, Hui Gao, Baolin Wang, Kun Huang, Xiaoyan Wu, Chunmei Liang, Shuangqin Yan, Yan Han, Peng Ding, Wei Wang, Sheng Wang, Peng Zhu, Kaiyong Liu, Yunxia Cao, Fangbiao Tao. 2022. Urinary tetracycline antibiotics exposure during pregnancy and maternal thyroid hormone parameters: A repeated measures study. *Science of The Total Environment* 838, 156146. [[Crossref](#)]
130. Ranit Hizkiyahu, Ahmad Badeghiesh, Haitham Baghlaf, Michael H. Dahan. 2022. Associations between hyperthyroidism and adverse obstetric and neonatal outcomes: A study of a population database including almost 17,000 women with hyperthyroidism. *Clinical Endocrinology* 97:3, 347-354. [[Crossref](#)]
131. Meilan Liu, Dongyan Wang, Liqiong Zhu, Jianlan Yin, Xiaohui Ji, Yilei Zhong, Yuan Gao, Jianping Zhang, Yukun Liu, Rui Zhang, Hui Chen. 2022. Association of thyroid peroxidase antibodies with the rate of first-trimester miscarriage in euthyroid women with unexplained recurrent spontaneous abortion. *Frontiers in Endocrinology* 13. . [[Crossref](#)]
132. Sanja Medenica, Dzihan Abazovic, Aleksandar Ljubić, Jovana Vukovic, Aleksa Begovic, Gaspare Cucinella, Simona Zaami, Giuseppe Gullo. 2022. The Role of Cell and Gene Therapies in the Treatment of Infertility in Patients with Thyroid Autoimmunity. *International Journal of Endocrinology* 2022, 1-10. [[Crossref](#)]
133. Nanna Maria Uldall Torp, Niels Henrik Bruun, Peter Astrup Christensen, Aase Handberg, Stig Andersen, Stine Linding Andersen. 2022. Thyrotropin Receptor Antibodies in Early Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* 107:9, e3705-e3713. [[Crossref](#)]
134. Yuchao Zhang, Yanli Liu, Zhen Li, Yichun Guan. 2022. Comparison of reproductive outcomes in subclinical hypothyroidism women with high-normal versus low-normal thyroid-stimulating hormone levels after treatment with levothyroxine. *Clinical Endocrinology* 16. . [[Crossref](#)]
135. S.O. Shurpyak. 2022. The substantiation of differentiated therapy of the combined disgormonal pathology of reproductive organs in women of reproductive age with comorbid conditions. *Reproductive health of woman* 27:4, 30-37. [[Crossref](#)]
136. Jennifer S R Mammen, David S Cooper. 2022. TSH Receptor Antibody Testing in Early Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* 31. . [[Crossref](#)]
137. Stephen P. Fitzgerald, Nigel G. Bean, Samuel P. Fitzgerald, Henrik Falhammar. 2022. The application of new concepts of the assessment of the thyroid state to pregnant women. *Frontiers in Endocrinology* 13. . [[Crossref](#)]
138. Dana Larsen, Sargun Singh, Maria Brito. 2022. Thyroid, Diet, and Alternative Approaches. *The Journal of Clinical Endocrinology & Metabolism* 24. . [[Crossref](#)]
139. Xue-Feng Jiao, Miao Zhang, Jingjing Chen, Qiang Wei, Linan Zeng, Dan Liu, Chuan Zhang, Hailong Li, Kun Zou, Li Zhang, Lingli Zhang. 2022. The impact of levothyroxine therapy on the pregnancy, neonatal and childhood outcomes of subclinical hypothyroidism during pregnancy: An updated systematic review, meta-analysis and trial sequential analysis. *Frontiers in Endocrinology* 13. . [[Crossref](#)]
140. Mario Rotondi, Giulia Bendotti, Laura Croce, Martina Molteni, Andrea Carbone, Flavia Magri, Elizabeth N. Pearce, Luca Chiovato. 2022. A unique presentation of Graves' disease in a pregnant woman with severe hypothyroidism. *Gynecological Endocrinology* 38:8, 697-701. [[Crossref](#)]
141. Benjamin J. Gigliotti. 2022. Prevalence and Clinical Significance of Thyrotropin Receptor Antibodies in Early Pregnancy: Lessons from a Danish Cohort and Biobank Study. *Clinical Thyroidology* 34:8, 339-342. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
142. Xin Hou, Haixia Guan, Shuang Sun, Yang Shi, Chenyan Li, Aihua Liu, Yongze Li, Xiaotong Gao, Yuanyuan Hou, Yang Yang, Yushu Li, Zhongyan Shan, Weiping Teng. 2022. Outcomes of Early-Pregnancy Antithyroid Drug Withdrawal in Graves' Disease: A Preliminary Prospective Follow-Up Study. *Thyroid* 32:8, 983-989. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
143. Ying Yang, Tonglei Guo, Jinrong Fu, Jun Zhao, Yuanyuan Wang, Yuan He, Zuoqi Peng, Ya Zhang, Hongguang Zhang, Yue Zhang, Qiaomei Wang, Haiping Shen, Yiping Zhang, Donghai Yan, Xu Ma, Haixia Guan. 2022. Association of Preconception Thyrotropin Levels With Fecundability and Risk of Spontaneous Abortion in China. *JAMA Network Open* 5:8, e2228892. [[Crossref](#)]
144. C. C-H. Peng, E. N. Pearce. 2022. An update on thyroid disorders in the postpartum period. *Journal of Endocrinological Investigation* 45:8, 1497-1506. [[Crossref](#)]
145. Andrea Corsello, Cesare Morgante, Pietro Locantore. 2022. Una giovane gestante con morbo di Graves-Basedow complicato: un problema di non facile gestione. *L'Endocrinologo* 23:4, 409-411. [[Crossref](#)]
146. Philippe Caron. 2022. Management of thyrotoxicosis and pregnancy: Review of the current literature and an update of the care pathway. *Annales d'Endocrinologie* 83:4, 226-231. [[Crossref](#)]

147. Carlos Alberto Maganha, Rosiane Mattar, Cleo Otaviano Mesa Júnior, Suemi Marui, Sara Toassa Gomes Solha, Patrícia de Fátima dos Santos Teixeira, Alberto Carlos Moreno Zaconeta, Renato Teixeira Souza. 2022. Screening, diagnosis and management of hyperthyroidism in pregnancy. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics* 44:08, 806-818. [[Crossref](#)]
148. Zohar Steinberg Ben-Zeev, Marina Peniakov, Clari Felszer, Scott A Weiner, Avishay Lahad, Shlomo Almashanu, Yardena Tenenbaum Rakover. 2022. Usefulness of thyroid function assessment in infants born to mothers with thyroid dysfunction during pregnancy. *European Thyroid Journal* 11:4. . [[Crossref](#)]
149. Maja Hjelm Lundgaard, Allan Carlé, Ulla Birgitte Christiansen, Anne Sørensen, Søren Risom Kristensen, Stine Linding Andersen. 2022. Prolonged APTT and autoimmune overt hypothyroidism identified postpartum: a case report. *European Thyroid Journal* 11:4. . [[Crossref](#)]
150. Mihaela Țarnă, Luminița Nicoleta Cima, Anca Maria Panaitescu, Carmen Sorina Martin, Anca Elena Sîrbu, Carmen Gabriela Barbu, Bogdan Pavel, Andreea Nicoleta Șerbănică, Simona Fica. 2022. Preconception Counseling in Patients with Hypothyroidism and/or Thyroid Autoimmunity. *Medicina* 58:8, 1122. [[Crossref](#)]
151. Dorota Filipowicz, Ewelina Szczepanek-Parulska, Małgorzata Kłobus, Krzysztof Szymanowski, Thilo Samson Chillon, Sabrina Asaad, Qian Sun, Aniceta A. Mikulska-Sauermann, Marta Karaźniewicz-Lada, Franciszek K. Głowska, Dominika Wietrzyk, Lutz Schomburg, Marek Ruchała. 2022. Selenium Status and Supplementation Effects in Pregnancy—A Study on Mother–Child Pairs from a Single-Center Cohort. *Nutrients* 14:15, 3082. [[Crossref](#)]
152. Dingqian Sun, Shenjiao Lu, Xiaowen Gan, Gendie E. Lash. 2022. Is there a place for Traditional Chinese Medicine (TCM) in the treatment of recurrent pregnancy loss?. *Journal of Reproductive Immunology* 152, 103636. [[Crossref](#)]
153. Zheng Zhou, Lixiang Liu, Meihui Jin, Bingxuan Ren, Fangang Meng, Dandan Wang, Jianshuang Li, Baoxiang Li, Yanhong He, Fan Li, Hongmei Shen. 2022. Relationships between the serum TPOAb and TGAb antibody distributions and water iodine concentrations, thyroid hormones and thyroid diseases: a cross-sectional study of 2503 adults in China. *British Journal of Nutrition* 33, 1-11. [[Crossref](#)]
154. Kelsey Olerich, David Soper, Shani Delaney, Mary Sterrett. 2022. Pregnancy Care for Patients With Super Morbid Obesity. *Frontiers in Pediatrics* 10. . [[Crossref](#)]
155. Carola Deischinger, Michael Krebs, Alexandra Kautzky-Willer. 2022. Nahrungsergänzungsmittel und die Schilddrüse – ein Update zur Supplementierung von Mikronährstoffen. *Journal für Klinische Endokrinologie und Stoffwechsel* 22. . [[Crossref](#)]
156. Regan L Bailey, Tieraona Low Dog, Abbie E Smith-Ryan, Sai Krupa Das, Fiona C Baker, Zeynep Madak-Erdogan, Billy R Hammond, Howard D Sesso, Alex Eapen, Susan H Mitmesser, Andrea Wong, Haiyuen Nguyen. 2022. Sex Differences Across the Life Course: A Focus On Unique Nutritional and Health Considerations among Women. *The Journal of Nutrition* 152:7, 1597-1610. [[Crossref](#)]
157. Klara Gustin, Marie Vahter, Malin Barman, Bo Jacobsson, Helena Skräder, Helena Filipsson Nyström, Anna Sandin, Ann-Sofie Sandberg, Agnes E Wold, Maria Kippler. 2022. Assessment of Joint Impact of Iodine, Selenium, and Zinc Status on Women's Third-Trimester Plasma Thyroid Hormone Concentrations. *The Journal of Nutrition* 152:7, 1737-1746. [[Crossref](#)]
158. Umit Yasemin Sert, Hatice Kansu Celik, Burcu Kısa Karakaya, Ali Turhan Caglar, Gokcen Erdogan, Yaprak Engin-Ustun. 2022. The Impact of First Trimester TSH Level on Fetal Growth and Post-Term Pregnancy. *Fetal and Pediatric Pathology* 41:4, 584-591. [[Crossref](#)]
159. Ioannis Ilias, Charalampos Milionis, Eftychia Koukkou. 2022. Further understanding of thyroid function in pregnant women. *Expert Review of Endocrinology & Metabolism* 17:4, 365-374. [[Crossref](#)]
160. Sibel Demiral Sezer, Omercan Topaloglu. 2022. Effects of thyroid hormones in women with gestational diabetes. *Gynecological Endocrinology* 38:7, 588-591. [[Crossref](#)]
161. Liliana Mota Ferreira. 2022. THYROID HEALTH, PREGNANCY AND IODINE SUPPLEMENTATION. *Health and Society* 2:03, 132-177. [[Crossref](#)]
162. Gary L. Francis. 2022. Screening for Neonatal Hyperthyroidism. *Clinical Thyroidology* 34:7, 282-285. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
163. Nianyu Li, Yueshuang Lu, Pinxin Si, Zhuqing Li, Yingying Qin, Xue Jiao. 2022. The Impact of Moderately High Preconception Thyrotropin Levels on Ovarian Reserve Among Euthyroid Infertile Women Undergoing Assisted Reproductive Technology. *Thyroid* 32:7, 841-848. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
164. Alexandra L. Solomon, Alan P. Farwell, Elizabeth N. Pearce, Sun Y. Lee. 2022. Letter to the Editor: A Patient-Centered Survey-Based Assessment of Prenatal Management of Hypothyroidism for Women of Reproductive Age. *Thyroid* 32:7, 871-874. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]

165. Sofie Bliddal, Arash Derakhshan, Yi Xiao, Liang-Miao Chen, Tuija Männistö, Ghalia Ashoor, Fangbiao Tao, Suzanne J. Brown, Marina Vafeiadi, Sachiko Itoh, Elena Nikolaevna Grineva, Peter Taylor, Farkhanda Ghafoor, Bijay Vaidya, Andrew Hattersley, Lorena Mosso, Emily Oken, Reiko Kishi, Erik K. Alexander, Spyridoula Maraka, Kun Huang, Layal Chaker, Judit Bassols, Amna Pirzada, Abel López-Bermejo, Laura Boucai, Robin P. Peeters, Elizabeth N. Pearce, Scott McGill Nelson, Leda Chatzi, Tanja G. Vrijkotte, Polina V. Popova, John P. Walsh, Kypros H. Nicolaides, Eila Suvanto, Xuemian Lu, Victor J.M. Pop, Julie Lyng Forman, Tim I.M. Korevaar, Ulla Feldt-Rasmussen. 2022. Association of Thyroid Peroxidase Antibodies and Thyroglobulin Antibodies with Thyroid Function in Pregnancy: An Individual Participant Data Meta-Analysis. *Thyroid* **32**:7, 828-840. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
166. Neda Shakerian, Maysam Mard-Soltani, Sima Nasri, Mohammad Javad Rasaei, Saeed Khalili. 2022. Different combinations of monoclonal antibodies and polyclonal antibodies in the design of neonatal hypothyroidism diagnostic kit. *Applied Biochemistry and Biotechnology* **194**:7, 3167-3181. [[Crossref](#)]
167. Luis Rene Quiroa Barreno, Julia Bette Homem de Mello, Mateus Camargo Barros-Filho, Ana Lucia Francisco, Thiago Celestino Chulam, Clovis Antonio Lopes Pinto, Joao Gonçalves-Filho, Luiz Paulo Kowalski. 2022. Characterization of BRAF mutation in patients older than 45 years with well-differentiated thyroid carcinoma. *Brazilian Journal of Otorhinolaryngology* **88**:4, 523-528. [[Crossref](#)]
168. Inés Velasco, Lluís Vila, Maria Goya, Amelia Oleaga, Maria de la Calle, Fco. Javier Santamaria. 2022. Resumen ejecutivo del documento de consenso SEEN-SEGO sobre el manejo de la disfunción tiroidea durante el embarazo. *Endocrinología, Diabetes y Nutrición* **27**. . [[Crossref](#)]
169. Elizabeth N. Pearce. 2022. Management of Hypothyroidism and Hypothyroxinemia During Pregnancy. *Endocrine Practice* **28**:7, 711-718. [[Crossref](#)]
170. Luisa Rodriguez, Catherine Dinuer, Gary Francis. 2022. Treatment of hypothyroidism in infants, children and adolescents. *Trends in Endocrinology & Metabolism* **33**:7, 522-532. [[Crossref](#)]
171. Carolyn Ee, Kate Levett, Caroline Smith, Mike Armour, Hannah G. Dahlen, Prakshi Chopra, Paulette Maroun, Vibhuti S. Rao, Nicole Avard, Suzanne Grant, Hazel Keedle, Susanne Armour, Susan Arentz, Adele E. Cave, Kerry Sutcliffe, Kate Templeman. 2022. Complementary medicines and therapies in clinical guidelines on pregnancy care: A systematic review. *Women and Birth* **35**:4, e303-e317. [[Crossref](#)]
172. Gilad Karavani, Lina Daoud-Sabag, Cherut Chay, David Gillis, David Strich. 2022. Is TSH a Reliable Indicator of Thyroid Hormone Status in Pregnancy?. *Hormone and Metabolic Research* **54**:07, 435-441. [[Crossref](#)]
173. 2022. Effect of levothyroxine on pregnancy outcomes in euthyroid women with antithyroid antibodies. *Drug and Therapeutics Bulletin* **60**:7, 101-101. [[Crossref](#)]
174. Ying Guo, Bin Wei, Wei Dai, Hongjian Xie. 2022. Establishment of trimester-specific reference intervals for thyroid stimulating hormone and free thyroxine during pregnancy in southwest China by indirect method. *Annals of Clinical Biochemistry: International Journal of Laboratory Medicine* **59**:4, 234-241. [[Crossref](#)]
175. Enrique González-Madrid, Ma. Andreina Rangel-Ramírez, María José Mendoza-León, Oscar Álvarez-Mardones, Pablo A. González, Alexis M. Kalergis, Ma. Cecilia Opazo, Claudia A. Riedel. 2022. Risk Factors from Pregnancy to Adulthood in Multiple Sclerosis Outcome. *International Journal of Molecular Sciences* **23**:13, 7080. [[Crossref](#)]
176. Juan Jesús Fernández Alba, María Castillo Lara, José Manuel Jiménez Heras, Rocío Moreno Cortés, Carmen González Macías, Ángel Vilar Sánchez, Florentino Carral San Laureano, Luis Javier Moreno Corral. 2022. High First Trimester Levels of TSH as an Independent Risk Factor for Gestational Diabetes Mellitus: A Retrospective Cohort Study. *Journal of Clinical Medicine* **11**:13, 3776. [[Crossref](#)]
177. Zagorka Milovanović, Dejan Filimonović, Ivan Soldatović, Nataša Karadžov Orlić. 2022. Can Thyroid Screening in the First Trimester Improve the Prediction of Gestational Diabetes Mellitus?. *Journal of Clinical Medicine* **11**:13, 3916. [[Crossref](#)]
178. Qingru Wang, Yangqian Jiang, Hong Lv, Qun Lu, Shiyao Tao, Rui Qin, Lei Huang, Cong Liu, Xin Xu, Siyuan Lv, Mei Li, Zhi Li, Jiangbo Du, Yuan Lin, Hongxia Ma, Xia Chi, Zhibin Hu, Tao Jiang, Guoying Zhang. 2022. Association of Maternal Mild Hypothyroidism With Offspring Neurodevelopment in TPOAb-Negative Women: A Prospective Cohort Study. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
179. Renata Świątkowska-Stodulska, Agata Berlińska, Katarzyna Stefańska, Maciej Zieliński, Sebastian Kwiatkowski, Joanna Połom, Elżbieta Andrysiak-Mamos, Piotr Wydra, Krzysztof Sworczak. 2022. Endocrine Autoimmunity in Pregnancy. *Frontiers in Immunology* **13**. . [[Crossref](#)]
180. Rima K. Dhillon-Smith, Kristien Boelaert, Yadava B. Jeve, Abha Maheshwari, Arri Coomarasamy. 2022. Subclinical hypothyroidism and antithyroid autoantibodies in women with subfertility or recurrent pregnancy loss. *BJOG: An International Journal of Obstetrics & Gynaecology* **189**. . [[Crossref](#)]

181. Rui Yang, Danhua Pu, Rongrong Tan, Jie Wu. 2022. Association of methylenetetrahydrofolate reductase (MTHFR) gene polymorphisms (C677T and A1298C) with thyroid dysfunction: A meta-analysis and trial sequential analysis. *Archives of Endocrinology and Metabolism* **375**. . [[Crossref](#)]
182. Jose Mario Alves Junior, Wanderley Marques Bernardo, Laura Sterian Ward, Danilo Villagelin. 2022. Effect of Hyperthyroidism Control During Pregnancy on Maternal and Fetal Outcome: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
183. Hanze Du, Danning Wu, Xiang Zhou, Hongbo Yang, Huijuan Zhu, Shi Chen, Hui Pan. 2022. Preconception TSH and Adverse Pregnancy Outcomes in China: A Nationwide Prospective Cohort Study. *The Journal of Clinical Endocrinology & Metabolism* **107**:7, e2770-e2776. [[Crossref](#)]
184. Daniela Pereira Carvalho, Ariane Fontes Dias, Amanda Nancy Sferruzzi-Perri, Tania Maria Ortiga-Carvalho. 2022. Gaps in the knowledge of thyroid hormones and placental biology. *Biology of Reproduction* **106**:6, 1033-1048. [[Crossref](#)]
185. Ceren YILDIZ EREN, Hulusi Göktuğ GÜRER, Özlem ÖZGÜR GÜRSOY, Ozge GODEK. 2022. PREVALENCE OF PREMATURE OVARIAN FAILURE IN PATIENTS WITH AUTOIMMUNE THYROIDITIS. *Health Care Academician Journal* . [[Crossref](#)]
186. Jesse Bertinato, Jeremiah Gaudet, Nimal De Silva, Smitarani Mohanty, Cunye Qiao, Matthew Herod, Nathalie Gharibeh, Hope Weiler. 2022. Iodine Status of Mother-Infant Dyads from Montréal, Canada: Secondary Analyses of a Vitamin D Supplementation Trial in Breastfed Infants. *The Journal of Nutrition* **152**:6, 1459-1466. [[Crossref](#)]
187. Iván Del Campo Cano, Raquel Alarza Cano, Begoña Encinas Padilla, Nerea Lacámara Ornaechea, Ana Royuela Vicente, Miguel Ángel Marín Gabriel. 2022. A prospective study among neonates born to mothers with active or past Graves disease. *Gynecological Endocrinology* **38**:6, 495-498. [[Crossref](#)]
188. Edith J. M. Feskens, Regan Bailey, Zulfiqar Bhutta, Hans-Konrad Biesalski, Heather Eicher-Miller, Klaus Krämer, Wen-Harn Pan, James C. Griffiths. 2022. Women's health: optimal nutrition throughout the lifecycle. *European Journal of Nutrition* **61**:S1, 1-23. [[Crossref](#)]
189. Evert F.S. van Velsen, Angela M. Leung, Tim I.M. Korevaar. 2022. Diagnostic and Treatment Considerations for Thyroid Cancer in Women of Reproductive Age and the Perinatal Period. *Endocrinology and Metabolism Clinics of North America* **51**:2, 403-416. [[Crossref](#)]
190. Rima K. Dhillon-Smith, Kristien Boelaert. 2022. Preconception Counseling and Care for Pregnant Women with Thyroid Disease. *Endocrinology and Metabolism Clinics of North America* **51**:2, 417-436. [[Crossref](#)]
191. Xiaotong Gao, Aihua Liu, Xichang Wang, Yutong Han, Haoyu Wang, Jiashu Li, Yuanyuan Hou, Yang Yang, Huiru Wang, Chenyu Zhang, Weiping Teng, Zhongyan Shan. 2022. Postpartum Levothyroxine Adjustment and Its Impact Factors in Women With Hypothyroidism in Pregnancy. *Endocrine Practice* **28**:6, 578-585. [[Crossref](#)]
192. Rachita Nanda, Prasanta Kumar Nayak, Suprava Patel, Eli Mohapatra, Sarita Agrawal. 2022. First-Trimester Reference Intervals for Thyroid Function Testing among Women Screened at a Tertiary Care Hospital in India. *Journal of Laboratory Physicians* **14**:02, 183-189. [[Crossref](#)]
193. Tianqing Deng, Xiaoyang Liao, Shaomi Zhu. 2022. Recent Advances in Treatment of Recurrent Spontaneous Abortion. *Obstetrical & Gynecological Survey* **77**:6, 355-366. [[Crossref](#)]
194. Demet Sengul, Ilker Sengul, José Maria Soares Junior. 2022. Repercussion of thyroid dysfunctions in thyroidology on the reproductive system: Conditio sine qua non?. *Revista da Associação Médica Brasileira* **68**:6, 721-722. [[Crossref](#)]
195. Justus Baumgarten, Christian Happel, Daniel Groener, Jennifer Staudt, Benjamin Bockisch, Amir Sabet, Frank Grünwald, Thomas Rink. 2022. Retrospective Analysis of the Development of Human Thyroglobulin during Pregnancy in Patients with Treated Non-Recurrent Differentiated Thyroid Cancer. *Current Oncology* **29**:6, 4012-4019. [[Crossref](#)]
196. Mateusz Winder, Zofia Koszyła, Aleksandra Boral, Piotr Kocelak, Jerzy Chudek. 2022. The Impact of Iodine Concentration Disorders on Health and Cancer. *Nutrients* **14**:11, 2209. [[Crossref](#)]
197. Marianna Minnetti, Valentina Sada, Tiziana Feola, Elisa Giannetta, Carlotta Pozza, Daniele Gianfrilli, Andrea M. Isidori, Alessia Cozzolino. 2022. Selenium Supplementation in Pregnant Women with Autoimmune Thyroiditis: A Practical Approach. *Nutrients* **14**:11, 2234. [[Crossref](#)]
198. Ines Bucci, Cesidio Giuliani, Giulia Di Dalmazi, Gloria Formoso, Giorgio Napolitano. 2022. Thyroid Autoimmunity in Female Infertility and Assisted Reproductive Technology Outcome. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
199. Min Wu, Yuxi Yang, Yali Fan, Shan Guo, Tianhe Li, Muqing Gu, Tingting Zhang, Huimin Gao, Ruixia Liu, Chenghong Yin. 2022. Characteristics of the Intestinal Flora of TPOAb-Positive Women With Subclinical Hypothyroidism in the Second Trimester of Pregnancy: A Single-Center Prospective Cohort Study. *Frontiers in Cellular and Infection Microbiology* **12**. . [[Crossref](#)]

200. Subhash C Dash, Nalinikanta Sahoo, Udaybhanu Rout, Sujata P Mishra, Jayashree Swain, Arijit G Mazumder. 2022. Outcomes With Levothyroxine Treatment in Early Pregnancy With Subclinical Hypothyroidism. *Cureus* **18**. . [\[Crossref\]](#)
201. Maria Andersson, Christian P Braegger. 2022. The Role of Iodine for Thyroid Function in Lactating Women and Infants. *Endocrine Reviews* **43**:3, 469-506. [\[Crossref\]](#)
202. Elizabeth K Cahoon, Dale Preston, Rui Zhang, Vibha Vij, Mark P Little, Kiyohiko Mabuchi, Vladimir Drozdovitch, Konstantin Chizhov, Vasilina V Yauseyenko, Alexander V Rozhko, Ilya V Velalkin. 2022. Breast cancer risk in residents of Belarus exposed to Chernobyl fallout while pregnant or lactating: standardized incidence ratio analysis, 1997 to 2016. *International Journal of Epidemiology* **51**:2, 547-554. [\[Crossref\]](#)
203. Sufia Athar, Stephen F Beer, Zeena Martis, Mohammad I Alloub. 2022. The Prevalence of Thyroid Autoimmunity in Pregnancy and Adverse Neonatal Outcomes at a Secondary Care Hospital in the Middle East. *Cureus* **87**. . [\[Crossref\]](#)
204. Anna Capozzi, Giovanni Scambia, Stefano Lello. 2022. Subclinical hypothyroidism in women's health: from pre- to post-menopause. *Gynecological Endocrinology* **38**:5, 357-367. [\[Crossref\]](#)
205. Beata Pyrzak, Małgorzata Rumińska, Ewelina Witkowska-Sędek, Anna Kucharska. 2022. Follow-Up of Thyroid Function in Children With Neonatal Hyperthyroidism. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)
206. Elio Monsour, Naykky Singh Ospina. 2022. No Benefit of Levothyroxine in Euthyroid Thyroid Peroxidase Antibody-Positive Women with Recurrent Pregnancy Loss. *Clinical Thyroidology* **34**:5, 208-212. [\[Citation\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
207. Maren Goeckenjan, Thomas Strowitzki. 2022. Stoffwechsel der Mutter vor und bei Schwangerschaftseintritt. *Gynäkologische Endokrinologie* **20**:2, 101-110. [\[Crossref\]](#)
208. Myrthe M van Dijk, Rosa Vissenberg, Eric Fliers, Joris A M van der Post, Marie-Louise P van der Hoorn, Sabina de Weerd, Walter K Kuchenbecker, Annemieke Hoek, J Marko Sikkema, Harold R Verhoeve, Kimiko A Broeze, Corry H de Koning, Willem Verpoest, Ole B Christiansen, Carolien Koks, Jan P de Bruin, Dimitri N M Papatsonis, Helen Torrance, Madelon van Wely, Peter H Bisschop, Mariëtte Goddijn. 2022. Levothyroxine in euthyroid thyroid peroxidase antibody positive women with recurrent pregnancy loss (T4LIFE trial): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *The Lancet Diabetes & Endocrinology* **10**:5, 322-329. [\[Crossref\]](#)
209. Nicole E. Marshall, Barbara Abrams, Linda A. Barbour, Patrick Catalano, Parul Christian, Jacob E. Friedman, William W. Hay, Teri L. Hernandez, Nancy F. Krebs, Emily Oken, Jonathan Q. Purnell, James M. Roberts, Hora Soltani, Jacqueline Wallace, Kent L. Thornburg. 2022. The importance of nutrition in pregnancy and lactation: lifelong consequences. *American Journal of Obstetrics and Gynecology* **226**:5, 607-632. [\[Crossref\]](#)
210. Erika S. O. Patriota, Isis C. C. Lima, Eduardo A. F. Nilson, Sylvia C. C. Franceschini, Vivian S. S. Gonçalves, Nathalia Pizato. 2022. Prevalence of insufficient iodine intake in pregnancy worldwide: a systematic review and meta-analysis. *European Journal of Clinical Nutrition* **76**:5, 703-715. [\[Crossref\]](#)
211. Caitlin E. Mayhew, Kristin R. Simonson, Esther R. Ellsworth Bowers. 2022. Antepartum Care for Pregnant People with Overt Hypothyroidism, Subclinical Hypothyroidism, and Positive Thyroid Autoantibodies. *Journal of Midwifery & Women's Health* **67**:3, 295-304. [\[Crossref\]](#)
212. L. V. Adamyan, E. N. Andreeva, Yu. S. Absatarova, O. R. Grigoryan, I. I. Dedov, G. A. Melnichenko, L. V. Suturina, O. S. Filippov, E. V. Sheremetyeva, G. E. Chernukha, M. I. Yarmolinskaya. 2022. Clinical guidelines «Polycystic Ovary Syndrome». *Problems of Endocrinology* **68**:2, 112-127. [\[Crossref\]](#)
213. Yoon Young Cho, Bongseong Kim, Dong Wook Shin, Hye Ryoung Jang, Bo-Yeon Kim, Chan-Hee Jung, Jae Hyeon Kim, Sun Wook Kim, Jae Hoon Chung, Kyungdo Han, Tae Hyuk Kim. 2022. Graves' Disease and the Risk of End-Stage Renal Disease: A Korean Population-Based Study. *Endocrinology and Metabolism* **37**:2, 281-289. [\[Crossref\]](#)
214. Olha Protsiuk. 2022. Diagnosis and Treatment of Primary Hypothyroidism in Primary Care. *Family Medicine* **5**:1-2, 76-80. [\[Crossref\]](#)
215. Xiaoqin Wang, Jing Liu, Cheng Li, Yifan Xu, Xinyi Wang, Yu Lu, Ting Zhang, Hongzhi Cao, Linjuan Huang, Zhongfu Wang. 2022. Pregnancy-Related Diseases and Delivery Mode can Affect the Content of Human Milk Oligosaccharides: A Preliminary Study. *Journal of Agricultural and Food Chemistry* **70**:16, 5207-5217. [\[Crossref\]](#)
216. M. S. Medvedeva, A. S. Lyashenko, E. N. Lyashenko. 2022. Iodine deficiency as a cause of hypothyroidosis in pregnant women: diagnostics and preventive measures. *Meditsinskiy sovet = Medical Council* :5, 70-77. [\[Crossref\]](#)
217. Qianqian Zhang, Xiaoyan Xin, Li Wang. 2022. A Bibliometric Analysis of 8271 Publications on Thyroid Nodules From 2000 to 2021. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)
218. Guohua Li, Yang Liu, Xiujuan Su, Shijia Huang, Xiaosong Liu, Qiaoling Du. 2022. Effect of Levothyroxine on Pregnancy Outcomes in Pregnant Women With Hypothyroxinemia: An Interventional Study. *Frontiers in Endocrinology* **13**. . [\[Crossref\]](#)

219. Loris Marin, Guido Ambrosini, Marco Noventa, Flavia Filippi, Eugenio Ragazzi, Francesco Dessole, Giampiero Capobianco, Alessandra Andrisani. 2022. Thyroid Dysfunction after Gonadotropin-Releasing Hormone Agonist Administration in Women with Thyroid Autoimmunity. *International Journal of Endocrinology* **2022**, 1-7. [[Crossref](#)]
220. Yang Liu, Guohua Li, Nafei Guo, Xiaosong Liu, Shijia Huang, Qiaoling Du. 2022. Association Between Maternal Characteristics and the Risk of Isolated Maternal Hypothyroxinemia. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
221. Swetha Bhaskar, Chuchin Lim, Tahir Mahmood. Pre-Pregnancy Care 255-276. [[Crossref](#)]
222. Sabriye Korkut, Nilgün Çaylan, Ayse S. Özgü-Erdinç, Mustafa Ş. Akın, Meryem Ceyhan, Fatih Kara, Başak Tezel, Şerife S. Oğuz. 2022. Effect of Maternal Subclinical Hypothyroidism on Congenital Hypothyroidism Screening Results: A Retrospective Cohort Study. *American Journal of Perinatology* **31**. . [[Crossref](#)]
223. Fernando Candanedo-Gonzalez, Javier Rios-Valencia, Dafne Noemi Pacheco-Garcilazo, Wilfredo Valenzuela-Gonzalez, Armando Gamboa-Dominguez. Morphology Aspects of Hypothyroidism . [[Crossref](#)]
224. Wissal Abassi, Nejmeddine Ouerghi, Anissa Bouassida. Hypothyroidism Therapy . [[Crossref](#)]
225. Feng Xie, Jiang-Nan Wu. 2022. Maternal thyroxine and risk of fetal malformations. *Gynecological Endocrinology* **38**:4, 273-274. [[Crossref](#)]
226. Arwa Albashaireh, Spyridoula Maraka. 2022. Thyroid Function Classification in Early Pregnancy Varies Considerably Across Repeated Testing Within an Interval of a Few Weeks. *Clinical Thyroidology* **34**:4, 153-156. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
227. Jiajia Chen, Jing Zhu, Xianping Huang, Shenzhi Zhao, Huiqiu Xiang, Panpan Zhou, Tong Zhou, Zhangye Xu. 2022. Subclinical Hypothyroidism with Negative for Thyroid Peroxidase Antibodies in Pregnancy: Intellectual Development of Offspring. *Thyroid* **32**:4, 449-458. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
228. Alex Stagnaro-Green. 2022. Thyroid Disease in Pregnancy: A Touch of Clarity. *Thyroid* **32**:4, 347-348. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
229. Sima Nazarpour, Fahimeh Ramezani Tehrani, Maryam Rahmati, Mina Amiri, Fereidoun Azizi. 2022. Effects of isolated maternal hypothyroxinemia on adverse pregnancy outcomes. *Archives of Gynecology and Obstetrics* **305**:4, 903-911. [[Crossref](#)]
230. Anna Aulinas, Nicole Stanton-yonge, Apolonia Garcia-Patterson, Juan M. Adelantado, Carmen Medina, Juan José Espinós, Esther López, Susan M. Webb, Rosa Corcoy. 2022. Hypopituitarism and pregnancy: clinical characteristics, management and pregnancy outcome. *Pituitary* **25**:2, 275-284. [[Crossref](#)]
231. Francesco Vermiglio, Mariacarla Moleti. 2022. From preeclampsia to thyroid dysfunction: a long and winding road. *The Lancet Diabetes & Endocrinology* **10**:4, 232-233. [[Crossref](#)]
232. Chunping Hu, Susu Wang, Deqing Wu, Chonghuai Yan, Meiqin Wu. 2022. Subclinical hypothyroidism in pregnancy rats impaired offspring's spatial learning and memory and the cerebellar development. *Biochemical and Biophysical Research Communications* **602**, 63-69. [[Crossref](#)]
233. Lingyu Li, Lin Cheng, Ri Sa, Xian Qiu, Libo Chen. 2022. Real-world insights into the efficacy and safety of tyrosine kinase inhibitors against thyroid cancers. *Critical Reviews in Oncology/Hematology* **172**, 103624. [[Crossref](#)]
234. Sujuan Zhang, Yi Wu, Wenjing Pan, Guoyi Li, Di Zhang, Shuyi Li, Qi Huang, Zhao-min Liu. 2022. Free-Triiodothyronine to Free-Thyroxine Ratio Mediated the Effect of Prepregnancy Body Mass Index or Maternal Weight Gain During Early Pregnancy on Gestational Diabetes Mellitus. *Endocrine Practice* **28**:4, 398-404. [[Crossref](#)]
235. Yanni Guo, Xia Xu, Weijiao Xu, Tingting Liao, Jie Liang, Jianying Yan. 2022. Subsequent perinatal outcomes of pregnancy with two consecutive pregnancies with gestational diabetes mellitus: A population-based cohort study. *Journal of Diabetes* **14**:4, 282-290. [[Crossref](#)]
236. Xin He, Qin Yan, Chazhen Liu, Zhengyuan Wang, Ping Liao, Tong Liu, Zehuan Shi, Qi Song, Xueying Cui, Wenjing Wang, Jiajie Zang. 2022. Association of maternal thyroid dysfunction and autoimmunity with adverse birth outcomes. *Endocrine Connections* **11**:4. . [[Crossref](#)]
237. Ning Yuan, Jianbin Sun, Xin Zhao, Jing Du, Min Nan, Qiaoling Zhang, Xiaomei Zhang. 2022. Untreated thyroid autoantibody-negative SCH increases the risk of spontaneous abortions. *Endocrine Connections* **11**:4. . [[Crossref](#)]
238. Kjersti S Bakken, Kristina Randjelovic Nermo, Bjørn Gunnar Nedrebø, Tim I M Korevaar, Tor A Strand. 2022. Antenatal thyroid hormone therapy and antithyroid drug use in Norway from 2004 to 2018. *Endocrine Connections* **11**:4. . [[Crossref](#)]
239. Sarah Feigl, Barbara Obermayer-Pietsch, Philipp Klaritsch, Gudrun Pregartner, Sereina Annik Herzog, Elisabeth Lerchbaum, Christian Trummer, Stefan Pilz, Martina Kollmann. 2022. Impact of Thyroid Function on Pregnancy and Neonatal Outcome in Women with and without PCOS. *Biomedicines* **10**:4, 750. [[Crossref](#)]

240. Charalambos Fotakis, Giorgos Moros, Anna Kontogeorgou, Nicoletta Iacovidou, Theodora Boutsikou, Panagiotis Zoumpoulakis. 2022. Uncontrolled Thyroid during Pregnancy Alters the Circulative and Exerted Metabolome. *International Journal of Molecular Sciences* **23**:8, 4248. [[Crossref](#)]
241. Maddalena Casale, Marina Itala Baldini, Patrizia Del Monte, Antonia Gigante, Anna Grandone, Raffaella Origa, Maurizio Poggi, Franco Gadda, Rosalba Lai, Monia Marchetti, Gian Luca Forni. 2022. Good Clinical Practice of the Italian Society of Thalassemia and Haemoglobinopathies (SITE) for the Management of Endocrine Complications in Patients with Haemoglobinopathies. *Journal of Clinical Medicine* **11**:7, 1826. [[Crossref](#)]
242. Carla A. Lopes, Susana Prazeres, José Martinez-de-Oliveira, Edward Limbert, Manuel C. Lemos. 2022. Iodine Supplementation in Pregnancy in an Iodine-Deficient Region: A Cross-Sectional Survey. *Nutrients* **14**:7, 1393. [[Crossref](#)]
243. P. Morcel, S. Hadjadj, C. Ansquer, A. Yan Lun, B. Cariou, A.-S. Delemazure Chesneau, M. Le Bras, E. Langlois, D. Drui. 2022. Démarche diagnostique et prise en charge thérapeutique de l'hyperthyroïdie. *La Revue de Médecine Interne* **43**:4, 233-241. [[Crossref](#)]
244. Young Sil Eom, Jessica R. Wilson, Victor J. Bernet. 2022. Links between Thyroid Disorders and Glucose Homeostasis. *Diabetes & Metabolism Journal* **46**:2, 239-256. [[Crossref](#)]
245. Michela Bottani, Aasne K. Aarsand, Giuseppe Banfi, Massimo Locatelli, Abdurrahman Coşkun, Jorge Díaz-Garzón, Pilar Fernandez-Calle, Sverre Sandberg, Ferruccio Ceriotti, Anna Carobene. 2022. European Biological Variation Study (EuBIVAS): within- and between-subject biological variation estimates for serum thyroid biomarkers based on weekly samplings from 91 healthy participants. *Clinical Chemistry and Laboratory Medicine (CCLM)* **60**:4, 523-532. [[Crossref](#)]
246. Da Xu, Haolin Zhong. 2022. Correlation Between Hypothyroidism During Pregnancy and Glucose and Lipid Metabolism in Pregnant Women and Its Influence on Pregnancy Outcome and Fetal Growth and Development. *Frontiers in Surgery* **9**. . [[Crossref](#)]
247. Maïa Banigé, Dulanjalee Kariyawasam, Valerie Gauthereau, Dominique Luton, Michel Polak. 2022. Neonatal Screening for Hyperthyroidism Proof of Concept. *The Journal of Clinical Endocrinology & Metabolism* **107**:4, e1374-e1381. [[Crossref](#)]
248. Davut SAKIZ, Murat ÇALAPKULU, Muhammed Erkam SENCAR, Bekir UCAN, İlknur ÖZTÜRK ÜNSAL, Mustafa ÖZBEK, Erman ÇAKAL. 2022. fT3 index/TSH index ratio and free thyroid hormone index in the differential diagnosis of thyrotoxicosis. *Journal of Health Sciences and Medicine* **5**:2, 586-591. [[Crossref](#)]
249. Jing Dong, Ting Peng, Ming-Qing Li, Feng Xie, Jiang-Nan Wu. 2022. Association between Maternal Thyroxine and Risk of Fetal Congenital Heart Defects: A Hospital-Based Cohort Study. *International Journal of Endocrinology* **2022**, 1-8. [[Crossref](#)]
250. Yiqun Miao, Yang Xu, Ping Teng, Aihua Wang, Yuanyuan Zhang, Yun Zhou, Wenwen Liu. 2022. Efficacy of propylthiouracil in the treatment of pregnancy with hyperthyroidism and its effect on pregnancy outcomes: A meta-analysis. *PLOS ONE* **17**:3, e0265085. [[Crossref](#)]
251. Jacqueline Jonklaas. 2022. Optimal Thyroid Hormone Replacement. *Endocrine Reviews* **43**:2, 366-404. [[Crossref](#)]
252. Claudia Ashkar, Shoshana Sztal-Mazer, Duncan J. Topliss. 2022. How to manage Graves' disease in women of childbearing potential. *Clinical Endocrinology* **1**. . [[Crossref](#)]
253. Zhao-min Liu, Guoyi Li, Yi Wu, Di Zhang, Sujuan Zhang, Yuan-Tao Hao, Weiqing Chen, Qi Huang, Shuyi Li, Yaojie Xie, Mingtong Ye, Chun He, Ping Chen, Wenjing Pan. 2022. Increased Central and Peripheral Thyroid Resistance Indices During the First Half of Gestation Were Associated With Lowered Risk of Gestational Diabetes—Analyses Based on Huizhou Birth Cohort in South China. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
254. Yue Han, Jun Wang, Xiaoying Wang, Ling Ouyang, Yan Li. 2022. Relationship Between Subclinical Hypothyroidism in Pregnancy and Hypertensive Disorder of Pregnancy: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
255. Farahnaz Mir, Hossein Chiti, Saeideh Mazloomzadeh. 2022. Short-Term Adverse Pregnancy Outcomes in Women with Subclinical Hypothyroidism: A Comparative Approach of Iranian and American Guidelines. *Journal of Thyroid Research* **2022**, 1-6. [[Crossref](#)]
256. Yaniv S. Ovadia, Shmuel Zangen, Shani R. Rosen, Dov Gefel, Shlomo Almashanu, Carlos Benbassat, Shlomo Fytlovich, Dorit Aharoni, Eyal Y. Anteby, Simon Shenhav. 2022. Maternal iodine deficiency: a newborns' overweight risk factor? A prospective study. *Archives of Gynecology and Obstetrics* **305**:3, 777-787. [[Crossref](#)]
257. Wei Zheng, Shu Heng Yang, Chen Yang, Bing Nan Ren, Si Min Sun, Yan Li Liu, Ru Jing Yang, Hua Lou, Lin Lin Zhang, Yi Chun Guan. 2022. Perinatal outcomes of singleton live births after preimplantation genetic testing during single frozen-thawed blastocyst transfer cycles: a propensity score-matched study. *Fertility and Sterility* **117**:3, 562-570. [[Crossref](#)]
258. Peng Zuo, Jingjing Yang. 2022. Pregnancy outcomes in papillary thyroid cancer survivors#A retrospective cohort study. *Gynecology and Obstetrics Clinical Medicine* **2**:1, 20-22. [[Crossref](#)]

259. Zheng Yang, Haoyu Wang, Yifu Liu, Yan Feng, Yang Xiang, Jing Li, Zhongyan Shan, Weiping Teng. 2022. The expression of anti-protein disulfide isomerase A3 autoantibody is associated with the increased risk of miscarriage in euthyroid women with thyroid autoimmunity. *International Immunopharmacology* **104**, 108507. [[Crossref](#)]
260. Sun Y. Lee, Elizabeth N. Pearce. 2022. Assessment and treatment of thyroid disorders in pregnancy and the postpartum period. *Nature Reviews Endocrinology* **18**:3, 158-171. [[Crossref](#)]
261. Shruthi Ravindra, Sahana Shetty. 2022. Neonatal thyrotoxicosis with maternal hypothyroidism. *BMJ Case Reports* **15**:3, e247865. [[Crossref](#)]
262. Martin Draznin, Preeti Borgohain, Shibani Kanungo. 2022. Newborn Screening in Pediatric Endocrine Disorders. *Endocrines* **3**:1, 107-114. [[Crossref](#)]
263. Iwona Magdalena Gawron, Rafał Baran, Kamil Derbisz, Robert Jach. 2022. Association of Subclinical Hypothyroidism with Present and Absent Anti-Thyroid Antibodies with PCOS Phenotypes and Metabolic Profile. *Journal of Clinical Medicine* **11**:6, 1547. [[Crossref](#)]
264. Karina dos Santos, Eliane Lopes Rosado, Ana Carolina Proença da Fonseca, Gabriella Pinto Belfort, Leticia Barbosa Gabriel da Silva, Marcelo Ribeiro-Alves, Verônica Marques Zembruski, J. Alfredo Martínez, Cláudia Saunders. 2022. FTO and ADRB2 Genetic Polymorphisms Are Risk Factors for Earlier Excessive Gestational Weight Gain in Pregnant Women with Pregestational Diabetes Mellitus: Results of a Randomized Nutrigenetic Trial. *Nutrients* **14**:5, 1050. [[Crossref](#)]
265. Anastasia Trouva, Michael Alvarsson, Jan Calissendorff, Bjørn Olav Åsvold, Eszter Vanky, Angelica Lindén Hirschberg. 2022. Thyroid Status During Pregnancy in Women With Polycystic Ovary Syndrome and the Effect of Metformin. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
266. Ran S Rotem, Gabriel Chodick, Michael Davidovitch, Andrea Bellavia, Marc G Weisskopf. 2022. Maternal Thyroid Anomalies and Attention-Deficit Hyperactivity Disorder in Progeny. *American Journal of Epidemiology* **191**:3, 430-440. [[Crossref](#)]
267. Mary Kathryn Bohn, Khosrow Adeli. 2022. Physiological and metabolic adaptations in pregnancy: importance of trimester-specific reference intervals to investigate maternal health and complications. *Critical Reviews in Clinical Laboratory Sciences* **59**:2, 76-92. [[Crossref](#)]
268. Juliane Léger, Clemence Delcour, Jean-Claude Carel. 2022. Fetal and Neonatal Thyroid Dysfunction. *The Journal of Clinical Endocrinology & Metabolism* **107**:3, 836-846. [[Crossref](#)]
269. Raffaella Di Girolamo, Marco Liberati, Claudia Silvi, Francesco D'Antonio. 2022. Levothyroxine Supplementation in Euthyroid Pregnant Women With Positive Autoantibodies: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
270. Qi Cao, Huili Zhu, Jiani Zhang, Yujing Li, Wei Huang. 2022. Pregnancy Outcomes in Thyroid Cancer Survivors: A Propensity Score-Matched Cohort Study. *Frontiers in Endocrinology* **13**. . [[Crossref](#)]
271. Ma. Cecilia Opazo, Irenice Coronado-Arrázola, Omar P. Vallejos, Rodrigo Moreno-Reyes, Carlos Fardella, Lorena Mosso, Alexis M. Kalergis, Susan M. Bueno, Claudia A. Riedel. 2022. The impact of the micronutrient iodine in health and diseases. *Critical Reviews in Food Science and Nutrition* **62**:6, 1466-1479. [[Crossref](#)]
272. Jan Krátký. 2022. Levothyroxine. *Vnitřní lékařství* **68**:1, 68-73. [[Crossref](#)]
273. Ping Li, Shuo Lin, Jinhui Cui, Xinjuan Chen, Zhaoran Meng, Jianhui Fan. 2022. Impact of Early Pregnancy Subclinical Hypothyroidism on Gestational Diabetes Mellitus: A Retrospective Study of 7,536 Cases. *Journal of Women's Health* **31**:2, 293-298. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
274. Hans Hauner. 2022. Stoffwechsel und Ernährung in der Schwangerschaft. *Monatsschrift Kinderheilkunde* **170**:2, 116-124. [[Crossref](#)]
275. Alan Chun Hong Lee, George J. Kahaly. 2022. Pathophysiology of thyroid-associated orbitopathy. *Best Practice & Research Clinical Endocrinology & Metabolism* **185**, 101620. [[Crossref](#)]
276. Aglaia Kyriilli, David Unuane, Kris G. Poppe. 2022. Thyroid autoimmunity and pregnancy in euthyroid women. *Best Practice & Research Clinical Endocrinology & Metabolism* **27**, 101632. [[Crossref](#)]
277. Jennifer Ngounda, Jeannine Baumgartner, Mariette Nel, Corinna May Walsh. 2022. Iodine status of pregnant women residing in the urban Free State Province of South Africa is borderline adequate: The NuEMI study. *Nutrition Research* **98**, 18-26. [[Crossref](#)]
278. Katharina Binz, Felix Huber, Corinne Chmiel, Uwe Beise. 2022. Schilddrüsenerkrankungen. *Praxis* **111**:2, 102-108. [[Crossref](#)]
279. Meiqin Wu, Yaqian Wang, Chonghuai Yan, Yan Zhao. 2022. Study on subclinical hypothyroidism in pregnancy: a bibliometric analysis via CiteSpace. *The Journal of Maternal-Fetal & Neonatal Medicine* **35**:3, 556-567. [[Crossref](#)]

280. Rachel A. Bradbury, Darshika Christie-David, Howard C. Smith, Karen Byth, Creswell J. Eastman. 2022. Prior iodine exposure and impact on thyroid function during controlled ovarian hyperstimulation: A prospective study. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **62**:1, 133-139. [[Crossref](#)]
281. Balsam Bohlega, Afshan Zahedi, George Tomlinson, Denise S. Feig. 2022. Weight-based thyroid dosing vs fixed dosing during pregnancy for subclinical hypothyroidism: A retrospective cohort study. *Clinical Endocrinology* **96**:2, 263-269. [[Crossref](#)]
282. Nykola Louise Kent, Sharat Chandra Atluri, James Sebastian Martin Cuffe. 2022. Maternal Hypothyroidism in Rats Reduces Placental Lactogen, Lowers Insulin Levels, and Causes Glucose Intolerance. *Endocrinology* **163**:2. . [[Crossref](#)]
283. Ning Yuan, Li Wang, Zhi Li, Xiaomei Zhang. 2022. Thyroid Diseases During Pregnancy: Bibliometric Analysis of Scientific Publications. *Endocrine, Metabolic & Immune Disorders - Drug Targets* **22**:2, 247-258. [[Crossref](#)]
284. Po-Keng Cheng, Hsin-Chang Chen, Pao-Lin Kuo, Jung-Wei Chang, Wan-Ting Chang, Po-Chin Huang. 2022. Associations between Oxidative/Nitrosative Stress and Thyroid Hormones in Pregnant Women—Tainan Birth Cohort Study (TBSCS). *Antioxidants* **11**:2, 334. [[Crossref](#)]
285. Eva Gerbier, Sereina M. Graber, Marlene Rauch, Carole A. Marxer, Christoph R. Meier, David Baud, Ursula Winterfeld, Eva Blozik, Daniel Surbek, Julia Spoendlin, Alice Panchaud. 2022. Use of Prescribed Drugs to Treat Chronic Diseases during Pregnancy in Outpatient Care in Switzerland between 2014 and 2018: Descriptive Analysis of Swiss Health Care Claims Data. *International Journal of Environmental Research and Public Health* **19**:3, 1456. [[Crossref](#)]
286. Qian Zhu, Kahindo P. Muyayalo, Qian-Han Xu, Jing Wang, Huan Wang, Ai-Hua Liao. 2022. *Prunella vulgaris* can improve the pregnancy outcomes of experimental autoimmune thyroiditis rats by inhibiting Th1/Th17 immune responses. *Journal of Reproductive Immunology* **149**, 103469. [[Crossref](#)]
287. Yanjun Cai, Yajuan Xu, Yanjie Ban, Jingjing Li, Zongzong Sun, Miao Zhang, Biao Wang, Xiaofeng Hou, Yingqi Hao, Qian Ouyang, Bo Wu, Mengqi Wang, Wentao Wang. 2022. Plasma Lipid Profile and Intestinal Microflora in Pregnancy Women With Hypothyroidism and Their Correlation With Pregnancy Outcomes. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
288. Ara L. Unanyan, Laura G. Pivazyan, Araksya A. Zakaryan, Archil A. Siordiya, Anatoliy I. Ishchenko. 2022. Treatment of habitual pregnancy miscarriage of various origins. *V.F.Negirev Archives of Obstetrics and Gynecology* **9**:1, 13-21. [[Crossref](#)]
289. Simten GENÇ, Başak CINGILLIOĞLU. 2022. Effects of subclinical hypothyroidism on maternal and obstetric outcomes during pregnancy. *The European Research Journal* . [[Crossref](#)]
290. Puneet Khanna, Soumya Sarkar. 2022. Parturient with Endocrine Disorders in the Intensive Care Unit. *Indian Journal of Critical Care Medicine* **25**:S3, S255-S260. [[Crossref](#)]
291. Carlos Alfonso Builes-Barrera. 2022. Función tiroidea y embarazo. *Medicina y Laboratorio* **26**:1, 11-13. [[Crossref](#)]
292. Myriam Vanessa Rueda-Galvis, Carlos Alfonso Builes-Barrera. 2022. Fisiología de la tiroides e hipotiroidismo en el embarazo. Revisión de tema. *Medicina y Laboratorio* **26**:1, 15-33. [[Crossref](#)]
293. Yu Meng, Jing Lin, Jianxia Fan. 2022. A Novel Nomogram for Predicting the Risk of Premature Delivery Based on the Thyroid Function in Pregnant Women. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
294. Sylwia Płaczkowska, Małgorzata Terpińska, Agnieszka Piwowar. 2022. Establishing laboratory-specific reference intervals for TSH and fT4 by use of the indirect Hoffman method. *PLOS ONE* **17**:1, e0261715. [[Crossref](#)]
295. Jingjing Li, Yajuan Xu, Yanjun Cai, Miao Zhang, Zongzong Sun, Yanjie Ban, Shanshan Zhai, Yingqi Hao, Qian Ouyang, Bo Wu, Mengqi Wang, Wentao Wang. 2022. Association of Differential Metabolites With Small Intestinal Microflora and Maternal Outcomes in Subclinical Hypothyroidism During Pregnancy. *Frontiers in Cellular and Infection Microbiology* **11**. . [[Crossref](#)]
296. Yuko Hasegawa, Yoshikazu Kitahara, Satoko Osuka, Yumiko Tsukui, Mio Kobayashi, Akira Iwase. 2022. Effect of hypothyroidism and thyroid autoimmunity on the ovarian reserve: A systematic review and meta-analysis. *Reproductive Medicine and Biology* **21**:1. . [[Crossref](#)]
297. Peter Kopp. Hypothyroidism 409-414. [[Crossref](#)]
298. Elizabeth N. Pearce. Introduction to Endocrine Disorders in Pregnancy 303-305. [[Crossref](#)]
299. Sun Y. Lee, Elizabeth N. Pearce. Diagnosis and Management of Thyroid Cancer in Pregnant Women 341-348. [[Crossref](#)]
300. Jad G. Sfeir, Hossein Gharib. Hypothyroidism 15-23. [[Crossref](#)]
301. Alessandro Delitala, Margherita Maioli, Francesco Dessole, Marco Petrillo, Giampiero Capobianco. Thyroid Function in Pregnancy 1-16. [[Crossref](#)]
302. Atieh Amouzegar. Thyroid Storm and Neonatal Hyperthyroidism 167-174. [[Crossref](#)]
303. Fahimeh Ramezani Tehrani. Autoimmune Thyroid Disease in Pregnancy 175-190. [[Crossref](#)]
304. Trevor E. Angell. Thyroid Nodules and Thyroid Cancer in the Pregnant Woman 191-209. [[Crossref](#)]

305. Caroline T. Nguyen. Postpartum Thyroiditis: Diagnosis and Management 211-222. [[Crossref](#)]
306. Fereidoun Azizi. Postpartum Thyrotoxicosis 223-231. [[Crossref](#)]
307. Jennifer B. Plotkin, Angela M. Leung. Iodine Requirements in Pregnancy 31-43. [[Crossref](#)]
308. Fereidoun Azizi. Screening of Thyroid Function in Pregnancy 45-55. [[Crossref](#)]
309. Theodora Pappa, Erik K. Alexander. Preconception Counseling 57-68. [[Crossref](#)]
310. Inés Velasco, Onyebuchi E. Okosieme. Maternal Clinical Hypothyroidism 69-91. [[Crossref](#)]
311. Creswell J. Eastman, Norman J. Blumenthal. Gestational Subclinical Hypothyroidism 93-108. [[Crossref](#)]
312. Eunice López-Muñoz. Isolated Hypothyroxinemia During Pregnancy 109-134. [[Crossref](#)]
313. K. Gkastaris, G. E. Krassas. Maternal Hyperthyroidism 135-154. [[Crossref](#)]
314. Lisa B. Nachtigall. Optimization of Care for Women with Complex Pituitary Tumors Who Seek Fertility 149-161. [[Crossref](#)]
315. Alyssa Dominguez, Rachel Danis, John D. Carmichael. Ovulation, Pregnancy, and Delivery in the Female Patient with Hypopituitarism 165-176. [[Crossref](#)]
316. Atılay Yaylacı, Murat Öztürk, Tania Sih. The Management of Neck Mass During Pregnancy and the Postpartum Period 687-698. [[Crossref](#)]
317. Başat Fethallah, Nuray Bayar Muluk, Felicia Manole. Thyroid Nodules and Cancer During Pregnancy and the Postpartum Period 723-744. [[Crossref](#)]
318. Ömer Bayir, Latif Akan, Hakan Korkmaz. General Principles of Ent Surgery in Pregnancy 963-973. [[Crossref](#)]
319. Peter Kopp. Hypothyroidism 1-6. [[Crossref](#)]
320. Erica M. Schindewolf, Julie S. Moldenhauer. Reproduction, prenatal screening, and diagnosis in 22q11.2 deletion syndrome 370-386. [[Crossref](#)]
321. Shannon M. Clark, Luis A. Monsivais. Antenatal thyroid disease and pharmacotherapy in pregnancy 339-355. [[Crossref](#)]
322. Samantha Newman, Terry F. Davies. Pharmacology of the Thyroid 155-172. [[Crossref](#)]
323. Renato Tozzoli, Nicola Bizzaro. TSH receptor autoantibodies in Graves' disease 69-82. [[Crossref](#)]
324. Katrina B. Mitchell, Margaret M. Fleming, Philip O. Anderson, Jamie G. Giesbrandt. ABM Clinical Protocol #31: Radiology and Nuclear Medicine Studies in Lactating Women 1017-1022. [[Crossref](#)]
325. Sara E. Lubitz. Hypopituitarism 239-258. [[Crossref](#)]
326. Caroline T. Nguyen, Jorge H. Mestman. Graves' Hyperthyroidism in Pregnancy 285-298. [[Crossref](#)]
327. Jamie M. Pitlick. Thyroid Ablation 244-247. [[Crossref](#)]
328. Kwan Cheng, Lubaina S. Presswala. Thyroid Uptake and Scan 248-253. [[Crossref](#)]
329. Yiqiu Wei, Songchen Cai, Jinli Ding, Yong Zeng, Tailang Yin, Lianghui Diao. Metabolic control of immune responses in women with recurrent pregnancy loss and recurrent implantation failure 379-394. [[Crossref](#)]
330. Elena Borodina, Alexander M. Gzgzryan, Lyailya Kh. Dzhemlikhanova, Dariko A. Niauri, Yehuda Shoenfeld. Antithyroid antibodies and reproductive function 153-164. [[Crossref](#)]
331. Véronique Raverot, Michèle d'Herbomez. Évaluation biologique de la fonction thyroïdienne 20-28. [[Crossref](#)]
332. Philippe Caron. Thyroïde et grossesse Prise en charge diagnostique et thérapeutique de l'hypothyroïdie et l'hyperthyroïdie au cours de la grossesse 231-246. [[Crossref](#)]
333. Ioannis Tsakiridis, Sonia Giouleka, Anargyros Kourtis, Apostolos Mamopoulos, Apostolos Athanasiadis, Themistoklis Dagklis. 2022. Thyroid Disease in Pregnancy: A Descriptive Review of Guidelines. *Obstetrical & Gynecological Survey* 77:1, 45-62. [[Crossref](#)]
334. Faustine Tanguy, Safouane Hamdi, Karim Chikh, Daniel Glinioer, Philippe Caron. 2022. Central hypothyroidism during pregnancy in a woman with Graves' disease. *Clinical Endocrinology* 96:1, 89-91. [[Crossref](#)]
335. Sicelosethu S. Siro, Lizelle Zandberg, Jennifer Ngounda, Amy Wise, Elizabeth A. Symington, Linda Malan, Cornelius M. Smuts, Jeannine Baumgartner. 2022. Iodine status of pregnant women living in urban Johannesburg, South Africa. *Maternal & Child Nutrition* 18:1. . [[Crossref](#)]
336. María Fernanda Toro-Wills, Angélica Imitola-Madero, Angelina Alvarez-Londoño, Abraham Hernández-Blanquissett, María Cristina Martínez-Ávila. 2022. Thyroid cancer in women of reproductive age: Key issues for the clinical team. *Women's Health* 18, 174550572211363. [[Crossref](#)]

337. Mizuho Minakata, Mitsuru Ito, Taketoshi Kishi, Mikiko Hada, Yuzuki Masaki, Tomohiko Nakamura, Kazuyoshi Kousaka, Toshihiko Kasahara, Takumi Kudo, Eijun Nishihara, Shuji Fukata, Mitsushige Nishikawa, Takashi Akamizu, Akira Miyauchi. 2022. The magnitude of increased Levothyroxine dose during pregnancy in patients on thyroid-stimulating hormone (TSH) suppression treatment after total thyroidectomy for papillary carcinoma. *Endocrine Journal* **69**:2, 165-172. [[Crossref](#)]
338. Natsuko Watanabe, Jaeduk Yoshimura Noh, Takashi Akamizu, Masanobu Yamada. 2022. Survey of the actual administration of thiamazole for hyperthyroidism in Japan by the Japan Thyroid Association. *Endocrine Journal* **69**:1, 35-43. [[Crossref](#)]
339. Koshi Hashimoto. 2022. Update on subclinical thyroid dysfunction. *Endocrine Journal* **69**:7, 725-738. [[Crossref](#)]
340. Wen-Yu Liu, Yu-Ting Zhu, Xian-Ming Xu, Jia-Rong Zhang, Xu-Pei Gan, De-Cui Cheng, Feng Lu, Cong Lu. 2022. Associations of Thyroid Function Tests with Lipid Levels and Adverse Pregnancy Outcomes During the First Trimester. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy* **Volume 15**, 973-981. [[Crossref](#)]
341. Jiayi Luo, Jialing Yuan. 2022. Effects of Levothyroxine Therapy on Pregnancy and Neonatal Outcomes in Subclinical Hypothyroidism. *International Journal of General Medicine* **Volume 15**, 6811-6820. [[Crossref](#)]
342. 2022. 15. Management of Diabetes in Pregnancy: Standards of Medical Care in Diabetes—2022. *Diabetes Care* **45**:Supplement_1, S232-S243. [[Crossref](#)]
343. Sarah Mathai. 2022. Diagnosis of hypothyroidism in pregnancy and screening of babies born to hypothyroid mothers. *Journal of Pediatric Endocrinology and Diabetes* **2**, 16-17. [[Crossref](#)]
344. JubbinJagan Jacob. 2022. TSH receptor antibodies (TRAb) assay: An underutilized test in India. *Indian Journal of Endocrinology and Metabolism* **26**:1, 1. [[Crossref](#)]
345. Sarika Palepu, ArvindKumar Singh, GoutamKumar Saharia, Suravi Patra, Sweta Singh, Manish Taywade, Vikas Bhatia. 2022. Hypothyroidism in pregnancy: An alarming concern in a rural community of Eastern India. *Indian Journal of Community Medicine*, ahead of print0. [[Crossref](#)]
346. Ayla Coussa, ThomasM Barber, Zakwan Khrait, Samer Cheaib, HayderA Hasan. 2022. Relationship between maternal serum thyroid-stimulating hormone and in vitro fertilisation-conceived pregnancy outcomes. *Journal of Human Reproductive Sciences*, ahead of print0. [[Crossref](#)]
347. SantoshKumar Singh, Sanchita Chakrabarti, AK Naik, BarunKumar Chakrabarty, N Nagaraja, Vimal Upreti. 2022. Universal thyroid screening in maternal health care in India: The need of the hour. *Journal of Marine Medical Society*, ahead of print0. [[Crossref](#)]
348. Uroš Pecikoza. 2022. Functional thyroid disorders in the pharmacy setting: How can we help our patients?. *Arhiv za farmaciju* **72**:5, 503-520. [[Crossref](#)]
349. Mirjana Stojković. 2022. Thyroid function disorders. *Arhiv za farmaciju* **72**:5, 429-443. [[Crossref](#)]
350. Sanja Medenica, Miloš Žarković. 2022. Thyroid autoimmunity and reproduction: Bidirectional relationship that continues to intrigue. *Medicinski glasnik Specijalne bolnice za bolesti štitaste žlezde i bolesti metabolizma* **27**:87, 7-27. [[Crossref](#)]
351. Željka Aleksić, Aleksandar Aleksić, Branka Đorđević. 2022. Subclinical hypothyroidism. *Timocki medicinski glasnik* **47**:1, 11-22. [[Crossref](#)]
352. Yozen Fuse, Yoshiya Ito, Yoshimasa Shishiba, Minoru Irie. 2022. Gestational trimester-specific reference ranges for serum thyrotropin and free thyroxine in Japanese. *Endocrine Journal* **69**:12, 1447-1455. [[Crossref](#)]
353. Fernanda Campos da Silva, Gustavo Mourão Rodrigues. Standards In Prenatal Care 157-169. [[Crossref](#)]
354. Henri Augusto Korkes, Renato José Bauer, Nelson Sass. Chronic Hypertension in Pregnancy 393-403. [[Crossref](#)]
355. Luciana C. Cima, Mariane T. Tauile, Viviane P. Monteiro, Isabela Bussade. Thyroid Disorders 469-482. [[Crossref](#)]
356. Lenita Zajdenverg, Carlos Antonio Negrato. Pregestational Diabetes Mellitus 405-425. [[Crossref](#)]
357. Ujjawal Bairagi, Brahmeshwar Mishra. An update on the development of advanced drug delivery systems for the treatment of hyperthyroidism 269-279. [[Crossref](#)]
358. Yuhan Zhou, Fen Chen, Lingyu Wang, Chunhui Tian, Shuo Zhang, Feifei Ding, Jie Deng. 2021. Establishment of an iodine model for prevention of iodine-excess-induced thyroid dysfunction in pregnant women. *Open Life Sciences* **16**:1, 1357-1364. [[Crossref](#)]
359. Hatice ÖZİŞİK, Şevki ÇETİNKALP, Aslı SUNER, İsmet HORTU, Ahmet Mete ERGENOĞLU, Mehmet ERDOĞAN. 2021. Comparison of Thyroid Function Tests in Pregnant Women with and without Gestational Diabetes. *Turkish Journal of Diabetes and Obesity* **5**:3, 285-290. [[Crossref](#)]
360. Jelte van der Vaart, Lynn Bosmans, Stijn F. Sijbesma, Kévin Knoop, Willine J. van de Wetering, Henny G. Otten, Harry Begthel, Inne H. M. Borel Rinkes, Jeroen Korving, Eef G. W. M. Lentjes, Carmen Lopez-Iglesias, Peter J. Peters, Hanneke M. van Santen,

- Menno R. Vriens, Hans Clevers. 2021. Adult mouse and human organoids derived from thyroid follicular cells and modeling of Graves' hyperthyroidism. *Proceedings of the National Academy of Sciences* **118**:51. . [\[Crossref\]](#)
361. Shaobo Chen, Yinzhen Pi, Haiyan Gong, Huaijun Wang, Shu Liu. 2021. Single-Photon Emission Computed Tomography Image-Assisted Diagnosis of Thyroid Diseases under Convolutional Network Neural Algorithm. *Scientific Programming* **2021**, 1-9. [\[Crossref\]](#)
362. Zheng Ding, Yindi Liu, Spyridoula Maraka, Nadia Abdelouahab, He-Feng Huang, William D. Fraser, Jianxia Fan. 2021. Pregnancy and Neonatal Outcomes With Levothyroxine Treatment in Women With Subclinical Hypothyroidism Based on New Diagnostic Criteria: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology* **12**. . [\[Crossref\]](#)
363. Guan-ying Nie, Rui Wang, Peng Liu, Ming Li, Dian-jun Sun. 2021. Mild Anemia May Affect Thyroid Function in Pregnant Chinese Women During the First Trimester. *Frontiers in Endocrinology* **12**. . [\[Crossref\]](#)
364. Tereza Planck, Mikael Lantz, Petros Perros, Enrico Papini, Roberto Attanasio, Endre V. Nagy, Laszlo Hegedüs. 2021. Use of Thyroid Hormones in Hypothyroid and Euthyroid Patients: A 2020 THESIS Questionnaire Survey of Members of the Swedish Endocrine Society. *Frontiers in Endocrinology* **12**. . [\[Crossref\]](#)
365. Jesse Bertinato, Cunye Qiao, Mary R L'Abbé. 2021. Iodine Status of Canadian Children, Adolescents, and Women of Childbearing Age. *The Journal of Nutrition* **151**:12, 3710-3717. [\[Crossref\]](#)
366. Louise Brough. 2021. Iodine Intake for Pregnant and Breastfeeding Women and Their Infants Remains a Global Concern. *The Journal of Nutrition* **151**:12, 3604-3605. [\[Crossref\]](#)
367. Anastasios Malakasis, Francoise H. Harlow. Thyroid Disease in Pregnancy 304-310. [\[Crossref\]](#)
368. Arwa Albashaireh, Spyridoula Maraka. 2021. Iron Deficiency Is Associated with Maternal Hypothyroxinemia in the Third Trimester of Pregnancy. *Clinical Thyroidology* **33**:12, 526-528. [\[Citation\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
369. Rodrigo Moreno-Reyes, Bernard Corvilain, Caroline Daelemans, Fleur Wolff, Camilo Fuentes Peña, Stefanie Vandevijvere. 2021. Iron Deficiency Is a Risk Factor for Thyroid Dysfunction During Pregnancy: A Population-Based Study in Belgium. *Thyroid* **31**:12, 1868-1877. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
370. Vedrana Höggqvist Tabor, Mikael Höggqvist Tabor, Sarai Keestra, Jean-Etienne Parrot, Alexandra Alvergne. 2021. Improving the Quality of Life of Patients with an Underactive Thyroid Through mHealth: A Patient-Centered Approach. *Women's Health Reports* **2**:1, 182-194. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#) [\[Supplementary Material\]](#)
371. Wenxia Tian, Wenqing Yan, Yang Liu, Fangfang Zhou, Haixia Wang, Wenguang Sun. 2021. The Status and Knowledge of Iodine among Pregnant Women in Shanghai. *Biological Trace Element Research* **199**:12, 4489-4497. [\[Crossref\]](#)
372. Bharti Goel, Poonam Goel, Jasbinder Kaur. 2021. Reference Levels for TSH in Iodine-Sufficient Low-Risk Pregnant Women. *The Journal of Obstetrics and Gynecology of India* **71**:6, 596-599. [\[Crossref\]](#)
373. Giovanna Savastano, Valentina Del Negro, Damiana Pompeo, Sara Sorrenti, Paola Galoppi, Roberto Brunelli, Maria Grazia Piccioni. 2021. Thyroid Disease, Pregnancy, and Selenium Supplementation. *Current Obstetrics and Gynecology Reports* **10**:4, 115-123. [\[Crossref\]](#)
374. Florence Scheffler, Rosalie Cabry, Marion Soye, Henri Copin, Moncef Ben Khalifa, Aviva Devaux, Rachel Desailoud. 2021. Growth hormone replacement improved oocyte quality in a patient with hypopituitarism: A study of follicular fluid. *Annales d'Endocrinologie* **82**:6, 590-596. [\[Crossref\]](#)
375. Navpreet Kaur, Raj Suryanarayanan. 2021. Levothyroxine Sodium Pentahydrate Tablets – Formulation Considerations. *Journal of Pharmaceutical Sciences* **110**:12, 3743-3756. [\[Crossref\]](#)
376. Sho Fukui, Yukihiko Ikeda, Yuko Kataoka, Haruyuki Yanaoka, Hiromichi Tamaki, Tokutarou Tsuda, Mitsumasa Kishimoto, Hiroshi Noto, Sachiko Ohde, Masato Okada. 2021. Clinical significance of monitoring hypothyroidism in patients with autoimmune rheumatic disease: a retrospective cohort study. *Scientific Reports* **11**:1. . [\[Crossref\]](#)
377. Jingjing Li, Yajuan Xu, Zongzong Sun, Yanjun Cai, Biao Wang, Miao Zhang, Yanjie Ban, Xiaofeng Hou, Yingqi Hao, Qian Ouyang, Bo Wu, Mengqi Wang, Wentao Wang. 2021. Differential lipids in pregnant women with subclinical hypothyroidism and their correlation to the pregnancy outcomes. *Scientific Reports* **11**:1. . [\[Crossref\]](#)
378. Ying Jin, Jane Coad, Shao J. Zhou, Sheila Skeaff, Thiagarajah Ramilan, Louise Brough. 2021. Prevalence of thyroid dysfunction in postpartum women with suboptimal iodine and selenium and adequate iron status. *Clinical Endocrinology* **95**:6, 873-881. [\[Crossref\]](#)
379. Jinju Dong, Shouyan Liu, Lingyun Wang, Xingjian Zhou, Qinghong Zhou, Congli Liu, Jingrui Zhu, Weilan Yuan, Wang-yang Xu, Jie Deng. 2021. Iodine monitoring models contribute to avoid adverse birth outcomes related more than adequate iodine intake. *BMC Pregnancy and Childbirth* **21**:1. . [\[Crossref\]](#)
380. Fereidoun Azizi, Hengameh Abdi, Atieh Amouzegar. 2021. Control of Graves' hyperthyroidism with very long-term methimazole treatment: a clinical trial. *BMC Endocrine Disorders* **21**:1. . [\[Crossref\]](#)

381. Magnus Bein, Oriana Hoi Yun Yu, Sonia Marzia Grandi, Francesca Y. E. Frati, Ihab Kandil, Kristian B. Filion. 2021. Levothyroxine and the risk of adverse pregnancy outcomes in women with subclinical hypothyroidism: a systematic review and meta-analysis. *BMC Endocrine Disorders* 21:1. . [[Crossref](#)]
382. Lijue Ren, Cuiying Wei, Feng Wei, Ruiting Ma, Yan Liu, Yonghong Zhang, Wei Wang, Jing Du, Lin Bai, Yexia Xue, Shaohua Cui. 2021. A case report of rhabdomyolysis and osteofascial compartment syndrome in a patient with hypothyroidism and diabetes. *BMC Endocrine Disorders* 21:1. . [[Crossref](#)]
383. Lale Susan Karakis, Huseyin Kiyak, Berfin Okmen, Cagdas Ozdemir, Engin Turkgeldi. 2021. Impact of preconceptional serum thyroid stimulating hormone values ranging between 2.5 and 4.5 mIU/L on live birth rates following ovulation induction and intrauterine insemination treatment for unexplained infertility. *BMC Women's Health* 21:1. . [[Crossref](#)]
384. Yuchao Zhang, Wenbin Wu, Yanli Liu, Xingling Wang, Yichun Guan, Liting Jia. 2021. Analysis of basal serum TSH, FT3, and FT4 levels based on age, sampling time in women with infertility. *BMC Women's Health* 21:1. . [[Crossref](#)]
385. Yixuan Lin, Diqun Chen, Jiani Wu, Zhihui Chen. 2021. Iodine status five years after the adjustment of universal salt iodization: a cross-sectional study in Fujian Province, China. *Nutrition Journal* 20:1. . [[Crossref](#)]
386. Stine Linding Andersen, Kasper Krogh Nielsen, Søren Risom Kristensen. 2021. The interrelationship between pregnancy, venous thromboembolism, and thyroid disease: a hypothesis-generating review. *Thyroid Research* 14:1. . [[Crossref](#)]
387. Silmar Paulo Moreira Rates, Flávio Diniz Capanema, Bárbara do Amaral, Cristina Marques Secundino, Leticia Miguez de Souza Michelli, Rafaela Catelan Martins Pereira, Fábio da Veiga Ued, Carlos Alberto Nogueira-de-Almeida. 2021. Nutritional status of iodine in pregnant and non-pregnant adolescents assisted at the Family Health Strategy in Vespasiano, MG. *Revista Brasileira de Saúde Materno Infantil* 21:4, 1035-1043. [[Crossref](#)]
388. Francesca Gorini, Laura Sabatino, Alessandro Pingitore, Cristina Vassalle. 2021. Selenium: An Element of Life Essential for Thyroid Function. *Molecules* 26:23, 7084. [[Crossref](#)]
389. Verónica Melero, Isabelle Runkle, Nuria Garcia de la Torre, Paz De Miguel, Johanna Valerio, Laura del Valle, Ana Barabash, Concepción Sanabria, Inmaculada Moraga, Cristina Familiar, Alejandra Durán, Maria Jose Torrejón, Jose Angel Diaz, Martin Cuesta, Jorge Grabiell Ruiz, Inés Jiménez, Mario Pazos, Miguel Angel Herraiz, Nuria Izquierdo, Noelia Pérez, Pilar Matia, Natalia Perez-Ferre, Clara Marcuello, Miguel Angel Rubio, Alfonso Luis Calle-Pascual. 2021. The Consumption of Food-Based Iodine in the Immediate Pre-Pregnancy Period in Madrid Is Insufficient. San Carlos and Pregnancy Cohort Study. *Nutrients* 13:12, 4458. [[Crossref](#)]
390. Ratna Acharya, Kiran Upadhyay. 2021. Unmasking of Gitelman Syndrome during Pregnancy in an Adolescent with Thyrotoxic Crisis. *Pediatric Reports* 13:4, 632-638. [[Crossref](#)]
391. Anca Maria Panaitescu. Pregnancy and Graves' Disease . [[Crossref](#)]
392. Ji Eun Jeong, So Hee Lee, Young Hyun Kim, Yoon Young Jang, Jin-Kyung Kim. 2021. A Case of Neonatal Thyrotoxicosis with Cardiac Insufficiency. *Neonatal Medicine* 28:4, 161-166. [[Crossref](#)]
393. Xiaotong Gao, Xichang Wang, Yutong Han, Haoyu Wang, Jiashu Li, Yuanyuan Hou, Yang Yang, Huiru Wang, Weiping Teng, Zhongyan Shan. 2021. Postpartum Thyroid Dysfunction in Women With Known and Newly Diagnosed Hypothyroidism in Early Pregnancy. *Frontiers in Endocrinology* 12. . [[Crossref](#)]
394. Daniela Vejrazkova, Josef Vcelak, Eliska Vaclavikova, Marketa Vankova, Katerina Zajickova, Jana Vrbikova, Michaela Duskova, Petra Pacesova, Zdenek Novak, Bela Bendlova. 2021. Recurrence of Graves' Disease: What Genetics of HLA and PTPN22 Can Tell Us. *Frontiers in Endocrinology* 12. . [[Crossref](#)]
395. Xiujuan Su, Yang Liu, Guohua Li, Xiaosong Liu, Shijia Huang, Tao Duan, Qiaoling Du. 2021. Associations of Hypothyroxinemia With Risk of Preeclampsia-Eclampsia and Gestational Hypertension. *Frontiers in Endocrinology* 12. . [[Crossref](#)]
396. Jan Tuzil, Jana Bartakova, Torquil Watt, Tomas Dolezal. 2021. Health-related quality of life in women with autoimmune thyroid disease during pregnancy and postpartum: systematic review including 321,850 pregnancies. *Expert Review of Pharmacoeconomics & Outcomes Research* 21:6, 1179-1193. [[Crossref](#)]
397. Stephen P. Fitzgerald, Nigel G. Bean, James V. Hennessey, Henrik Falhammar. 2021. Thyroid testing paradigm switch from thyrotropin to thyroid hormones—Future directions and opportunities in clinical medicine and research. *Endocrine* 74:2, 285-289. [[Crossref](#)]
398. Jean-Philippe Rousseau, Luana Tenorio-Lopes, Sergio Cortez Ghio, Pascale Desjardins, Stéphanie Fournier, Richard Kinkead. 2021. Thyroid hormones during the perinatal period are necessary to respiratory network development of newborn rats. *Experimental Neurology* 345, 113813. [[Crossref](#)]
399. Anne Ottney, Lydia Lebeau. 2021. A “secret shopper” survey of community pharmacist prenatal vitamin recommendations. *Journal of the American Pharmacists Association* 61:6, e60-e64. [[Crossref](#)]

400. Rekha Ramot, Garima Kachhawa, Vidushi Kulshrestha, Vishnubhatla Sreenivas, Devasenathipathy Kandasamy, Rajesh Khadgawat. 2021. Bone mass in newborns and its predictors. *Clinical Endocrinology* **95**:5, 727-734. [[Crossref](#)]
401. Małgorzata Karbownik-Lewińska, Jan Stępnia, Andrzej Lewiński. 2021. Potential Risk Factors for Isolated Hypothyroxinemia in Women of Childbearing Age—Results from Retrospective Analysis. *Journal of Clinical Medicine* **10**:22, 5384. [[Crossref](#)]
402. Vija Veisa, Ieva Kalere, Tatjana Zake, Ieva Strele, Marina Makrecka-Kuka, Sabine Upmale-Engela, Andrejs Skesters, Dace Rezeberga, Aivars Lejnicks, Iveta Pudule, Daiga Grinberga, Biruta Velika, Maija Dambrova, Ilze Konrade. 2021. Assessment of Iodine and Selenium Nutritional Status in Women of Reproductive Age in Latvia. *Medicina* **57**:11, 1211. [[Crossref](#)]
403. Aniceta A. Mikulska, Dorota Filipowicz, Franciszek K. Główska, Ewelina Szczepanek-Parulska, Marek Ruchała, Michał Bartecki, Marta Karaźniewicz-Łada. 2021. HPLC Analysis of the Urinary Iodine Concentration in Pregnant Women. *Molecules* **26**:22, 6797. [[Crossref](#)]
404. A. I. Denysenko, V. I. Cherniy. 2021. PERIOPERATIVE METABOLISM IN PATIENTS WITH THYROTOXICOSIS, POSSIBILITIES OF ITS CORRECTION. *Клінічна та профілактична медицина* **3**:17, 36-48. [[Crossref](#)]
405. Zheng Ding, Fei Guo, Yulai Zhou, Xiaoyi Huang, Zhiwei Liu, Jianxia Fan. 2021. Thyroxine Supplementation in Pregnant Women After Thyroidectomy for Thyroid Cancer and Neonatal Birth Weight. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
406. Yuantong Sun, Weiwei Zheng, Ling Zhang, Huijuan Zhao, Xun Li, Chao Zhang, Wuren Ma, Dajun Tian, Kun-Hsing Yu, Shuo Xiao, Liping Jin, Jing Hua. 2021. Quantifying the Impacts of Pre- and Post-Conception TSH Levels on Birth Outcomes: An Examination of Different Machine Learning Models. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
407. Saadi JS AlJadir. 2021. Thyroid dysfunction in pregnancy: practical perspective. *Obstetrics & Gynecology International Journal* **12**:5, 316-321. [[Crossref](#)]
408. Isabelle Runkle, María Paz de Miguel, Ana Barabash, Martin Cuesta, Ángel Diaz, Alejandra Duran, Cristina Familiar, Nuria García de la Torre, Miguel Ángel Herraiz, Nuria Izquierdo, Ángel Diaz, Clara Marcuello, Pilar Matia, Verónica Melero, Carmen Montañez, Inmaculada Moraga, Natalia Perez-Ferre, Noelia Perez, Carla Assaf-Balut, Miguel Ángel Rubio, Jorge Gabriel Ruiz-Sanchez, Concepción Sanabria, María José Torrejon, Johanna Valerio, Laura del Valle, Alfonso Calle-Pascual. 2021. Early Levothyroxine Treatment for Subclinical Hypothyroidism or Hypothyroxinemia in Pregnancy: The St Carlos Gestational and Thyroid Protocol. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
409. N. A. Kurmacheva, M. Yu. Svinarev, Yu. V. Chernenkov, O. S. Panina, K. V. Yakovleva, S. V. Anisimova. 2021. Iodine deficiency in pregnant women and newborns in the Saratov region: current state of the problem. *Meditinskiy sovet = Medical Council* :13, 156-161. [[Crossref](#)]
410. Maria C. Opazo, Juan Carlos Rivera, Pablo A. Gonzalez, Susan M. Bueno, Alexis M. Kalergis, Claudia A. Riedel. 2021. Thyroid Gene Mutations in Pregnant and Breastfeeding Women Diagnosed With Transient Congenital Hypothyroidism: Implications for the Offspring's Health. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
411. Soo Myoung Shin, Ka Hee Yi. 2021. Antithyroid drug therapy for Graves' disease. *Journal of the Korean Medical Association* **64**:10, 690-698. [[Crossref](#)]
412. Ya-Hui Yu, Kristian B. Filion, Pauline Reynier, Robert W. Platt, Oriana H. Y. Yu, Sonia M. Grandi. 2021. Use of levothyroxine among pregnant women with subclinical hypothyroidism in the United Kingdom: A population-based assessment. *Pharmacology Research & Perspectives* **9**:5. . [[Crossref](#)]
413. Nykola L. Kent, Sophia L. Young, Lisa K. Akison, James S. M. Cuffe. 2021. Is the link between elevated TSH and gestational diabetes mellitus dependant on diagnostic criteria and thyroid antibody status: a systematic review and meta-analysis. *Endocrine* **74**:1, 38-49. [[Crossref](#)]
414. Martin Grußendorf. 2021. Hyperthyreose: Ursachen, Diagnostik, Behandlung. *MMW - Fortschritte der Medizin* **163**:17, 50-59. [[Crossref](#)]
415. Rongrong Li, Wei Chen, Yanping Liu, Liangkun Ma, Ling Qiu, Jianhua Han, Rui Li. 2021. The Impact of Preconceptional Hysterosalpingography with Oil-based Contrast on Maternal and Neonatal Iodine Status. *Reproductive Sciences* **28**:10, 2887-2894. [[Crossref](#)]
416. Yongze Li, Zhongyan Shan, Weiping Teng. 2021. The Iodine Status and Prevalence of Thyroid Disorders Among Women of Childbearing Age in China: National Cross-sectional Study. *Endocrine Practice* **27**:10, 1028-1033. [[Crossref](#)]
417. Marion Lecorguillé, Juliane Léger, Anne Forhan, Marie Cheminat, Marie-Noëlle Dufourg, Barbara Heude, Marie-Aline Charles. 2021. Pregnancy outcomes in women with preexisting thyroid diseases: a French cohort study. *Journal of Developmental Origins of Health and Disease* **12**:5, 704-713. [[Crossref](#)]
418. Stephanie Allelein, Matthias Schott. 2021. Morbus Basedow. *DMW - Deutsche Medizinische Wochenschrift* **146**:20, 1337-1343. [[Crossref](#)]

419. Maja Avramovska, Neda Milevska Kostova, Borislav Karanfilski, Sandra Hunziker, Olivija Vaskova, Goran Dimitrov, Elena Dzikova, Ana Daneva Markova, Marija Hadzi Lega, Gligor Tofoski, Aleksandar Sikole. 2021. Thyroid Function of Pregnant Women and Perinatal Outcomes in North Macedonia. *Revista Brasileira de Ginecologia e Obstetricia / RBGO Gynecology and Obstetrics* **43**:10, 736-742. [[Crossref](#)]
420. Synnøve Næss, Maria W Markhus, Tor A Strand, Marian Kjellefold, Lisbeth Dahl, Ann-Elin M Stokland, Bjørn G Nedrebø, Inger Aakre. 2021. Iodine Nutrition and Iodine Supplement Initiation in Association with Thyroid Function in Mildly-to-Moderately Iodine-Deficient Pregnant and Postpartum Women. *The Journal of Nutrition* **151**:10, 3187-3196. [[Crossref](#)]
421. Daniel R. Morales, Lionel Fonkwen, Hedvig M. E. Nordeng. 2021. Antithyroid drug use during pregnancy and the risk of birth defects in offspring: systematic review and meta-analysis of observational studies with methodological considerations. *British Journal of Clinical Pharmacology* **87**:10, 3890-3900. [[Crossref](#)]
422. Nikolina Zdraveska, Mirjana Kocova. 2021. Thyroid function and dysfunction in preterm infants—Challenges in evaluation, diagnosis and therapy. *Clinical Endocrinology* **95**:4, 556-570. [[Crossref](#)]
423. Karin Säljö, Anders Thornell, Chunsheng Jin, Olov Norlén, Susann Teneberg. 2021. Characterization of Human Medullary Thyroid Carcinoma Glycosphingolipids Identifies Potential Cancer Markers. *International Journal of Molecular Sciences* **22**:19, 10463. [[Crossref](#)]
424. L. A. Suplotova, O. B. Makarova, E. N. Maksyukova, L. N. Smolina. 2021. Analysis of the thyroid status of pregnant women in the region of mild iodine deficiency. *Meditinskiy sovet = Medical Council* :12, 276-284. [[Crossref](#)]
425. Masoume Nodehi, Abolghasem Ajami, Maryam Izad, Hossein Asgarian Omran, Fatemeh Esfahanian, Saeed Yekaninejad, Mahbobeh Hemmatbadi, Atieh Amouzegar, Reza Chahardoli, Fatemeh Mansouri, Ali Akbar Saboor-Yaraghi. 2021. The Frequency of CD4+ T Cells in Women with Hashimoto's Thyroiditis. *International Journal of Endocrinology and Metabolism* **19**:4. . [[Crossref](#)]
426. Maya Berlin, Dana Barchel, Anna Brik, Elkana Kohn, Ayelet Livne, Rimona Keidar, Josef Tovbin, Moshe Betser, Miki Moskovich, Dror Mandel, Ronit Lubetzky, Amit Ovental, Pam Factor-Litvak, Malka Britzi, Tomer Ziv-Baran, Ronit Koren, Chagit Klieger, Matitiah Berkovitch, Ilan Matok, Ronella Marom. 2021. Maternal and Newborn Thyroid Hormone, and the Association With Polychlorinated Biphenyls (PCBs) Burden: The EHF (Environmental Health Fund) Birth Cohort. *Frontiers in Pediatrics* **9**. . [[Crossref](#)]
427. Yue Yu, Qian-Qi Liu, De-Yun Liu, Dan-Dan Wang, Li-Qi Yang, Shu-Ming Ye. 2021. Antibodies against thyroid-stimulating hormone receptor cause maternal-neonatal transmission of Graves' Disease. *Experimental and Therapeutic Medicine* **22**:5. . [[Crossref](#)]
428. Jinrong Fu, Guofeng Zhang, Pei Xu, Rui Guo, Jiarong Li, Haixia Guan, Yushu Li. 2021. Seasonal Changes of Thyroid Function Parameters in Women of Reproductive Age Between 2012 and 2018: A Retrospective, Observational, Single-Center Study. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
429. Sofia Manousou, Robert Eggertsen, Lena Hulthén, Helena Filipsson Nyström. 2021. A randomized, double-blind study of iodine supplementation during pregnancy in Sweden: pilot evaluation of maternal iodine status and thyroid function. *European Journal of Nutrition* **60**:6, 3411-3422. [[Crossref](#)]
430. Danpei Li, Sitao Hu, Xiaoyu Meng, Xuefeng Yu. 2021. Changes in thyroid function during controlled ovarian hyperstimulation (COH) and its impact on assisted reproduction technology (ART) outcomes: a systematic review and meta-analysis. *Journal of Assisted Reproduction and Genetics* **38**:9, 2227-2235. [[Crossref](#)]
431. Irfan Ahmed, Victor Ma, Yuanchao Liu, Muhammad Shehzad Khan, Zhenhui Liu, Chi Zhang, Santosh Kumar Paidi, Francis A. M. Manno, Noreen Amjad, Sinai H. C. Manno, Rafay Ahmed, Alan W. L. Law, Ahmed Ali, Faizan Raza, Yanpeng Zhang, William C. S Cho, Ishan Barman, Martin Alda, Veerle Bergink, Condon Lau. 2021. Lithium from breast-milk inhibits thyroid iodine uptake and hormone production, which are remedied by maternal iodine supplementation. *Bipolar Disorders* **23**:6, 615-625. [[Crossref](#)]
432. Marianne Skovsager Andersen, Tina Kold Jensen, Anja Fenger Dreyer, Jeppe Buur Madsen, Henrik Thybo Christesen, Ivan Brandslund, Niels Bilenberg, Dorte Glintborg. 2021. Free thyroxine in early pregnancy is an independent negative predictor of 3rd trimester HbA1c. Odense child cohort. *Clinical Endocrinology* **95**:3, 508-519. [[Crossref](#)]
433. Erin Farah, Mary K. Barger, Carrie Klima, Beverly Rossman, Patricia Hershberger. 2021. Impaired Lactation: Review of Delayed Lactogenesis and Insufficient Lactation. *Journal of Midwifery & Women's Health* **66**:5, 631-640. [[Crossref](#)]
434. Weiqi Liu, Qiaoling Liu, Weiling Liu, Cuiqing Qiu. 2021. Maternal risk factors and pregnancy complications associated with low birth weight neonates in preterm birth. *Journal of Obstetrics and Gynaecology Research* **47**:9, 3196-3202. [[Crossref](#)]

435. Jesmine Banu, Mumtahena Amir, Shakeela Ishrat, Sadia Afrin Munmun, Nighat Sultana. 2021. Successful Outcome Following Hysteroscopic Septoplasty in a Woman with Seven Consecutive Pregnancy Losses. *Fertility & Reproduction* **03**:03, 84-87. [[Crossref](#)]
436. Onyebuchi E Okosieme, Medha Agrawal, Danyal Usman, Carol Evans. 2021. Method-dependent variation in TSH and FT4 reference intervals in pregnancy: A systematic review. *Annals of Clinical Biochemistry: International Journal of Laboratory Medicine* **58**:5, 537-546. [[Crossref](#)]
437. Pietro LOCANTORE, Andrea CORSELLO, Caterina POLICOLA, Alfredo PONTECORVI. 2021. Subclinical thyroid diseases and isolated hypothyroxinemia during pregnancy. *Minerva Endocrinology* **46**:3. . [[Crossref](#)]
438. Alfredo PONTECORVI, Silvia GELLI. 2021. Subclinical endocrine diseases: a never-ending challenge. *Minerva Endocrinology* **46**:3. . [[Crossref](#)]
439. Sahar Mazloomi, Narges Alizadeh, Gholamreza Shafiee, Somayeh Ghorbani, Massoume Afsharirad. 2021. The Effects of Hypothyroidism on Maternal Health. *Journal of Advances in Medical and Biomedical Research* **29**:136, 279-285. [[Crossref](#)]
440. Sonia Santander Ballestín, Marta Isabel Giménez Campos, Jara Ballestín Ballestín, María José Luesma Bartolomé. 2021. Is Supplementation with Micronutrients Still Necessary during Pregnancy? A Review. *Nutrients* **13**:9, 3134. [[Crossref](#)]
441. Felicity Coad, Charlotte Frise. 2021. Tachycardia in pregnancy: when to worry?. *Clinical Medicine* **21**:5, e434-e437. [[Crossref](#)]
442. Arsenio Spinillo, Irene De Maggio, Beatrice Ruspini, Camilla Bellingeri, Chiara Cavagnoli, Serena Giannico, Anna Boschetti, Flavia Magri, Elisabetta Lovati, Fausta Beneventi. 2021. Placental pathologic features in thyroid autoimmunity. *Placenta* **112**, 66-72. [[Crossref](#)]
443. Matthew Lumchee, Mimi Yue, Josephine Laurie, Adam Morton. 2021. Therapeutic plasma exchange for Graves' disease in pregnancy. *Obstetric Medicine* **84**, 1753495X2110313. [[Crossref](#)]
444. Min Song, Mengyuan Lei, Chenghan Luo, Zanyang Shi, Xinru Cheng, Wenqian Ding, Wenjun Cao, Jingdi Zhang, Jian Ge, Mengmeng Wang, Peige Xia, Fengxia Mao, Li Wang, Qian Zhang. 2021. Development of a Nomogram for Moderate-to-Severe Bronchopulmonary Dysplasia or Death: Role of N-Terminal Pro-brain Natriuretic Peptide as a Biomarker. *Frontiers in Pediatrics* **9**. . [[Crossref](#)]
445. Vera M. Kodentsova, Dmitry V. Risnik, Stanislav V. Pavlovich, Olga B. Ladodo. 2021. Optimization of the micronutrients sufficiency of feeding women and children on exclusively breastfeeding by enriching of the woman diet. *Gynecology* **23**:3, 222-227. [[Crossref](#)]
446. Kurnia Alisaputri, Sony Wibisono. 2021. A CASE REPORT: CHALLENGES IN THE MANAGEMENT OF TOXIC MULTI NODULAR THYROID IN PREGNANCY WHO HAD CRISIS THYROID. *Current Internal Medicine Research and Practice Surabaya Journal* **2**:2, 56. [[Crossref](#)]
447. Kagan Gungor, Nur Dokuzeylul Gungor. 2021. Antithyroid antibodies may predict serum beta HCG levels and biochemical pregnancy losses in euthyroid women with IVF single embryo transfer. *Gynecological Endocrinology* **37**:8, 702-705. [[Crossref](#)]
448. Jinyoung Kim, Min Sun Choi, Jun Park, Hyunju Park, Hye Won Jang, Jun-Ho Choe, Jung-Han Kim, Jee Soo Kim, Young Seok Cho, Joon Young Choi, Tae Hyuk Kim, Jae Hoon Chung, Sun Wook Kim. 2021. Changes in Thyrotropin Receptor Antibody Levels Following Total Thyroidectomy or Radioiodine Therapy in Patients with Refractory Graves' Disease. *Thyroid* **31**:8, 1264-1271. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
449. Ning Huang, Lin Zeng, Jie Yan, Hongbin Chi, Jie Qiao. 2021. Impact of thyroid cancer treatment on assisted reproductive technology outcomes in women with infertility. *Journal of Assisted Reproduction and Genetics* **38**:8, 2121-2128. [[Crossref](#)]
450. Naykky Singh Ospina, Ramzi G. Salloum, Spyridoula Maraka, Juan P. Brito. 2021. De-implementing low-value care in endocrinology. *Endocrine* **73**:2, 292-300. [[Crossref](#)]
451. Jue Wang, Xiao-Hui Gong, Ting Peng, Jiang-Nan Wu. 2021. Association of Thyroid Function During Pregnancy With the Risk of Pre-eclampsia and Gestational Diabetes Mellitus. *Endocrine Practice* **27**:8, 819-825. [[Crossref](#)]
452. Jackie Gilbert. 2021. Hypothyroidism. *Medicine* **49**:8, 512-514. [[Crossref](#)]
453. Ömercan Topaloğlu, Arzu Yavuz, Ahsen Banu Tiryaki Ayılıkcı. 2021. Evaluation of adherence to levothyroxine and out-of-range thyroid-stimulating hormone levels in pregnant women with primary hypothyroidism. *International Journal of Clinical Practice* **75**:8. . [[Crossref](#)]
454. Andres F. Espinoza, Eyal Krispin, Raphael C. Sun, Jimmy Espinoza, Ahmed Nassr, Alireza A. Shamshirsaz. 2021. Overtreatment of Transient Maternal Hyperthyroidism Resulting in Fetal Goiter. *NeoReviews* **22**:8, e564-e569. [[Crossref](#)]
455. Halina B. Röllin, Kalavati Channa, Bukola Olutola, Jon Øyvind Odland. 2021. Selenium Status, Its Interaction with Selected Essential and Toxic Elements, and a Possible Sex-Dependent Response In Utero, in a South African Birth Cohort. *International Journal of Environmental Research and Public Health* **18**:16, 8344. [[Crossref](#)]

456. Monica Livia Gheorghiu, Roxana Georgiana Bors, Ancuta-Augustina Gheorghisan-Galateanu, Anca Lucia Pop, Dragos Cretoiu, Valentin Nicolae Varlas. 2021. Hyperthyroidism in Pregnancy: The Delicate Balance between Too Much or Too Little Antithyroid Drug. *Journal of Clinical Medicine* **10**:16, 3742. [[Crossref](#)]
457. Sina Jasim, Hengameh Abdi, Hossein Gharib, Bernadette Biondi. 2021. A Clinical Debate: Subclinical Hypothyroidism. *International Journal of Endocrinology and Metabolism* **19**:3. . [[Crossref](#)]
458. F. M. Abdulkhabirova, O. B. Bezlepkina, D. N. Brovin, T. A. Vadina, G. A. Melnichenko, E. V. Nagaeva, L. V. Nikankina, V. A. Peterkova, N. M. Platonova, A. A. Rybakova, T. V. Soldatova, E. A. Troshina, T. Y. Shiryayeva. 2021. Clinical practice guidelines "Management of iodine deficiency disorders". *Problems of Endocrinology* **67**:3, 10-25. [[Crossref](#)]
459. Eren Gürkan, Kenan Dolapçioğlu, Emre Dirican. 2021. Perinatal effects of maternal FT3/FT4 ratio on gestational transient thyrotoxicosis. *Archives of Endocrinology and Metabolism* . [[Crossref](#)]
460. Mihai Dumitrascu, Adina-Elena Nenciu, Sandru Florica, Catalin Nenciu, Aida Petca, Răzvan-Cosmin Petca, Adrian Comănici. 2021. Hyperthyroidism management during pregnancy and lactation (Review). *Experimental and Therapeutic Medicine* **22**:3. . [[Crossref](#)]
461. Sun Y. Lee. 2021. Maternal Iodine Supplementation During Lactation May Improve Child's Cognitive Development at 3 Years of Age in a Mild-to-Moderately Iodine-Deficient Population. *Clinical Thyroidology* **33**:7, 302-304. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
462. P. Merviel, J. Hannigsberg, C. Tremouilhac, S. Herrmann, A.H. Saliou, P.F. Dupré, C. De Moreuil, E. Pasquier, E. Le Moigne, F. Couturaud. 2021. Abortos espontáneos recurrentes. *EMC - Ginecología-Obstetricia* **57**:3, 1-17. [[Crossref](#)]
463. Ahed Hamad, Nawras Alhalabi, Nazht Nmr, Fatima Abbas, Hisham Al-Hammami, Nazir Ibrahim, Marwan Alhalabi. 2021. Impact of Thyroid Autoimmunity in euthyroid women on the outcomes of In Vitro Fertilization. *Annals of Medicine and Surgery* **67**, 102473. [[Crossref](#)]
464. Isha Sekhon, Elizabeth N. Pearce, Xuemei He, Sun Y. Lee. 2021. Iodine Content in Meal Replacements in the United States. *Endocrine Practice* **27**:7, 668-672. [[Crossref](#)]
465. Andrea Busnelli, Federico Cirillo, Paolo Emanuele Levi-Setti. 2021. Thyroid function modifications in women undergoing controlled ovarian hyperstimulation for in vitro fertilization: a systematic review and meta-analysis. *Fertility and Sterility* **116**:1, 218-231. [[Crossref](#)]
466. Tim I.M. Korevaar, Catherine E. de Keyser. 2021. Controlled ovarian hyperstimulation as a stress test for the thyroid. *Fertility and Sterility* **116**:1, 85-86. [[Crossref](#)]
467. Karin Frank-Raue. 2021. Schilddrüse und Schwangerschaft. *Geburtshilfe und Frauenheilkunde* **81**:07, 745-748. [[Crossref](#)]
468. Luca Persani. 2021. Rescue of Neurological Development in Congenital Hypothyroidism: We Should Leave No Stone Unturned. *The Journal of Clinical Endocrinology & Metabolism* **31**. . [[Crossref](#)]
469. Line Sletner, Anne Karen Jenum, Elisabeth Qvigstad, Sara Salehi Hammerstad. 2021. Thyroid Function During Pregnancy in A Multiethnic Population in Norway. *Journal of the Endocrine Society* **5**:7. . [[Crossref](#)]
470. J.G. Vázquez-Rodríguez, A.C. Andrade-Rodríguez. 2021. Enfermedades tiroideas y embarazo en una unidad de cuidados intensivos. Experiencia 2014-2019. *Clínica e Investigación en Ginecología y Obstetricia* **48**:3, 100662. [[Crossref](#)]
471. Won Sang Yoo, Hyun Kyung Chung. 2021. Subclinical Hypothyroidism: Prevalence, Health Impact, and Treatment Landscape. *Endocrinology and Metabolism* **36**:3, 500-513. [[Crossref](#)]
472. Jae Hoon Chung. 2021. Antithyroid Drug Treatment in Graves' Disease. *Endocrinology and Metabolism* **36**:3, 491-499. [[Crossref](#)]
473. Cemal Aktürk. 2021. Bibliometric Analysis of Clinical Decision Support Systems. *Acta Informatica Pragensia* **10**:1, 61-74. [[Crossref](#)]
474. Duygu SEZGİN, Yasemin AYDIN KARTAL. 2021. Gebelik Döneminde Kanıta Dayalı Yaklaşımların Güncel Rehberler Doğrultusunda İncelenmesi. *Ankara Sağlık Bilimleri Dergisi* **10**:1, 92-107. [[Crossref](#)]
475. Almeida A. L. Machamba, Francilene M. Azevedo, Karen O. Fracalossi, Sylvia do C. C. Franceschini. 2021. Effect of iodine supplementation in pregnancy on neurocognitive development on offspring in iodine deficiency areas: a systematic review. *Archives of Endocrinology and Metabolism* **65**:3, 352-367. [[Crossref](#)]
476. Rachita Nanda, Suprava Patel, Prasant Kumar Nayak, Eli Mohapatra, Sarita Agrawal. 2021. Thyroid Peroxidase Antibodies as a Marker of Iodine Status in Healthy Euthyroid Women in First Trimester of Pregnancy Visiting a Tertiary Care Hospital in Raipur, India. *Journal of Evolution of Medical and Dental Sciences* **10**:25, 1857-1861. [[Crossref](#)]
477. Jennifer S. R. Mammen, Anne R. Cappola. 2021. Autoimmune Thyroid Disease in Women. *JAMA* **325**:23, 2392. [[Crossref](#)]
478. Shunli Zhang, Fei Cheng, Hua Wang, Jiangping Wen, Jie Zeng, Chuanbao Zhang, Wensong Liu, Ning Wang, Tingting Jia, Mo Wang, Rui Zhang, Yuhong Yue, Jing Xu, Zhanyong Wang, Yilong Li, Wenxiang Chen, Qingtao Wang. 2021. Comparability of

- thyroid-stimulating hormone immunoassays using fresh frozen human sera and external quality assessment data. *PLOS ONE* **16**:6, e0253324. [[Crossref](#)]
479. Stephen H. LaFranchi. 2021. Thyroid Function in Preterm/Low Birth Weight Infants: Impact on Diagnosis and Management of Thyroid Dysfunction. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]
480. Soumya Thumma, Spyridoula Maraka. 2021. Iodine Supplementation in Pregnant Women with Hashimoto's Thyroiditis. *Clinical Thyroidology* **33**:6, 257-259. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
481. Giovanni Dambrosio, Luigi Laviola. 2021. Trattamento dell'ipertiroidismo di Graves con tionamidi: un documento di sintesi sulle indicazioni e la sicurezza in gravidanza. *L'Endocrinologo* **22**:3, 268-269. [[Crossref](#)]
482. Jörg Bojunga, Ludwig Hofbauer. 2021. Schilddrüse und Schwangerschaft. *Journal für Gynäkologische Endokrinologie/Schweiz* **24**:2, 82-92. [[Crossref](#)]
483. Ayla Coussa, Hayder A. Hasan, Thomas M. Barber. 2021. Early Predictors of Gestational Diabetes Mellitus in IVF-Conceived Pregnancies. *Endocrine Practice* **27**:6, 579-585. [[Crossref](#)]
484. Jing Hua, Jiajin Shen, Jiajia Zhang, Yingchun Zhou, Wenchong Du, Gareth J. Williams. 2021. The association between COVID-19 pandemic and maternal isolated hypothyroxinemia in first and second trimesters. *Psychoneuroendocrinology* **128**, 105210. [[Crossref](#)]
485. Marita Hennessy, Rebecca Dennehy, Sarah Meaney, Laura Linehan, Declan Devane, Rachel Rice, Keelin O'Donoghue. 2021. Clinical practice guidelines for recurrent miscarriage in high-income countries: a systematic review. *Reproductive BioMedicine Online* **42**:6, 1146-1171. [[Crossref](#)]
486. Inez Roest, Kimmy Rosielle, Nienke van Welie, Kim Dreyer, Marlies Bongers, Velja Mijatovic, Ben W. Mol, Carolien Koks. 2021. Safety of oil-based contrast medium for hysterosalpingography: a systematic review. *Reproductive BioMedicine Online* **42**:6, 1119-1129. [[Crossref](#)]
487. Chieh-Chen Wu, Md. Mohaimenul Islam, Phung-Anh Nguyen, Tahmina Nasrin Poly, Ching-Huan Wang, Usman Iqbal, Yu-Chuan (Jack) Li, Hsuan-Chia Yang. 2021. Risk of cancer in long-term levothyroxine users: Retrospective population-based study. *Cancer Science* **112**:6, 2533-2541. [[Crossref](#)]
488. Tejas Kalaria, Anna Sanders, Jonathan Fenn, Helen L. Ashby, Pervaz Mohammed, Harit N. Buch, Clare Ford, Rousseau Gama. 2021. The diagnosis and management of subclinical hypothyroidism is assay-dependent– Implications for clinical practice. *Clinical Endocrinology* **94**:6, 1012-1016. [[Crossref](#)]
489. Sara Salehi Hammerstad, Elisabeth G Celius, Henrik Husby, Ingvild M Sørensen, Ingrid E Norheim. 2021. Management of Severe Graves' Hyperthyroidism in Pregnancy Following Immune Reconstitution Therapy in Multiple Sclerosis. *Journal of the Endocrine Society* **5**:6. . [[Crossref](#)]
490. Nikolina Docheva, Erin Tran. 2021. A Diagnosis of Maternal Hyperthyroidism. *NeoReviews* **22**:6, e409-e414. [[Crossref](#)]
491. Nikolaos Vrachnis, Orestis Tsonis, Dionisios Vrachnis, Nikolaos Antonakopoulos, George Paltoglou, Stavroula Barbounaki, George Mastorakos, Minas Paschopoulos, Zoi Iliodromiti. 2021. The Effect of Thyrotropin-Releasing Hormone and Antithyroid Drugs on Fetal Thyroid Function. *Children* **8**:6, 454. [[Crossref](#)]
492. José María Hernández, Berta Soldevila, Inés Velasco, Fernando Moreno-Flores, Laura Ferrer, Alejandra Pérez-Montes de Oca, Cecilia Santillán, Carla Muñoz, Sílvia Ballesta, Cristina Canal, Manel Puig-Domingo, María Luisa Granada. 2021. Reference Intervals of Thyroid Function Tests Assessed by Immunoassay and Mass Spectrometry in Healthy Pregnant Women Living in Catalonia. *Journal of Clinical Medicine* **10**:11, 2444. [[Crossref](#)]
493. Silvia González-Martínez, María Riestra-Fernández, Eduardo Martínez-Morillo, Noelia Avello-Llano, Elías Delgado-Álvarez, Edelmiro Luis Menéndez-Torre. 2021. Nutritional Iodine Status in Pregnant Women from Health Area IV in Asturias (Spain): Iodised Salt Is Enough. *Nutrients* **13**:6, 1816. [[Crossref](#)]
494. James B. Adams, Jacob C. Sorenson, Elena L. Pollard, Jasmine K. Kirby, Tapan Audhya. 2021. Evidence-Based Recommendations for an Optimal Prenatal Supplement for Women in the U.S., Part Two: Minerals. *Nutrients* **13**:6, 1849. [[Crossref](#)]
495. Jia-Wei Wang, Xiao-Xian Liao, Tuo Li. 2021. Thyroid autoimmunity in adverse fertility and pregnancy outcomes: Timing of assisted reproductive technology in AITD women. *Journal of Translational Internal Medicine* **9**:2, 76-83. [[Crossref](#)]
496. Juergen Kratzsch, Nikola A. Baumann, Ferruccio Ceriotti, Zhong X. Lu, Matthias Schott, Antonius E. van Herwaarden, José Gilberto Henriques Vieira, Dusanka Kasapic, Luca Giovanella. 2021. Global FT4 immunoassay standardization: an expert opinion review. *Clinical Chemistry and Laboratory Medicine (CCLM)* **59**:6, 1013-1023. [[Crossref](#)]
497. Yunhai Li, Cheng Jin, Jie Li, Mingkun Tong, Mengxue Wang, Jiefeng Huang, Yi Ning, Guosheng Ren. 2021. Prevalence of Thyroid Nodules in China: A Health Examination Cohort-Based Study. *Frontiers in Endocrinology* **12**. . [[Crossref](#)]

498. Oshini Mallawa Kankanamalage, Qiongjie Zhou, Xiaotian Li. 2021. Understanding the Pathogenesis of Gestational Hypothyroidism. *Frontiers in Endocrinology* 12. . [\[Crossref\]](#)
499. Dustin R Bunch, Kyle Firmender, Roa Harb, Joe M El-Khoury. 2021. First- and Second-Trimester Reference Intervals for Thyroid Function Testing in a US Population. *American Journal of Clinical Pathology* 155:6, 776-780. [\[Crossref\]](#)
500. Lauren N. Krumeich, Rachel R. Kelz. 2021. Editorial: Time to Surgery and Thyroid Cancer Survival in the United States. *Annals of Surgical Oncology* 84. . [\[Crossref\]](#)
501. Yanrui Huang, Haixia Guan. 2021. High Serum TRAb Levels at Birth May Result in Various Types of Neonatal Thyroid Dysfunction. *Clinical Thyroidology* 33:5, 213-216. [\[Citation\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
502. Patricia Lemieux, Jennifer M. Yamamoto, Kara A. Nerenberg, Amy Metcalfe, Alex Chin, Rshmi Khurana, Lois Elizabeth Donovan. 2021. Thyroid Laboratory Testing and Management in Women on Thyroid Replacement Before Pregnancy and Associated Pregnancy Outcomes. *Thyroid* 31:5, 841-849. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#) [\[Supplementary Material\]](#)
503. Lei Tang, Ping Li, Hua Zhou, Ling Li. 2021. A longitudinal study of thyroid markers during pregnancy and the risk of gestational diabetes mellitus and post-partum glucose metabolism. *Diabetes/Metabolism Research and Reviews* 37:4. . [\[Crossref\]](#)
504. Cheng Huang, Ying Wu, Linong Chen, Zhiya Yuan, Shuzhe Yang, Chenggui Liu. 2021. Establishment of assay method- and trimester-specific reference intervals for thyroid hormones during pregnancy in Chengdu, China. *Journal of Clinical Laboratory Analysis* 35:5. . [\[Crossref\]](#)
505. Bassem Refaat, Firas Azzeh. 2021. Factors Associated with Thyroid Disorders and Iodine Adequacy in Pregnant Saudi Women. *Biological Trace Element Research* 199:5, 1715-1728. [\[Crossref\]](#)
506. Francisco J. Barrera, Karina Raygoza-Cortez, Mariana García-Leal, Juan P. Brito, Naykky M. Singh Ospina, René Rodríguez-Gutiérrez. 2021. Are American follow-up recommendations in endocrinology actionable? A systematic review of clinical practice guidelines. *Endocrine* 72:2, 375-384. [\[Crossref\]](#)
507. Juliane Léger. 2021. Dépistage de l'hypothyroïdie congénitale. *médecine/sciences* 37:5, 474-481. [\[Crossref\]](#)
508. Dária Madej, Pauline Wimberger, Elena Tsourdi, Maren Goeckenjan. 2021. Präkonzeptionelle Beratung bei Diabetes mellitus. *Diabetes aktuell* 19:03, 102-109. [\[Crossref\]](#)
509. Çiğdem Tura Bahadır, Merve Yilmaz, Elif Kılıçkan. 2021. Free triiodothyronine to free thyroxine ratio in the differential diagnosis of thyrotoxicosis and hyperthyroidism: A retrospective study. *International Journal of Clinical Practice* 75:5. . [\[Crossref\]](#)
510. Alessandro Dal Lago, Francesco Galanti, Donatella Miriello, Antonella Marcoccia, Micol Massimiani, Luisa Campagnolo, Costanzo Moretti, Rocco Rago. 2021. Positive Impact of Levothyroxine Treatment on Pregnancy Outcome in Euthyroid Women with Thyroid Autoimmunity Affected by Recurrent Miscarriage. *Journal of Clinical Medicine* 10:10, 2105. [\[Crossref\]](#)
511. Arri Coomarasamy, Rima K Dhillon-Smith, Argyro Papadopoulou, Maya Al-Memar, Jane Brewin, Vikki M Abrahams, Abha Maheshwari, Ole B Christiansen, Mary D Stephenson, Mariëtte Goddijn, Olufemi T Oladapo, Chandrika N Wijeyaratne, Debra Bick, Hassan Shehata, Rachel Small, Phillip R Bennett, Lesley Regan, Raj Rai, Tom Bourne, Rajinder Kaur, Oonagh Pickering, Jan J Brosens, Adam J Devall, Ioannis D Gallos, Siobhan Quenby. 2021. Recurrent miscarriage: evidence to accelerate action. *The Lancet* 397:10285, 1675-1682. [\[Crossref\]](#)
512. K Aaron Geno, Matthew S Reed, Mark A Cervinski, Robert D Nerenz. 2021. Evaluation of Thyroid Function in Pregnant Women Using Automated Immunoassays. *Clinical Chemistry* 67:5, 772-780. [\[Crossref\]](#)
513. Samantha M Siskind, Sun Y Lee, Elizabeth N Pearce. 2021. Investigating hypothyroidism. *BMJ* 129, n993. [\[Crossref\]](#)
514. Sue Marty, Manon Beekhuijzen, Alex Charlton, Nina Hallmark, Bethany R. Hannas, Sylvia Jacobi, Stephanie Melching-Kollmuss, Ursula G. Sauer, Larry P. Sheets, Volker Strauss, Daniel Urbisch, Philip A. Botham, Bennard van Ravenzwaay. 2021. Towards a science-based testing strategy to identify maternal thyroid hormone imbalance and neurodevelopmental effects in the progeny – part II: how can key events of relevant adverse outcome pathways be addressed in toxicological assessments?. *Critical Reviews in Toxicology* 51:4, 328-358. [\[Crossref\]](#)
515. Mohammad Moin Shahid, K. M. Thouhidur Rahman, Richmond Ronald Gomes, Menoka Ferdous, Sharmin Ferdousi, Tamannur Zahan. 2021. Association of gestational diabetes mellitus and thyroid status during pregnancy: a cross-sectional study in a tertiary health care center of Bangladesh. *Gynecological Endocrinology* 37:4, 312-314. [\[Crossref\]](#)
516. Zhirou Chen, Xi Yang, Chen Zhang, Zheng Ding, Yong Zhang, Tim I.M. Korevaar, Jianxia Fan. 2021. Thyroid Function Test Abnormalities in Twin Pregnancies. *Thyroid* 31:4, 572-579. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#) [\[Supplementary Material\]](#)
517. Freddy J.K. Toloza, Sarah E. Theriot, Naykky M. Singh Ospina, Sameen Nooruddin, Brooke Keathley, Stacey M. Johnson, Nalin Payakachat, Elena Ambrogini, Rene Rodriguez-Gutierrez, Derek T. O'Keeffe, Juan P. Brito, Victor M. Montori, Nafisa K. Dajani, Spyridoula Maraka. 2021. Knowledge, Attitudes, Beliefs, and Treatment Burden Related to the Use of Levothyroxine

- in Hypothyroid Pregnant Women in the United States. *Thyroid* 31:4, 669-677. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
518. Ying Yang, Tonglei Guo, Jinrong Fu, Jian Kuang, Yuanyuan Wang, Ya Zhang, Hongguang Zhang, Yuan He, Zuoqi Peng, Qiaomei Wang, Haiping Shen, Yiping Zhang, Donghai Yan, Xu Ma, Haixia Guan. 2021. Preconception Thyrotropin Levels and Risk of Adverse Pregnancy Outcomes in Chinese Women Aged 20 to 49 Years. *JAMA Network Open* 4:4, e215723. [[Crossref](#)]
 519. Hortense Didier-Mathon, Hanane Bouchghoul, Marie-Victoire Senat, Jacques Young, Dominique Luton. 2021. Prenatal management of fetal goiter alternating between hypothyroidism and hyperthyroidism in a mother with Graves' disease. *Clinical Case Reports* 9:4, 2281-2284. [[Crossref](#)]
 520. Liliana Fonseca, Vânia Silva Benido, João Pessanha, Maria Teresa Pereira, Joana Vilaverde, Jorge Soares. 2021. Being alert on thyroid function tests interpretation: A case report of heterophile antibody interference in serum in a pregnant woman. *Annales d'Endocrinologie* 82:2, 121-123. [[Crossref](#)]
 521. Cord Sturgeon, Brian W. Kim. 2021. An evolving role for GRADE-trained methodologists in the American Thyroid Association Clinical Practice Guidelines Development Process. *Journal of Clinical Epidemiology* 132, 147-148. [[Crossref](#)]
 522. Lynne T. Haber, Rita S. Schoeny, Bruce C. Allen. 2021. Impact of updated BMD modeling methods on perchlorate and chlorate assessments of human health hazard. *Toxicology Letters* 340, 89-100. [[Crossref](#)]
 523. Chaochao Ma, Xiaoqi Li, Lixin Liu, Xinqi Cheng, Fang Xue, Jie Wu, Liangyu Xia, Yicong Yin, Danchen Wang, Yutong Zou, Ling Qiu, Juntao Liu. 2021. Establishment of Early Pregnancy Related Thyroid Hormone Models and Reference Intervals for Pregnant Women in China Based on Real World Data. *Hormone and Metabolic Research* 53:04, 272-279. [[Crossref](#)]
 524. Jayne V. Woodside, Karen R. Mullan. 2021. Iodine status in UK—An accidental public health triumph gone sour. *Clinical Endocrinology* 94:4, 692-699. [[Crossref](#)]
 525. I. A. Tsanova, S. V. Bulgakova, A. V. Melikova. 2021. Subclinical hypothyroidism: treat or watch?. *Bulletin of the Medical Institute "REAVIZ" (REHABILITATION, DOCTOR AND HEALTH)* :6, 98-108. [[Crossref](#)]
 526. Sanjay Kalra, Sameer Aggarwal, Deepak Khandelwal. 2021. Thyroid Dysfunction and Dysmetabolic Syndrome: The Need for Enhanced Thyrovigilance Strategies. *International Journal of Endocrinology* 2021, 1-11. [[Crossref](#)]
 527. Shriya Ganju. 2021. Predictive value of thyroid stimulating hormone as a biomarker in intrauterine growth restriction. *Indian Journal of Obstetrics and Gynecology Research* 8:1, 53-56. [[Crossref](#)]
 528. Ashley S. P. Boggs, Lisa E. Kilpatrick, Carolyn Q. Burdette, Denise S. Tevis, Zachary A. Fultz, Michael A. Nelson, Jeffery M. Jarrett, Jennifer V. Kemp, Ravinder J. Singh, Stefan K. G. Grebe, Stephen A. Wise, Brittany L. Kassim, Stephen E. Long. 2021. Development of a pregnancy-specific reference material for thyroid biomarkers, vitamin D, and nutritional trace elements in serum. *Clinical Chemistry and Laboratory Medicine (CCLM)* 59:4, 671-679. [[Crossref](#)]
 529. Elizabeth O Buschur, Sarit Polsky. 2021. Type 1 Diabetes: Management in Women From Preconception to Postpartum. *The Journal of Clinical Endocrinology & Metabolism* 106:4, e952-e967. [[Crossref](#)]
 530. Elizabeth N. Pearce. 2021. Comparison of ATA and Updated ACOG Guidelines for Thyroid Disease in Pregnancy. Russian translation. *Clinical and experimental thyroidology* 16:3, 12-15. [[Crossref](#)]
 531. Delshad Hossein, Azizi Fereidoun. 2021. Mild to moderate iodine deficiency in pregnancy: A matter of debate. *Archives of Food and Nutritional Science* 5:1, 018-026. [[Crossref](#)]
 532. Sun Y Lee, Elizabeth N Pearce. 2021. Testing, Monitoring, and Treatment of Thyroid Dysfunction in Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* 106:3, 883-892. [[Crossref](#)]
 533. Owain Leng, Salman Razvi. 2021. Treatment of subclinical hypothyroidism: assessing when treatment is likely to be beneficial. *Expert Review of Endocrinology & Metabolism* 16:2, 73-86. [[Crossref](#)]
 534. Riley Epp, Janine Malcolm, Khiera Jolin-Dahel, Michaela Clermont, Erin Keely. 2021. Postpartum thyroiditis. *BMJ* 39, n495. [[Crossref](#)]
 535. Paul van Trotsenburg, Athanasia Stoupa, Juliane Léger, Tilman Rohrer, Catherine Peters, Laura Fugazzola, Alessandra Cassio, Claudine Heinrichs, Veronique Beauloye, Joachim Pohlenz, Patrice Rodien, Régis Coutant, Gabor Szinnai, Philip Murray, Beate Bartés, Dominique Luton, Mariacarla Salerno, Luisa de Sanctis, Mariacristina Vigone, Heiko Krude, Luca Persani, Michel Polak. 2021. Congenital Hypothyroidism: A 2020–2021 Consensus Guidelines Update—An ENDO-European Reference Network Initiative Endorsed by the European Society for Pediatric Endocrinology and the European Society for Endocrinology. *Thyroid* 31:3, 387-419. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
 536. Simona Gaberšček, Brina Gaberšček, Katja Zaletel. 2021. Incidence of thyroid disorders in the second decade of adequate iodine supply in Slovenia. *Wiener klinische Wochenschrift* 133:5-6, 182-187. [[Crossref](#)]

537. Serena Girardelli, Giorgia Mangili, Stefania Cosio, Emanuela Rabaiotti, Antonio Fanucchi, Luca Valsecchi, Massimo Candiani, Angiolo Gadducci. 2021. A narrative review of pregnancy after malignancies in young women that don't originate in the female genital organs or in the breast. *Critical Reviews in Oncology/Hematology* **159**, 103240. [[Crossref](#)]
538. Eduardo Rodriguez-Diaz, Jinetsy I. Rivera-Ortiz, Sun Y. Lee, Loida A. Gonzalez-Rodriguez, Xuemei He, Elizabeth N. Pearce. 2021. Iodine Status in Pregnant Women of Puerto Rico. *Endocrine Practice* **27**:3, 241-244. [[Crossref](#)]
539. Karen Tsai, Angela M. Leung. 2021. Subclinical Hyperthyroidism: A Review of the Clinical Literature. *Endocrine Practice* **27**:3, 254-260. [[Crossref](#)]
540. Timo Deutschbein, Cornelia Jaursch-Hancke, Ulrich J. Knappe, Wolfgang Saeger, Jörg Flitsch, Jörg Bojunga, Michael Buchfelder, Beate Ditzen, Rüdiger Gerlach, Elfriede Gertzen, Jürgen Honegger, Gerhard A. Horstmann, Arend Koch, Ilonka Kreitschmann-Andermahr, Mirjam Kunz, Wolf A. Lagrèze, Nils H. Nicolay, Werner Paulus, Martin Reincke, Manuel A. Schmidt, Matthias M. Weber, Helmut Wilhelm, Martin Fassnacht. 2021. First German Guideline on Diagnostics and Therapy of Clinically Non-Functioning Pituitary Tumors. *Experimental and Clinical Endocrinology & Diabetes* **129**:03, 250-264. [[Crossref](#)]
541. Shuting Si, Yu Shen, Xing Xin, Minjia Mo, Bule Shao, Shuojia Wang, Wenliang Luo, Zexin Chen, Hui Liu, Danqing Chen, Yunxian Yu. 2021. Hemoglobin concentration and iron supplement during pregnancy were associated with an increased risk of gestational diabetes mellitus. *Journal of Diabetes* **13**:3, 211-221. [[Crossref](#)]
542. Hiroyuki Yoshihara, Mayumi Sugiura-Ogasawara, Shinobu Goto, Tamao Kitaori. 2021. Levothyroxine and subclinical hypothyroidism in patients with recurrent pregnancy loss. *American Journal of Reproductive Immunology* **85**:3. . [[Crossref](#)]
543. Stine Linding Andersen, Stig Andersen, Zeyan Liew, Peter Vestergaard, Søren Lundbye-Christensen, Thorkild I. A. Sørensen, Jørn Olsen. 2021. Maternal thyroid disease and adiposity in mother and child. *Clinical Endocrinology* **94**:3, 484-493. [[Crossref](#)]
544. Ai Yoshihara, Jaeduk Yoshimura Noh, Natsuko Watanabe, Miho Fukushima, Masako Matsumoto, Nami Suzuki, Ayako Hoshiyama, Ai Suzuki, Takako Mitsumatsu, Aya Kinoshita, Kentaro Mikura, Ran Yoshimura, Kiminori Sugino, Koichi Ito. 2021. Exposure to Propylthiouracil in the First Trimester of Pregnancy and Birth Defects: A Study at a Single Institution. *Journal of the Endocrine Society* **5**:3. . [[Crossref](#)]
545. Lidia Arce-Sánchez, Salvatore Giovanni Vitale, Claudia Montserrat Flores-Robles, Myrna Souraye Godines-Enriquez, Marco Noventa, Carmen Marcela Urquía-Figueroa, Nayeli Martínez-Cruz, Guadalupe Estrada-Gutierrez, Salvador Espino y Sosa, José Romo-Yañez, Araceli Montoya-Estrada, Enrique Reyes-Muñoz. 2021. Effect of the Cut-Off Level for Thyroid-Stimulating Hormone on the Prevalence of Subclinical Hypothyroidism among Infertile Mexican Women. *Diagnostics* **11**:3, 417. [[Crossref](#)]
546. Maria Teresa Murillo-Llorente, Francisco Llorca-Colomer, Marcelino Pérez-Bermejo. 2021. Relationship between Thyroid Status during the First Trimester of Pregnancy and Neonatal Well-Being. *Nutrients* **13**:3, 872. [[Crossref](#)]
547. Heraldio Mendes Garmes, César Luiz Boguszewski, Paulo Augusto Carvalho Miranda, Manoel Ricardo Alves Martins, Silvia Regina Correa da Silva, Julio Zaki Abucham, Nina Rosa de Castro Musolino, Lucio Vilar, Luiz Henrique Corrêa Portari, Mônica Roberto Gadelha, Leandro Kasuki, Luciana Ansaneli Naves, Mauro Antônio Czepielewski, Tobias Skrebsky de Almeida, Felipe Henning Gaia Duarte, Andrea Glezer, Marcello Delano Bronstein. 2021. Management of hypopituitarism: a perspective from the Brazilian Society of Endocrinology and Metabolism. *Archives of Endocrinology and Metabolism* . [[Crossref](#)]
548. Wenjun Cao, Chenghan Luo, Mengyuan Lei, Min Shen, Wenqian Ding, Mengmeng Wang, Min Song, Jian Ge, Qian Zhang. 2021. Development and Validation of a Dynamic Nomogram to Predict the Risk of Neonatal White Matter Damage. *Frontiers in Human Neuroscience* **14**. . [[Crossref](#)]
549. Vikas Yadav, Deepti Dabar, Akhil D. Goel, Mohan Bairwa, Akanksha Sood, Pankaj Prasad, Sanjay S. Agarwal, Sunil Nandeshwar. 2021. Prevalence of Hypothyroidism in Pregnant Women in India: A Meta-Analysis of Observational Studies. *Journal of Thyroid Research* **2021**, 1-19. [[Crossref](#)]
550. Geng-Dong Chen, Ting-Ting Pang, Xia-Fen Lu, Peng-Sheng Li, Zi-Xing Zhou, Shao-Xin Ye, Jie Yang, Xiu-Yin Shen, Dong-Xin Lin, Da-Zhi Fan, De-Mei Lu, Zheng-Ping Liu. 2021. Associations Between Maternal Thyroid Function and Birth Outcomes in Chinese Mother-Child Dyads: A Retrospective Cohort Study. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
551. Freddy J.K. Toloza, Spyridoula Maraka. 2021. Isolated Maternal Hypothyroxinemia Is Associated with Preterm Delivery. *Clinical Thyroidology* **33**:2, 80-82. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
552. Cari M Kitahara, Dagrun Slettebø Daltveit, Anders Ekblom, Anders Engeland, Mika Gissler, Ingrid Glimelius, Tom Grotmol, Ylva Trolle Lagerros, Laura Madanat-Harjuoja, Tuija Männistö, Henrik Toft Sørensen, Rebecca Troisi, Tone Bjørge. 2021. Maternal health, in-utero, and perinatal exposures and risk of thyroid cancer in offspring: a Nordic population-based nested case-control study. *The Lancet Diabetes & Endocrinology* **9**:2, 94-105. [[Crossref](#)]
553. Koji Sakamaki, Katsuhiko Tsunekawa, Nobuyoshi Ishiyama, Mizuho Kudo, Kimiko Ando, Masako Akuzawa, Katsuyuki Nakajima, Yohnosuke Shimomura, Osamu Araki, Takao Kimura, Masami Murakami. 2021. Association between high normal-range

- thyrotropin concentration and carotid intima-media thickness in euthyroid premenopausal, perimenopausal and postmenopausal women. *Maturitas* **144**, 29-36. [[Crossref](#)]
554. Simone Canovi, Silvia Vezzani, Alessandra Polese, Andrea Frasoldati, Clara Schiatti, Cristina Preda, Michele Corradini Zini, Antonio Vitiello, Marco Foracchia, Giuseppina Comitini, Lorenzo Aguzzoli, Tommaso Fasano, Luigi Vecchia. 2021. Pregnancy-related reference intervals for serum thyrotropin based on real-life clinical data. *Gynecological Endocrinology* **37**:2, 113-116. [[Crossref](#)]
 555. Lorraine Lau, Jamie L Benham, Patricia Lemieux, Jennifer Yamamoto, Lois E Donovan. 2021. Impact of levothyroxine in women with positive thyroid antibodies on pregnancy outcomes: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* **11**:2, e043751. [[Crossref](#)]
 556. Katsuhiko Hamada, Tetsuya Mizokami, Tetsushi Maruta, Kiichiro Higashi, Kaoru Konishi, Naoko Momotani, Junichi Tajiri. 2021. Thyroid Function of Infants Breastfed by Mothers with Graves Disease Treated with Inorganic Iodine: A Study of 100 Cases. *Journal of the Endocrine Society* **5**:2. . [[Crossref](#)]
 557. Kilian Vomstein, Katharina Feil, Laura Strobel, Anna Aulitzky, Susanne Hofer-Tollinger, Ruben-Jeremias Kuon, Bettina Toth. 2021. Immunological Risk Factors in Recurrent Pregnancy Loss: Guidelines Versus Current State of the Art. *Journal of Clinical Medicine* **10**:4, 869. [[Crossref](#)]
 558. Myrna Souraye Godines-Enriquez, Silvia Miranda-Velásquez, María Magdalena Enriquez-Pérez, Lidia Arce-Sánchez, Nayeli Martínez-Cruz, Claudia Montserrat Flores-Robles, Patricia Aguayo-González, Fela Vanessa Morales-Hernández, Alma Villarreal-Barranca, Blanca Vianey Suárez-Rico, Araceli Montoya-Estrada, José Romo-Yáñez, Enrique Reyes-Muñoz. 2021. Prevalence of Thyroid Autoimmunity in Women with Recurrent Pregnancy Loss. *Medicina* **57**:2, 96. [[Crossref](#)]
 559. Iwona Krela-Kaźmierczak, Agata Czarnywojtek, Kinga Skoracka, Anna Maria Rychter, Alicja Ewa Ratajczak, Aleksandra Szymczak-Tomczak, Marek Ruchała, Agnieszka Dobrowolska. 2021. Is There an Ideal Diet to Protect against Iodine Deficiency?. *Nutrients* **13**:2, 513. [[Crossref](#)]
 560. Huijia Lin, Mudan Ye, Zhenwen Zhou, Lianxiong Yuan, Gendie E. Lash, Guanglan Zhang, Li Li. 2021. Reference values and the effect of clinical parameters on thyroid hormone levels during early pregnancy. *Bioscience Reports* **41**:1. . [[Crossref](#)]
 561. Divya M Mathews, Neil P Johnson, Robert G Sim, Susannah O'Sullivan, Jane M Peart, Paul L Hofman. 2021. Iodine and fertility: do we know enough?. *Human Reproduction* **36**:2, 265-274. [[Crossref](#)]
 562. Jun Li, Litao Bai, Fan Wei, Maoying Wei, Yao Xiao, Weitian Yan, Junping Wei. 2021. Effect of Addition of Thyroxine in the Treatment of Graves' Disease: A Systematic Review. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
 563. S. V. Khabarov. 2021. Screening determination of thyroid stimulating hormone in diagnosis of subclinical hypothyroidism in pregnant women: versions and contraversions. *Medical alphabet* :34, 59-65. [[Crossref](#)]
 564. Victor Pop, Johannes Krabbe, Wolfgang Maret, Margaret Rayman. 2021. Plasma mineral (selenium, zinc or copper) concentrations in the general pregnant population, adjusted for supplement intake, in relation to thyroid function. *British Journal of Nutrition* **125**:1, 71-78. [[Crossref](#)]
 565. Danrong Chen, Jiani Liu, Wu Yan, Kacey Fang, Yankai Xia, Wei Lv, Zhonghua Shi. 2021. Associations of Prenatal Exposure to Triclosan and Maternal Thyroid Hormone Levels: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
 566. T van der Made, M van de Vyver, M Conradie-Smit, Magda Conradie. 2021. Prevalence and aetiology of thyrotoxicosis in patients with hyperemesis gravidarum presenting to a tertiary hospital in Cape Town, South Africa. *Journal of Endocrinology, Metabolism and Diabetes of South Africa* **26**:1, 1-8. [[Crossref](#)]
 567. Anna M. Sawka, Erik K. Alexander, Antonio C. Bianco, Roger Chou, Bryan R. Haugen, Peter A. Kopp, Elizabeth N. Pearce, Douglas S. Ross, Robert C. Smallridge, Jacqueline Jonklaas. 2021. Challenges in Developing Recommendations Based on Low-Quality Evidence in Thyroid Guidelines. *Thyroid* **31**:1, 3-7. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
 568. Sarah E. Cusick, Amanda Barks, Michael K. Georgieff. Nutrition and Brain Development 131-165. [[Crossref](#)]
 569. Julia R. Donner, Lisa Swartz Topor. Thyroid Studies 149-156. [[Crossref](#)]
 570. Lorraine Lau, Ralf Paschke. Antithyroid Drugs 212-218. [[Crossref](#)]
 571. Zachary Simons, David S. Cooper. Papillary Thyroid Carcinoma Diagnosed During Pregnancy 141-145. [[Crossref](#)]
 572. Takashi Akamizu. Practical Application of Levothyroxine-Based Therapy 121-129. [[Crossref](#)]
 573. Hans-Peter Lipp. Administration and Pharmacokinetics of Levothyroxine 13-22. [[Crossref](#)]
 574. Kris Gustave Poppe. Levothyroxine in Pregnancy 47-60. [[Crossref](#)]
 575. Aura Meliza Mejia Monroy. Thyroid Emergency and Pregnancy 139-149. [[Crossref](#)]

576. Sabrina Corbetta. Classification of Thyroid Diseases 21-35. [[Crossref](#)]
577. Hang Zhou, Yiming Lu, Binyu Pan, Qihua Zhao, Zheng Fei Ma. 2021. Iodine Deficiency as Assessed by Neonatal TSH in a Sample of Mother-and-Newborn Pairs in Jiangsu Province, China. *Biological Trace Element Research* **199**:1, 70-75. [[Crossref](#)]
578. K. Aaron Geno, Mark A. Cervinski, Robert D. Nerenz. Pregnancy and the fetus 543-579. [[Crossref](#)]
579. Johnny Deladoey, Julia Elisabeth von Oettingen, Guy Van Vliet. Disorders of the Thyroid in the Newborn and Infant 202-219. [[Crossref](#)]
580. Angela M. Leung, Gregory A. Brent. Thyroid Physiology and Thyroid Function Testing 26-38.e3. [[Crossref](#)]
581. Trevor E. Angell, Matthew I. Kim, Victor J. Bernet. Thyroiditis 40-49.e4. [[Crossref](#)]
582. Whitney Liddy, Juliana Bonilla-Velez, Frédéric Triponez, Dipti Kamani, Gregory Randolph. Principles in Thyroid Surgery 272-293.e5. [[Crossref](#)]
583. Lisa Antonia Dröge, Thorsten Fischer, Dieter Grab, Thomas Grubert, Peer Hantschmann, Wolfgang Henrich, Ralph Kästner, Franz Kainer, Christina Kentenich, Walter Klockenbusch, Frank Lammert, Frank Louwen, Ioannis Mylonas, Stephanie Pildner von Steinburg, Ute M. Schäfer-Graf, Ekkehard Schleußner, Ralf Schmitz, Stefan Verlohren. Maternale Erkrankungen in der Schwangerschaft 337-609. [[Crossref](#)]
584. Jesse Bertinato. Iodine nutrition: Disorders, monitoring and policies 365-415. [[Crossref](#)]
585. Jean M. Kerver, Elizabeth N. Pearce, Tengfei Ma, Monica Gentchev, Michael R. Elliott, Nigel Paneth. 2021. Prevalence of inadequate and excessive iodine intake in a US pregnancy cohort. *American Journal of Obstetrics and Gynecology* **224**:1, 82.e1-82.e8. [[Crossref](#)]
586. Qian Zhu, Qian-Han Xu, Ting Xie, Li-Ling Wang, Hong Liu, Kahindo P. Muyayalo, Xiao-Bo Huang, Si-Jia Zhao, Ai-Hua Liao. 2021. Recent insights into the impact of immune dysfunction on reproduction in autoimmune thyroiditis. *Clinical Immunology* 108663. [[Crossref](#)]
587. Charles Bitamazire Businge, Benjamin Longo-Mbenza, Andre Pascal Kengne. 2021. Mildly elevated thyroid-stimulating hormone is associated with endothelial dysfunction and severe preeclampsia among pregnant women with insufficient iodine intake in Eastern Cape province, South Africa. *Annals of Medicine* **53**:1, 1084-1090. [[Crossref](#)]
588. Zareen Kiran, Aisha Sheikh, Khadija Nuzhat Humayun, Najmul Islam. 2021. Neonatal outcomes and congenital anomalies in pregnancies affected by hypothyroidism. *Annals of Medicine* **53**:1, 1560-1568. [[Crossref](#)]
589. Luyang Han, Yan Ma, Zhaoxia Liang, Danqing Chen. 2021. Laboratory characteristics analysis of the efficacy of levothyroxine on subclinical hypothyroidism during pregnancy: a single-center retrospective study. *Bioengineered* **12**:1, 4183-4190. [[Crossref](#)]
590. Yongchi Zhan, Tingting Xu, Xiaodong Wang. 2021. Perinatal Management and Outcomes of Pregnancy Following Sheehan Syndrome: A Case Report and Literature Review. *Maternal-Fetal Medicine* **3**:3, 213-220. [[Crossref](#)]
591. Divya M. Mathews, Jane M. Peart, Robert G. Sim, Neil P. Johnson, Susannah O'Sullivan, José G.B. Derraik, Paul L. Hofman. 2021. The effect of acute and chronic iodine excess on thyroid profile and reproductive function of women using Lipiodol during hysterosalpingography and the potential impact on thyroid function of their offspring: The SELF study protocol. *Medicine: Case Reports and Study Protocols* **2**:8, e0148. [[Crossref](#)]
592. Sofia Manousou, Hanna Augustin, Robert Eggertsen, Lena Hulthén, Helena Filipsson Nyström. 2021. Inadequate iodine intake in lactating women in Sweden: A pilot 1-year, prospective, observational study. *Acta Obstetrica et Gynecologica Scandinavica* **100**:1, 48-57. [[Crossref](#)]
593. Solène Castellnou, Patricia Bretones, Juliette Abeillon, Myriam Moret, Pauline Perrin, Karim Chikh, Véronique Raverot. 2021. Congenital Hypothyroidism due to a Low Level of Maternal Thyrotropin Receptor-Blocking Antibodies. *European Thyroid Journal* **10**:2, 174-178. [[Crossref](#)]
594. Hiroyuki Iwaki, Kenji Ohba, Eisaku Okada, Takeshi Murakoshi, Yumiko Kashiwabara, Chiga Hayashi, Akio Matsushita, Shigekazu Sasaki, Takafumi Suda, Yutaka Oki, Rieko Gemma. 2021. Dose-Dependent Influence of Antithyroid Drugs on the Difference in Free Thyroxine Levels between Mothers with Graves' Hyperthyroidism and Their Neonates. *European Thyroid Journal* **10**:5, 372-381. [[Crossref](#)]
595. Constance Repelaer van Driel-Delprat, Eveline van Dam, Peter van de Ven, Khadija Aissa, Melanie ter Haar, Yikke Feenstra, Aletta de Roos, Gaby Beelen, Roel Schats, Cornelis Lambalk. 2021. More Live Births in Primary Subfertile Intracytoplasmic Sperm Injection-Treated Women with High Normal TSH Levels. *Gynecologic and Obstetric Investigation* **86**:4, 398-407. [[Crossref](#)]
596. Ping Zhu, Ruifeng Chu, Shilei Pan, Xiaolan Lai, Jianmin Ran, Xiuhong Li. 2021. Impact of TPOAb-negative maternal subclinical hypothyroidism in early pregnancy on adverse pregnancy outcomes. *Therapeutic Advances in Endocrinology and Metabolism* **12**, 204201882110546. [[Crossref](#)]

- 597.## #. 2021. Stratified Analysis of the Effects of Subclinical Hypothyroidism on the Clinical Outcomes of Assisted Pregnancy Techniques. *Advances in Clinical Medicine* 11:01, 312-318. [[Crossref](#)]
- 598.## #. 2021. Study on Reference Value Range of Serum Thyroid Hormones in Pregnancy in Zhangye, Gansu Province. *Advances in Clinical Medicine* 11:11, 5279-5283. [[Crossref](#)]
- 599.## #. 2021. Predictive Value of TRAb in the Early Drug Treatment of GD. *Advances in Clinical Medicine* 11:12, 5827-5834. [[Crossref](#)]
600. Laura Linehan, Marita Hennessy, Keelin O'Donoghue. 2021. Infertility and subsequent recurrent miscarriage: Current state of the literature and future considerations for practice and research. *HRB Open Research* 4, 100. [[Crossref](#)]
601. Asako Tsunemi, Toyoyoshi Uchida, Keiji Kuroda, Yuko Ikemoto, Asako Ochiai, Hiromasa Goto, Rikikazu Sugiyama, Hiroaki Satoh, Atsuo Itakura, Hirotaka Watada. 2021. Effect of thyroxine treatment on pregnancy outcomes in infertile Japanese women with TSH levels between 2.5 μ IU/mL and the upper reference limit: a retrospective study. *Endocrine Journal* 68:2, 171-177. [[Crossref](#)]
602. 2021. 14. Management of Diabetes in Pregnancy: Standards of Medical Care in Diabetes—2021. *Diabetes Care* 44:Supplement_1, S200-S210. [[Crossref](#)]
603. 2021. Peculiarities of the clinical course of hypothyroidism during pregnancy. *Actual problems of modern medicine* :7. . [[Crossref](#)]
604. Hui-Hui Shen, Zhen-Zhen Lai, Hui-Li Yang, Jia-Wei Shi, Ming-Qing Li. 2021. Role of autoantibodies in infertility, miscarriage, and assisted reproductive technology outcomes. *Reproductive and Developmental Medicine*, ahead of print0. [[Crossref](#)]
605. Dheeman Futela, K Maheswari, Tejasvini Khanna. 2021. Fasting versus postprandial state: Impact on thyroid function testing. *Thyroid Research and Practice* 18:2, 61. [[Crossref](#)]
606. Mai Nouchi, Muneo Nakaya, Naoyuki Matsumoto, Hiroko Hojo, Wataru Kida, Yasuhiro Inayoshi, Hitoshi Iwamura. 2021. A case of aggressive follicular thyroid carcinoma in pregnancy. *JOURNAL OF JAPAN SOCIETY FOR HEAD AND NECK SURGERY* 31:2, 209-214. [[Crossref](#)]
607. Khulood Hussein. 2021. Hypothyroidism and Hypothyroxinemia in Pregnancy: An Overview in the Time of Coronavirus Disease 2019. *Pharmacophore* 12:2, 22-31. [[Crossref](#)]
608. Guangqin Zhou, Meiyang Zhou, Xuan Duan, Weihong Li. 2021. Glucocorticoid supplementation improves reproductive outcomes in infertile women with antithyroid autoimmunity undergoing ART. *Medicine* 100:16, e25554. [[Crossref](#)]
609. I-Ju Tu, Shih-Ming Huang, Ying-Ren Chen. 2021. Pregnancy exacerbated lymph node metastasis of well differentiated thyroid carcinoma. *Medicine* 100:51, e28264. [[Crossref](#)]
610. E. S. Shilova, N. V. Borovik, P. V. Popova, M. I. Yarmolinskaya. 2020. Diagnostic and treatment of subclinical hypothyroidism in pregnant and planning pregnancy patients: modern view of the problem. *Problems of Endocrinology* 66:6, 65-73. [[Crossref](#)]
611. Kaminskiy Viacheslav Volodymyrovych, Suslikova Lidiya Viktorivna, Serbeniuk Anastasia Valeriivna, Dercach Andriy Dmytrovych. 2020. THYROID DYSFUNCTION IN WOMEN OF REPRODUCTIVE AGE WHO PARTICIPATED IN HOSTILITIES AND SUFFERED CONTUSIONS. *Science Review* :8(35). . [[Crossref](#)]
612. Elizabeth O. Buschur, Stephen F. Thung. Thyroid Disorders 231-239. [[Crossref](#)]
613. Mustafa BAKIRCI, Fatih KILIÇ, Burak ERS AK, Hakan ARSLAN, İlker SELCUK. 2020. GEBELERDE İLK TRİMESTER TSH SEVİYELERİNİN FETAL DOĞUM AĞIRLIKLARINA VE MATERNAL HEMOGLOBİN DEĞERLERİ ÜZERİNE ETKİSİ. *Jinekoloji-Obstetrik ve Neonatoloji Tıp Dergisi* . [[Crossref](#)]
614. Tal Schiller, Arnon Agmon, Viviana Ostrovsky, Gabi Shefer, Hilla Knobler, Taiba Zornitzki. 2020. Moderate Iodine Deficiency Is Common in Pregnancy but Does Not Alter Maternal and Neonatal Thyroid Function Tests. *Frontiers in Endocrinology* 11. . [[Crossref](#)]
615. Ting-Ting Lin, Chen Zhang, Han-Qiu Zhang, Yu Wang, Lei Chen, Cindy-Lee Dennis, Hefeng Huang, Yan-Ting Wu. 2020. Thyroid Hormone Changes in Early Pregnancy Along With the COVID-19 Pandemic. *Frontiers in Endocrinology* 11. . [[Crossref](#)]
616. Nathalie Silva de Moraes, Débora Ayres Saraiva, Carolina Corcino, Tatiana Berbara, Annie Schtscherbyna, Karina Moreira, Mario Vaisman, Erik K. Alexander, Patrícia Teixeira. 2020. Consequences of Iodine Deficiency and Excess in Pregnancy and Neonatal Outcomes: A Prospective Cohort Study in Rio de Janeiro, Brazil. *Thyroid* 30:12, 1792-1801. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
617. Xi Yang, Yamei Yu, Chen Zhang, Yong Zhang, Zhirou Chen, Lise Dubois, He-Feng Huang, William D. Fraser, Jianxia Fan. 2020. The Association Between Isolated Maternal Hypothyroxinemia in Early Pregnancy and Preterm Birth. *Thyroid* 30:12, 1724-1731. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
618. Lei Zhao, Di Teng, Xiaoguang Shi, Yongze Li, Jianming Ba, Bing Chen, Jianling Du, Lanjie He, Xiaoyang Lai, Yanbo Li, Haiyi Chi, Eryuan Liao, Chao Liu, Libin Liu, Guijun Qin, Yingfen Qin, Huibiao Quan, Bingyin Shi, Hui Sun, Xulei Tang, Nanwei

- Tong, Guixia Wang, Jin-an Zhang, Youmin Wang, Yuanming Xue, Li Yan, Jing Yang, Lihui Yang, Yongli Yao, Zhen Ye, Qiao Zhang, Lihui Zhang, Jun Zhu, Mei Zhu, Zhongyan Shan, Weiping Teng. 2020. The Effect of Increased Iodine Intake on Serum Thyrotropin: A Cross-Sectional, Chinese Nationwide Study. *Thyroid* **30**:12, 1810-1819. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
619. M. Dolores Ollero, Juan Pablo Martínez, Javier Pineda, Marta Toni, Mercedes Espada, Emma Anda. 2020. Change over time in the iodine nutritional status of pregnant women from the Pamplona healthcare region. *Endocrinología, Diabetes y Nutrición (English ed.)* **67**:10, 643-649. [[Crossref](#)]
 620. M. Dolores Ollero, Juan Pablo Martínez, Javier Pineda, Marta Toni, Mercedes Espada, Emma Anda. 2020. Evolución del estado de nutrición de yodo en gestantes del area sanitaria de Pamplona. *Endocrinología, Diabetes y Nutrición* **67**:10, 643-649. [[Crossref](#)]
 621. Xing Wang, Yu Zhang, Huiwen Tan, Yu Bai, Liangxue Zhou, Fang Fang, Andrew Faramand, Weelic Chong, Yang Hai. 2020. Effect of levothyroxine on pregnancy outcomes in women with thyroid autoimmunity: a systematic review with meta-analysis of randomized controlled trials. *Fertility and Sterility* **114**:6, 1306-1314. [[Crossref](#)]
 622. Terry F. Davies, Stig Andersen, Rauf Latif, Yuji Nagayama, Giuseppe Barbesino, Maria Brito, Anja K. Eckstein, Alex Stagnaro-Green, George J. Kahaly. 2020. Graves' disease. *Nature Reviews Disease Primers* **6**:1. . [[Crossref](#)]
 623. Andréa Harumy de Lima Hirata, Luiz Antônio de Jesus Rocha, Valdelena Alessandra da Silva, Robson José de Almeida, Lucas dos Santos Bacigalupo, Patrícia Varela, Leonardo Martins, João Bosco Pesquero, Humberto Dellê, Cleber Pinto Camacho. 2020. Circulating RNA Transcriptome of Pregnant Women with TSH Just Above the Trimester-Specific Reference and its Correlation with the Hypertensive Phenotype. *Scientific Reports* **10**:1. . [[Crossref](#)]
 624. Bruno Ramalho de Carvalho, Andrea Prestes Nacul, Cristina Laguna Benetti-Pinto, Ana Carolina Japur de Sá Rosa-e-Silva, José Maria Soares Júnior, Gustavo Arantes Rosa Maciel, Edmund Chada Baracat. 2020. Reproductive Outcomes in Cases of Subclinical Hypothyroidism and Thyroid Autoimmunity: A Narrative Review. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics* **42**:12, 829-833. [[Crossref](#)]
 625. Masakazu Notsu, Mika Yamauchi, Toshitsugu Sugimoto, Keizo Kanasaki. 2020. A pregnant woman with an autonomously functioning thyroid nodule: a case report. *Gynecological Endocrinology* **36**:12, 1140-1143. [[Crossref](#)]
 626. Qiaohua He, Yujing Zhang, Wenyi Qiu, Jingjing Fan, Cuilian Zhang, Joanne Kwak-Kim. 2020. Does thyroid autoimmunity affect the reproductive outcome in women with thyroid autoimmunity undergoing assisted reproductive technology?. *American Journal of Reproductive Immunology* **84**:6. . [[Crossref](#)]
 627. Krzysztof C. Lewandowski, Karolina Garnysz, Wojciech Horzelski, Joanna Kawalec, Karolina Budzen, Mariusz Grzesiak, Andrzej Lewinski. 2020. Subclinical thyroid dysfunction in the first trimester of pregnancy: 'Disease' versus physiological (pulsatile) variation in TSH concentrations. *Clinical Endocrinology* **93**:6, 739-745. [[Crossref](#)]
 628. Ayla Coussa, Hayder A. Hasan, Thomas M. Barber. 2020. Effects of in vitro fertilization (IVF) therapies on metabolic, endocrine and inflammatory status in IVF-conceived pregnancy. *Clinical Endocrinology* **93**:6, 705-712. [[Crossref](#)]
 629. Suvi Turunen, Marja Väärasmäki, Anna-Maria Lahesmaa-Korpinen, Maarit K. Leinonen, Mika Gissler, Tuija Männistö, Eila Suvanto. 2020. Maternal hyperthyroidism and pregnancy outcomes: A population-based cohort study. *Clinical Endocrinology* **93**:6, 721-728. [[Crossref](#)]
 630. Biao Wang, Yajuan Xu, Miao Zhang, Jingzhe Zhang, Xiaofeng Hou, Jingjing Li, Yanjun Cai, Zongzong Sun, Yanjie Ban, Wentao Wang. 2020. Oral and intestinal microbial features in pregnant women with hypothyroidism and their correlations with pregnancy outcomes. *American Journal of Physiology-Endocrinology and Metabolism* **319**:6, E1044-E1052. [[Crossref](#)]
 631. Maike Huisinga, Lise Bertrand, Ronnie Chamanza, Isabelle Damiani, Jeff Engelhardt, Sabine Francke, Alexius Freyberger, Takanori Harada, Johannes Harleman, Wolfgang Kaufmann, Kevin Keane, Josef Köhrle, Barbara Lenz, M. Sue Marty, Stephanie Melching-Kollmuss, Xavier Palazzi, Gabriele Pohlmeier-Esch, Andreas Popp, Thomas J. Rosol, Volker Strauss, Hetty Van den Brink-Knol, Charles E. Wood, Midori Yoshida. 2020. Adversity Considerations for Thyroid Follicular Cell Hypertrophy and Hyperplasia in Nonclinical Toxicity Studies: Results From the 6th ESTP International Expert Workshop. *Toxicologic Pathology* **48**:8, 920-938. [[Crossref](#)]
 632. Akiko Fujishima, Akira Sato, Hiroshi Miura, Yuki Shimoda, Saeko Kameyama, Chika Ariake, Hiroyuki Adachi, Yuki Fukuoka, Yukihiko Terada. 2020. Fetal goiter identified in a pregnant woman with triiodothyronine-predominant graves' disease: a case report. *BMC Pregnancy and Childbirth* **20**:1. . [[Crossref](#)]
 633. Farnaz Sepasi, Tayebbeh Rashidian, Mehdi Shokri, Gholamreza Badfar, Fatemeh Kazemi, Milad Azami. 2020. Thyroid dysfunction in Iranian pregnant women: a systematic review and meta-analysis. *BMC Pregnancy and Childbirth* **20**:1. . [[Crossref](#)]
 634. Ning Yuan, Jianbin Sun, Zhi Li, Sanbao Chai, Xiaomei Zhang, Linong Ji. 2020. Relationship between anti-thyroid peroxidase antibody positivity and pregnancy-related and fetal outcomes in Euthyroid women: a single-center cohort study. *BMC Pregnancy and Childbirth* **20**:1. . [[Crossref](#)]

635. Yuchao Zhang, Wenbin Wu, Yanli Liu, Yichun Guan, Xingling Wang, Liting Jia. 2020. The impact of TSH levels on clinical outcomes 14 days after frozen-thawed embryo transfer. *BMC Pregnancy and Childbirth* **20**:1. . [\[Crossref\]](#)
636. Kalpana Mahadik, Payal Choudhary, P. K. Roy. 2020. Study of thyroid function in pregnancy, its feto-maternal outcome; a prospective observational study. *BMC Pregnancy and Childbirth* **20**:1. . [\[Crossref\]](#)
637. Juan Gui, Wangming Xu, Jie Zhang. 2020. Association between thyroid dysfunction and perinatal outcomes in women with gestational hypertension: a retrospective study. *BMC Pregnancy and Childbirth* **20**:1. . [\[Crossref\]](#)
638. Malgorzata Karbownik-Lewinska, Jan Stepniak, Magdalena Marcinkowska, Adrian Krygier, Andrzej Lewinski. 2020. High normal TSH is associated with lower mannan-binding lectin in women of childbearing age. *BMC Endocrine Disorders* **20**:1. . [\[Crossref\]](#)
639. Janos K. Aranyosi, Tamas Deli, Annamaria Erdei, Geza Toth, Attila Jakab, Mariann Fodor, Endre V. Nagy, Bernadett Ujhelyi. 2020. Unusual onset of thyroid associated orbitopathy during pregnancy: case report and review of literature. *BMC Endocrine Disorders* **20**:1. . [\[Crossref\]](#)
640. Sophie A. H. Boesen, Manhai Long, Maria Wielsøe, Vicente Mustieles, Mariana F. Fernandez, Eva C. Bonefeld-Jørgensen. 2020. Exposure to Perfluoroalkyl acids and foetal and maternal thyroid status: a review. *Environmental Health* **19**:1. . [\[Crossref\]](#)
641. Ashley Venables, Wilbert Wong, Mandy Way, Hayden Anthony Homer. 2020. Thyroid autoimmunity and IVF/ICSI outcomes in euthyroid women: a systematic review and meta-analysis. *Reproductive Biology and Endocrinology* **18**:1. . [\[Crossref\]](#)
642. Stine Linding Andersen, Stig Andersen. 2020. Antithyroid drugs and birth defects. *Thyroid Research* **13**:1. . [\[Crossref\]](#)
643. Subhodip Pramanik, Pradip Mukhopadhyay, Sujoy Ghosh. 2020. Total T4 rise in pregnancy: a relook?. *Thyroid Research* **13**:1. . [\[Crossref\]](#)
644. Leonidas H. Duntas. 2020. Selenium and at-risk pregnancy: challenges and controversies. *Thyroid Research* **13**:1. . [\[Crossref\]](#)
645. Shani R. Rosen, Yaniv S. Ovadia, Eyal Y. Anteby, Shlomo Fytlovich, Dorit Aharoni, Doron Zamir, Dov Gefel, Simon Shenhav. 2020. Low intake of iodized salt and iodine containing supplements among pregnant women with apparently insufficient iodine status - time to change policy?. *Israel Journal of Health Policy Research* **9**:1. . [\[Crossref\]](#)
646. Vehbi Yavuz Tokgoz, Berker Isim, Ahmet Basar Tekin. 2020. The impact of thyroid autoantibodies on the cycle outcome and embryo quality in women undergoing intracytoplasmic sperm injection. *Middle East Fertility Society Journal* **25**:1. . [\[Crossref\]](#)
647. Grace Mengqin Ge, Miriam T Y Leung, Kenneth K C Man, Wing Cheong Leung, Patrick Ip, Gloria H Y Li, Ian C K Wong, Annie W C Kung, Ching-Lung Cheung. 2020. Maternal Thyroid Dysfunction During Pregnancy and the Risk of Adverse Outcomes in the Offspring: A Systematic Review and Meta-Analysis. *The Journal of Clinical Endocrinology & Metabolism* **105**:12, 3821-3841. [\[Crossref\]](#)
648. George J Kahaly. 2020. Management of Graves Thyroidal and Extrathyroidal Disease: An Update. *The Journal of Clinical Endocrinology & Metabolism* **105**:12, 3704-3720. [\[Crossref\]](#)
649. Kosuke Inoue, Atsushi Goto, Takehiro Sugiyama, Cecilia Høst Ramlau-Hansen, Zeyan Liew. 2020. The Confounder-Mediator Dilemma: Should We Control for Obesity to Estimate the Effect of Perfluoroalkyl Substances on Health Outcomes?. *Toxics* **8**:4, 125. [\[Crossref\]](#)
650. Jae Hoon Chung. 2020. Recent Issues Related to Thyroid Disease in Pregnancy. *International Journal of Thyroidology* **13**:2, 85-94. [\[Crossref\]](#)
651. Fadeev Fadeev. 2020. Review of European Thyroid Association Guideline (2018) for the Manage-ment of Graves' Hyperthyroidism. *Clinical and experimental thyroidology* **16**:1, 4-20. [\[Crossref\]](#)
652. Ina Kusriani, Jessica Farebrother, Donny Kristanto Mulyantoro. 2020. Adequately iodized salt is an important strategy to prevent iodine insufficiency in pregnant women living in Central Java, Indonesia. *PLOS ONE* **15**:11, e0242575. [\[Crossref\]](#)
653. Yuchao Zhang, Wenbin Wu, Yanli Liu, Yichun Guan, Xingling Wang, Liting Jia. 2020. The Impact of Preconception TSH on the Reproductive Outcomes of Infertile Women Undergoing the First Fresh D3 Embryo Transfer Cycle. *International Journal of Endocrinology* **2020**, 1-7. [\[Crossref\]](#)
654. Elizabeth N. Pearce. 2020. The Proportion of In-Range TSH Values Does Not Differ in Users of Desiccated Thyroid Compared to Levothyroxine. *Clinical Thyroidology* **32**:11, 505-507. [\[Citation\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
655. Na Li, Jiaying Yang, Xi Chen, Jingjing Huang, Mengyu Lai, Fang Fang, Liping Gu, Yu-fan Wang, Yong-de Peng. 2020. Postpartum Follow-Up of Patients with Subclinical Hypothyroidism During Pregnancy. *Thyroid* **30**:11, 1566-1573. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#) [\[Supplementary Material\]](#)
656. Zixiao Wang, Chenyang Li, Ying Teng, Yunping Guan, Li Zhang, Xi Jia, Dandan Cui, Jing Li, Haixia Guan. 2020. The Effect of Iodine-Containing Vitamin Supplementation During Pregnancy on Thyroid Function in Late Pregnancy and Postpartum Depression in an Iodine-Sufficient Area. *Biological Trace Element Research* **198**:1, 1-7. [\[Crossref\]](#)

657. Urs D. Lichtenauer, Markus Reincke. 2020. Latente Hypothyreose - (k)ein Problem?. *MMW - Fortschritte der Medizin* **162**:S3, 76-83. [[Crossref](#)]
658. Alberto López García-Franco, José Antonio Baeyens Fernández, Emilia Bailón Muñoz, M. José Iglesias Piñeiro, Amparo Ortega del Moral, Pablo Alonso Coello, Cristina Ruiz Cabello, Jacinta Landa Goñi, Lorenzo Arribas Mir. 2020. Actividades preventivas en la mujer. Actualización PAPPS 2020. *Atención Primaria* **52**, 125-148. [[Crossref](#)]
659. Fernanda de Oliveira Lopes, Fernanda Valente Mendes Soares, Danielle Aparecida da Silva, Maria Elisabeth Lopes Moreira. 2020. Do Thyroid Diseases during Pregnancy and Lactation Affect the Nutritional Composition of Human Milk?. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics* **42**:11, 752-758. [[Crossref](#)]
660. Stine Linding Andersen, Peter Astrup Christensen, Louise Knøsgaard, Stig Andersen, Aase Handberg, Annebirthe Bo Hansen, Peter Vestergaard. 2020. Classification of Thyroid Dysfunction in Pregnant Women Differs by Analytical Method and Type of Thyroid Function Test. *The Journal of Clinical Endocrinology & Metabolism* **105**:11, e4012-e4022. [[Crossref](#)]
661. Ursula G. Sauer, Alex Asiimwe, Philip A. Botham, Alex Charlton, Nina Hallmark, Sylvia Jacobi, Sue Marty, Stephanie Melching-Kollmuss, Joana A. Palha, Volker Strauss, Bennard van Ravenzwaay, Gerard Swaen. 2020. Toward a science-based testing strategy to identify maternal thyroid hormone imbalance and neurodevelopmental effects in the progeny – part I: which parameters from human studies are most relevant for toxicological assessments?. *Critical Reviews in Toxicology* **50**:9, 740-763. [[Crossref](#)]
662. Liangmiao Chen, Hong Yang, Enling Ye, Zhenzhen Lin, Mengmeng Peng, Hai Lin, Lechu Yu, Zhuhua Cai, Xuemian Lu. 2020. Insignificant Effect of Isolated Hypothyroxinemia on Pregnancy Outcomes During the First and Second Trimester of Pregnancy. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
663. A. Z. Bulgakova, E. A. Fazlyeva, G. A. Galieva, R. A. Izmailova. 2020. Experience in organizing screening of thyroid pathology during pregnancy in the region of iodine deficiency. *Meditinskiy sovet = Medical Council* :13, 58-64. [[Crossref](#)]
664. Mei-Fang Li, Li Ma, Qi-Ming Feng, Yue Zhu, Tian-Pei Yu, Jiang-Feng Ke, Zhi-Hui Zhang, Yun Liu, Lian-Xi Li. 2020. Effects of Maternal Subclinical Hypothyroidism in Early Pregnancy Diagnosed by Different Criteria on Adverse Perinatal Outcomes in Chinese Women With Negative TPOAb. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
665. Nicole J. E. Verhagen, Sueppong Gowachirapant, Pattanee Winichagoon, Maria Andersson, Alida Melse-Boonstra, Michael B. Zimmermann. 2020. Iodine Supplementation in Mildly Iodine-Deficient Pregnant Women Does Not Improve Maternal Thyroid Function or Child Development: A Secondary Analysis of a Randomized Controlled Trial. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
666. Sun Y. Lee. 2020. No Differences in Neurodevelopmental Scores of Children Born to Hypothyroid Women Treated with Levothyroxine During Preconception Versus Early Pregnancy. *Clinical Thyroidology* **32**:10, 464-467. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
667. Alan P. Farwell. 2020. Commentary on “Long Work Hours Are Associated with Hypothyroidism: A Cross-Sectional Study with Population-Representative Data” by Lee et al. *Thyroid* **30**:10, 1395-1396. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
668. Tanja Groten. 2020. 30-jährige schwangere Patientin mit Diabetes mellitus Typ 1. *Der Gynäkologe* **53**:S1, 58-65. [[Crossref](#)]
669. Satoru Funaki, Nagayoshi Umehara, Hidetoshi Mezawa, Masamitsu Kurakazu, Sachio Matsushima, Naoko Arata, Aikou Okamoto, Haruhiko Sago. 2020. Ultrasonographic assessment of fetal thyroid in Japan: thyroid circumference and distal femoral and proximal tibial ossification. *Journal of Medical Ultrasonics* **47**:4, 603-608. [[Crossref](#)]
670. Luciana Sant’Ana Leone de Souza, Renata de Oliveira Campos, Vitor dos Santos Alves, Taíse Lima Oliveira Cerqueira, Thiago Magalhães da Silva, Leonardo Sena Gomes Teixeira, Alina Coutinho Rodrigues Feitosa, Crésio de Aragão Dantas Alves, Helton Estrela Ramos. 2020. Hypertension and Salt-Restrictive Diet Promotes Low Urinary Iodine Concentration in High-Risk Pregnant Women: Results from a Cross-Sectional Study Conducted After Salt Iodination Reduction in Brazil. *Biological Trace Element Research* **197**:2, 445-453. [[Crossref](#)]
671. T. Álvarez Fernández, M.M. González Vilanova, M.J. Bal Alvarado, M.C. Sánchez Pérez-Mel, L. Máiz Suárez. 2020. Valores de referencia de hormonas tiroideas en la población de gestantes en Lugo. *Clínica e Investigación en Ginecología y Obstetrícia* **47**:4, 143-148. [[Crossref](#)]
672. Lois Donovan, Heather Cockwell, Niamh Tallon, Jennifer Yamamoto. 2020. Opinion de comité no 407 : Pathologie thyroïdienne et infertilité. *Journal of Obstetrics and Gynaecology Canada* **42**:10, 1283-1286. [[Crossref](#)]
673. Lois Donovan, Heather Cockwell, Niamh Tallon, Jennifer Yamamoto. 2020. Committee Opinion No. 407: Thyroid Disease and Infertility. *Journal of Obstetrics and Gynaecology Canada* **42**:10, 1279-1282. [[Crossref](#)]
674. Bevin E. Blake, Suzanne E. Fenton. 2020. Early life exposure to per- and polyfluoroalkyl substances (PFAS) and latent health outcomes: A review including the placenta as a target tissue and possible driver of peri- and postnatal effects. *Toxicology* **443**, 152565. [[Crossref](#)]

675. Aino Lintula, Leea Keski-Nisula, Heidi Sahlman. 2020. Hypothyroidism and the increased risk of preeclampsia – interpretative factors?. *Hypertension in Pregnancy* 39:4, 411-417. [[Crossref](#)]
676. Margreet Meems, Lianne Hulsbosch, Madelon Riem, Christina Meyers, Tila Pronk, Maarten Broeren, Karin Nabbe, Guid Oei, Stefan Bogaerts, Victor Pop. 2020. The Brabant study: design of a large prospective perinatal cohort study among pregnant women investigating obstetric outcome from a biopsychosocial perspective. *BMJ Open* 10:10, e038891. [[Crossref](#)]
677. Megan J. Ritter, Suruchi Gupta, James V. Hennessey. 2020. Alternative routes of levothyroxine administration for hypothyroidism. *Current Opinion in Endocrinology, Diabetes & Obesity* 27:5, 318-322. [[Crossref](#)]
678. Andrea Padoan, Aldo Clerico, Martina Zaninotto, Tommaso Trenti, Renato Tozzoli, Rosalia Aloe, Antonio Alfano, Sara Rizzardi, Ruggero Dittadi, Marco Migliardi, Marcello Bagnasco, Mario Plebani. 2020. Percentile transformation and recalibration functions allow harmonization of thyroid-stimulating hormone (TSH) immunoassay results. *Clinical Chemistry and Laboratory Medicine (CCLM)* 58:10, 1663-1672. [[Crossref](#)]
679. Shahin Safian, Farzaneh Esna-Ashari, Shiva Borzouei. 2020. Thyroid Dysfunction in Pregnant Women with Gestational Diabetes Mellitus. *Current Diabetes Reviews* 16:8, 895-899. [[Crossref](#)]
680. Mariza Kampouri, Katerina Margetaki, Katerina Koutra, Andriani Kyriklaki, Polyxeni Karakosta, Despoina Anousaki, Georgia Chalkiadaki, Marina Vafeiadi, Manolis Kogevas, Leda Chatzi. 2020. Maternal mild thyroid dysfunction and offspring cognitive and motor development from infancy to childhood: the Rhea mother-child cohort study in Crete, Greece. *Journal of Epidemiology and Community Health* 3, jech-2019-213309. [[Crossref](#)]
681. A. S. Loktionova, I. A. Ilvayskaya. 2020. Basal level of luteinizing hormone as a key marker of the idiopathic central hypogonadism in women. *Almanac of Clinical Medicine* 48. . [[Crossref](#)]
682. Kamilla G. Eriksen, Maria Andersson, Sandra Hunziker, Michael B. Zimmermann, Sophie E. Moore. 2020. Effects of an Iodine-Containing Prenatal Multiple Micronutrient on Maternal and Infant Iodine Status and Thyroid Function: A Randomized Trial in The Gambia. *Thyroid* 30:9, 1355-1365. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
683. Alfred N. Romeo, Sarah G. Običan. 2020. Teratogen update: Antithyroid medications. *Birth Defects Research* 112:15, 1150-1170. [[Crossref](#)]
684. Sofia Manousou, Maria Andersson, Robert Eggertsen, Sandra Hunziker, Lena Hulthén, Helena Filipsson Nyström. 2020. Iodine deficiency in pregnant women in Sweden: a national cross-sectional study. *European Journal of Nutrition* 59:6, 2535-2545. [[Crossref](#)]
685. Stefan Pilz, Verena Theiler-Schwetz, Oliver Malle, Eva Steinberger, Marlene Pandis, Elisabeth Lerchbaum, Christian Trummer. 2020. Schilddrüse: Fertilität, Schwangerschaft und Laktation. *Journal für Klinische Endokrinologie und Stoffwechsel* 13:3, 106-114. [[Crossref](#)]
686. Aleksandra Mikołajczak, Maria Katarzyna Borszewska-Kornacka, Ewa Romejko-Wolniewicz, Renata Bokiniec. 2020. Comparison of the offspring ultrasound thyroid volume in hypothyroid mothers treated with different levothyroxine doses: A cohort study. *Advances in Medical Sciences* 65:2, 332-337. [[Crossref](#)]
687. T. Wasilewski, M. Łukaszewicz-Zajac, J. Wasilewska, B. Mroczko. 2020. Biochemistry of infertility. *Clinica Chimica Acta* 508, 185-190. [[Crossref](#)]
688. Stefano Luisi, Matteo Giorgi, Silvia Riggi, Gabriele Messina, Filiberto Maria Severi. 2020. Neonatal outcome in pregnancy hypothyroid women. *Gynecological Endocrinology* 36:9, 772-775. [[Crossref](#)]
689. Maja Cigrovski Berković, Davorka Herman Mahečić, Jelena Marinković Radošević, Mateja Strinović Morić, Ines Bilić-Ćurčić. 2020. Hypothyroidism and pregnancy: still a controversial issue. *Gynecological Endocrinology* 36:9, 776-780. [[Crossref](#)]
690. Louise Knøsgaard, Stig Andersen, Annebirthe Bo Hansen, Peter Vestergaard, Stine Linding Andersen. 2020. Thyroid function abnormalities and thyroid autoantibodies in Danish pregnant women. *Clinical Endocrinology* 93:3, 329-338. [[Crossref](#)]
691. S. O. Shurpyak. 2020. The substantiation of differentiated therapy of the combined disgormonal pathology of reproductive organs in women of reproductive age with comorbid conditions. *Reproductive health of woman* 2:2, 5-10. [[Crossref](#)]
692. Anastasia Linardi, Ioannis Kakoulidis, Ioannis Ilias, Aikaterini Michou, Athina Pappa, Evangelia Venaki, Eftychia Koukkou. 2020. Postpartum Follow-Up of Women Who Developed Subclinical Hypothyroidism during Pregnancy. *Medical Sciences* 8:3, 29. [[Crossref](#)]
693. Mendes AB, Seabra FCA, de Sousa Bechtluft IM, Costa JA, Ajeje LS, Guimarães MM, Rodrigues VO, de Paiva AR, Silva AC, Panconi CR, de Souza HD, Zimmermann JB. 2020. Urinary Iodine Concentration in 24-Hour Urine of Pregnant Women and Its Association with Food and Salt Intake. *Asploro Journal of Biomedical and Clinical Case Reports* 3:2, 119-127. [[Crossref](#)]

694. Yuchao Zhang, Wenbin Wu, Yanli Liu, Yichun Guan, Xingling Wang, Liting Jia. 2020. High-Normal Preconception TSH Levels Have No Adverse Effects on Reproductive Outcomes in Infertile Women Undergoing the First Single Fresh D5 Blastocyst Transfer. *International Journal of Endocrinology* **2020**, 1-8. [[Crossref](#)]
695. Shigeo Iijima. 2020. Pitfalls in the assessment of gestational transient thyrotoxicosis. *Gynecological Endocrinology* **36**:8, 662-667. [[Crossref](#)]
696. Fabienne Delay, Vincent Dochez, Florence Biquard, Marie-Thérèse Cheve, Philippe Gillard, C. J. Arthuis, Norbert Winer. 2020. Management of fetal goiters: 6-year retrospective observational study in three prenatal diagnosis and treatment centers of the Pays De Loire Perinatal Network. *The Journal of Maternal-Fetal & Neonatal Medicine* **33**:15, 2561-2569. [[Crossref](#)]
697. Sun Y. Lee. 2020. High Prevalence of Subclinical Hypothyroidism and TPO Antibody Positivity Among Women with a History of Miscarriage or Infertility. *Clinical Thyroidology* **32**:8, 368-371. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
698. Tim I.M. Korevaar. 2020. Levothyroxine Overtreatment During Pregnancy May Decrease Offspring Thyroid Gland Volume. *Clinical Thyroidology* **32**:8, 372-374. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
699. Kris Poppe, Candice Autin, Flora Veltri, Georgiana Sitoris, Pierre Kleynen, Jean-Philippe Praet, Serge Rozenberg. 2020. Thyroid Disorders and In Vitro Outcomes of Assisted Reproductive Technology: An Unfortunate Combination?. *Thyroid* **30**:8, 1177-1185. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
700. Sandra A. Lowe. 2020. Ionizing radiation for maternal medical indications. *Prenatal Diagnosis* **40**:9, 1150-1155. [[Crossref](#)]
701. Sawyer Huget-Penner, Denice S. Feig. 2020. Maternal thyroid disease and its effects on the fetus and perinatal outcomes. *Prenatal Diagnosis* **40**:9, 1077-1084. [[Crossref](#)]
702. Jamil M. Kazma, John van den Anker, Karel Allegaert, André Dallmann, Homa K. Ahmadzia. 2020. Anatomical and physiological alterations of pregnancy. *Journal of Pharmacokinetics and Pharmacodynamics* **47**:4, 271-285. [[Crossref](#)]
703. Susan Ogg, James L. Klosky, Wassim Chemaitilly, Deo Kumar Srivastava, Mingjuan Wang, Ginger Carney, Rohit Ojha, Leslie L. Robison, Cheryl L. Cox, Melissa M. Hudson. 2020. Breastfeeding practices among childhood cancer survivors. *Journal of Cancer Survivorship* **14**:4, 586-599. [[Crossref](#)]
704. Nazarena Betella, Walter Vena, Emilia Biamonte, Elisabetta Lavezzi, Andrea Lania. 2020. Gli adenomi ipofisari in gravidanza. *L'Endocrinologo* **21**:4, 246-251. [[Crossref](#)]
705. Elisabetta Torlone, Maria Angela Sculli, Matteo Bonomo, Antonino Di Benedetto, Graziano Di Cianni, Camilla Festa, Gloria Formoso, Annunziata Lapolla, Domenico Mannino, Angela Napoli, Marina Scavini, Elena Succurro, Ester Vitacolonna, Laura Sciacca. 2020. Recommendations and management of hyperglycaemia in pregnancy during COVID-19 pandemic in Italy. *Diabetes Research and Clinical Practice* **166**, 108345. [[Crossref](#)]
706. Mariana Dineva, Harry Fishpool, Margaret P Rayman, Jeewaka Mendis, Sarah C Bath. 2020. Systematic review and meta-analysis of the effects of iodine supplementation on thyroid function and child neurodevelopment in mildly-to-moderately iodine-deficient pregnant women. *The American Journal of Clinical Nutrition* **112**:2, 389-412. [[Crossref](#)]
707. Michael B Zimmermann. 2020. Iodine supplements for mildly iodine-deficient pregnant women: are they worthwhile?. *The American Journal of Clinical Nutrition* **112**:2, 247-248. [[Crossref](#)]
708. Jon T. Andersen, Tina Bergmann Futtrup. 2020. Drugs during lactation. *Adverse Drug Reaction Bulletin* **323**:1, 1251-1254. [[Crossref](#)]
709. Jane Michele Peart, Robert Sim. 2020. Lipiodol hysterosalpingogram: A modified HSG technique to minimize risks associated with lipiodol use. *Journal of Medical Imaging and Radiation Oncology* **64**:4, 516-521. [[Crossref](#)]
710. Louisa M Churcher, Julia A Wood, Kim A Ramjan, Anna C Gill. 2020. Reducing unnecessary neonatal testing in infants of mothers with thyroid disease. *Journal of Paediatrics and Child Health* **56**:8, 1185-1188. [[Crossref](#)]
711. Rima K Dhillon-Smith, Aurelio Tobias, Paul P Smith, Lee J Middleton, Kirandeep K Sunner, Krystyna Baker, Samantha Farrell-Carver, Ruth Bender-Atik, Rina Agrawal, Kalsang Bhatia, Justin J Chu, Edmond Edi-Osagie, Ayman Ewies, Tarek Ghobara, Pratima Gupta, Davor Jurkovic, Yacoub Khalaf, Khashia Mulbagal, Natalie Nunes, Caroline Overton, Siobhan Quenby, Raj Rai, Nick Raine-Fenning, Lynne Robinson, Jackie Ross, Andrew Sizer, Rachel Small, Martyn Underwood, Mark D Kilby, Jane Daniels, Shakila Thangaratinam, Shiao Chan, Kristien Boelaert, Arri Coomarasamy. 2020. The Prevalence of Thyroid Dysfunction and Autoimmunity in Women With History of Miscarriage or Subfertility. *The Journal of Clinical Endocrinology & Metabolism* **105**:8, 2667-2677. [[Crossref](#)]
712. Anca Marina Ciobanu, Andreea Elena Dumitru, Nicolae Gica, Radu Botezatu, Gheorghe Peltecu, Anca Maria Panaitescu. 2020. Benefits and Risks of IgG Transplacental Transfer. *Diagnostics* **10**:8, 583. [[Crossref](#)]

713. Simona Censi, Jacopo Manso, Susi Barollo, Alberto Mondin, Loris Bertazza, Massimo De Marchi, Caterina Mian, on behalf of the Food and Nutrition Hygiene Services (SIAN). 2020. Changing Dietary Habits in Veneto Region over Two Decades: Still a Long Road to Go to Reach an Iodine-Sufficient Status. *Nutrients* **12**:8, 2399. [[Crossref](#)]
714. Hannah S. Clausen, Ninna H. Ebdrup, Ida M. Barsøe, Julie Lyngsø, Jörg Schullehner, Cecilia H. Ramlau-Hansen, Bjørn Bay, Ulla B. Knudsen. 2020. Association between Drinking Water Nitrate and Adverse Reproductive Outcomes: A Systematic PRISMA Review. *Water* **12**:8, 2287. [[Crossref](#)]
715. Sangyoong Park, Soron Choi, Joonho Jeong, Jeongho Kim. 2020. Spinal anesthesia for urgent Cesarean section in a patient with uncontrolled hyperthyroidism due to Graves' disease - A case report -. *Anesthesia and Pain Medicine* **15**:3, 319-324. [[Crossref](#)]
716. Juan José CORRALES-HERNÁNDEZ, Ana Isabel SÁNCHEZ-MARCOS, José María RECIO-CÓRDOVA, Rosa Ana IGLESIAS-LÓPEZ, María Teresa MORIES ALVÁREZ. 2020. Tratamiento médico del hipertiroidismo. *Revista ORL* **11**:3, 8. [[Crossref](#)]
717. Guocheng Wang, Guojun Zhang. 2020. The Diverse Upper Reference Limits of Serum Thyroid-Stimulating Hormone on the Same Platform for Pregnant Women in China. *Laboratory Medicine* **51**:4, 416-422. [[Crossref](#)]
718. Michael T. McDermott. 2020. Hypothyroidism. *Annals of Internal Medicine* **173**:1, ITC1-ITC16. [[Crossref](#)]
719. Elizabeth N. Pearce. 2020. A Comparison of ATA and Updated ACOG Guidelines for Thyroid Disease in Pregnancy. *Clinical Thyroidology* **32**:7, 317-320. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
720. Shuhei So, Fumiko Tawara. 2020. Risk factors of subclinical hypothyroidism and the potential contribution to miscarriage: A review. *Reproductive Medicine and Biology* **19**:3, 232-242. [[Crossref](#)]
721. Elizabeth C. Ailes, John Zimmerman, Jennifer N. Lind, Fanghui Fan, Kun Shi, Jennita Reefhuis, Cheryl S. Broussard, Meghan T. Frey, Janet D. Cragan, Emily E. Petersen, Kara D. Polen, Margaret A. Honein, Suzanne M. Gilboa. 2020. Using Supervised Learning Methods to Develop a List of Prescription Medications of Greatest Concern during Pregnancy. *Maternal and Child Health Journal* **24**:7, 901-910. [[Crossref](#)]
722. Ali S. Alzahrani, Mourad Al Mourad, Kevin Hafez, Abdulrahman M. Almaghamsy, Fahad Abdulrahman Alamri, Nasser R. Al Juhani, Alhussien Sagr Alhazmi, Mohammad Yahya Saeedi, Saud Alsefri, Musa Daif Allah Alzahrani, Nadia Al Ali, Wiam I. Hussein, Mohamed Ismail, Ahmed Adel, Hisham El Bahtimy, Eslam Abdelhamid. 2020. Diagnosis and Management of Hypothyroidism in Gulf Cooperation Council (GCC) Countries. *Advances in Therapy* **37**:7, 3097-3111. [[Crossref](#)]
723. Alex Stagnaro-Green, Allan Dong, Mary D. Stephenson. 2020. Universal screening for thyroid disease during pregnancy should be performed. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101320. [[Crossref](#)]
724. Maria Papaleontiou, Megan R. Haymart. 2020. Thyroid nodules and cancer during pregnancy, post-partum and preconception planning: Addressing the uncertainties and challenges. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101363. [[Crossref](#)]
725. David Unuane, Brigitte Velkeniers. 2020. Impact of thyroid disease on fertility and assisted conception. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101378. [[Crossref](#)]
726. Stine Linding Andersen, Louise Knøsgaard. 2020. Management of thyrotoxicosis during pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101414. [[Crossref](#)]
727. A.H. van der Spek, P.H. Bisschop. 2020. Universal screening for thyroid disease SHOULD NOT be recommended before and during pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101429. [[Crossref](#)]
728. Eduardo Rodriguez-Diaz, Elizabeth N. Pearce. 2020. Iodine status and supplementation before, during, and after pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101430. [[Crossref](#)]
729. W. Edward Visser, Robin P. Peeters. 2020. Interpretation of thyroid function tests during pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101431. [[Crossref](#)]
730. Peter N. Taylor, Ilaria Muller, Melanie Nana, Ines Velasco, John H. Lazarus. 2020. Indications for treatment of subclinical hypothyroidism and isolated hypothyroxinaemia in pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101436. [[Crossref](#)]
731. A.S. Paul van Trotsenburg. 2020. Management of neonates born to mothers with thyroid dysfunction, and points for attention during pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101437. [[Crossref](#)]
732. Nobuyuki Amino, Naoko Arata. 2020. Thyroid dysfunction following pregnancy and implications for breastfeeding. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101438. [[Crossref](#)]
733. Sarah Weiling Li, Shiao-Yng Chan. 2020. Management of overt hypothyroidism during pregnancy. *Best Practice & Research Clinical Endocrinology & Metabolism* **34**:4, 101439. [[Crossref](#)]

734. Karen Gabriela Bautista-Orduno, Edgar Gerardo Dorsey-Trevino, Jose Gerardo Gonzalez-Gonzalez, Dalia A. Castillo-Gonzalez, Mariana Garcia-Leal, Karina Raygoza-Cortez, Michael R. Gionfriddo, Rene Rodriguez-Gutierrez. 2020. American thyroid association guidelines are inconsistent with Grading of Recommendations Assessment, Development, and Evaluations—A meta-epidemiologic study. *Journal of Clinical Epidemiology* **123**, 180-188.e2. [[Crossref](#)]
735. Vanessa Y. De La Rosa, Joseph Hoover, Ruofei Du, Elizabeth Yakes Jimenez, Debra MacKenzie, Johnnye Lewis. 2020. Diet quality among pregnant women in the Navajo Birth Cohort Study. *Maternal & Child Nutrition* **16**:3. . [[Crossref](#)]
736. Iliara Muller, Peter N Taylor, Rhian M Daniel, Charlotte Hales, Anna Scholz, Toby Candler, Rebecca J Pettit, William D Evans, Dionne Shillabeer, Mohd S Draman, Colin M Dayan, Hiu K C Tang, Onyebuchi Okosieme, John W Gregory, John H Lazarus, D Aled Rees, Marian E Ludgate. 2020. CATS II Long-term Anthropometric and Metabolic Effects of Maternal Sub-optimal Thyroid Function in Offspring and Mothers. *The Journal of Clinical Endocrinology & Metabolism* **105**:7, 2150-2161. [[Crossref](#)]
737. Mariacarla Moleti, Maria Di Mauro, Angela Alibrandi, Roberto Vita, Salvatore Benvenga, Francesco Vermiglio. 2020. Postpartum Thyroiditis in Women With Euthyroid and Hypothyroid Hashimoto's Thyroiditis Antedating Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* **105**:7, e2421-e2428. [[Crossref](#)]
738. Elizabeth N Pearce. 2020. Thyroid Autoimmunity Is Associated With Postpartum Thyroiditis Risk, Even in Women With Treated Hypothyroidism. *The Journal of Clinical Endocrinology & Metabolism* **105**:7. . [[Crossref](#)]
739. Anna Karen Gárate-Escamilla, Edelmiro Garza-Padilla, Agustín Carvajal Rivera, Celina Salas-Castro, Emmanuel Andrés, Amir Hajjam El Hassani. 2020. Cluster Analysis: A New Approach for Identification of Underlying Risk Factors and Demographic Features of First Trimester Pregnancy Women. *Journal of Clinical Medicine* **9**:7, 2247. [[Crossref](#)]
740. Christopher Spoke, Christopher Martin. 2020. Maternal Graves Disease and Abnormal CYP2D6 Genotype with Fetal Hyperthyroidism. *AACE Clinical Case Reports* **6**:4, e161-e164. [[Crossref](#)]
741. Hengameh Abdi, Atieh Amouzegar. 2020. Management of Graves' Hyperthyroidism: More Than a Century of Progression. *International Journal of Endocrinology and Metabolism* **18**:Suppl. . [[Crossref](#)]
742. Freddy J. K. Toloza, Hooman Motahari, Spyridoula Maraka. 2020. Consequences of Severe Iodine Deficiency in Pregnancy: Evidence in Humans. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
743. Dalia A. Castillo-Gonzalez, Edgar G. Dorsey-Trevino, Jose G. Gonzalez-Gonzalez, Mariana Garcia-Leal, Karen G. Bautista-Orduño, Karina Raygoza, Michael R. Gionfriddo, Naykky M. Singh Ospina, Rene Rodriguez-Gutierrez. 2020. A deeper analysis in thyroid research: A meta-epidemiological study of the American Thyroid Association clinical guidelines. *PLOS ONE* **15**:6, e0234297. [[Crossref](#)]
744. Yuanyuan Chen, Fengxia Xue. 2020. The impact of gestational hypothyroxinemia on the cognitive and motor development of offspring. *The Journal of Maternal-Fetal & Neonatal Medicine* **33**:11, 1940-1945. [[Crossref](#)]
745. Haixia Guan. 2020. Maintaining Euthyroidism Without Levothyroxine Initiation or Dose Adjustment Increases Postpartum Thyroiditis Risk in Women with Hashimoto's Thyroiditis. *Clinical Thyroidology* **32**:6, 263-265. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
746. Tim I.M. Korevaar. 2020. Do Women with Gestational Subclinical Hypothyroidism Require Postpartum Monitoring?. *Clinical Thyroidology* **32**:6, 266-269. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
747. Shashank R. Joshi. 2020. Dynamic Live Guidelines Are Needed for Thyroid Disease in Pregnancy. *Clinical Thyroidology* **32**:6, 270-272. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
748. Michael C. Topf, Jared A. Shenson, F. Christopher Holsinger, Samuel H. Wald, Lisa J. Cianfichi, Eben L. Rosenthal, John B. Sunwoo. 2020. Framework for prioritizing head and neck surgery during the COVID -19 pandemic. *Head & Neck* **42**:6, 1159-1167. [[Crossref](#)]
749. Jacob Young, Calan Savoy, Ian Colman, Mark Ferro, Ryan J. Van Lieshout. 2020. Psychiatric Disorders in the Adolescent Offspring of Mothers with Thyroid Problems During Pregnancy. *Child Psychiatry & Human Development* **51**:3, 461-470. [[Crossref](#)]
750. Ayla Coussa, Hayder A. Hasan, Thomas M. Barber. 2020. Impact of contraception and IVF hormones on metabolic, endocrine, and inflammatory status. *Journal of Assisted Reproduction and Genetics* **37**:6, 1267-1272. [[Crossref](#)]
751. Miriam Promintzer-Schifferl, Michael Krebs. 2020. Schilddrüse und Schwangerschaft: Schein und Sein. *Journal für Klinische Endokrinologie und Stoffwechsel* **13**:2, 59-64. [[Crossref](#)]
752. Arash Derakhshan, Robin P Peeters, Peter N Taylor, Sofie Bliddal, David M Carty, Margreet Meems, Bijay Vaidya, Liangmiao Chen, Bridget A Knight, Farkhanda Ghafoor, Polina V Popova, Lorena Mosso, Emily Oken, Eila Suvanto, Aya Hisada, Jun Yoshinaga, Suzanne J Brown, Judit Bassols, Juha Auvinen, Wichor M Bramer, Abel López-Bermejo, Colin M Dayan, Robert French, Laura Boucai, Marina Vafeiadi, Elena N Grineva, Victor J M Pop, Tanja G Vrijkotte, Leda Chatzi, Jordi Sunyer, Ana Jiménez-Zabala, Isolina Riaño, Marisa Rebagliato, Xuemian Lu, Amna Pirzada, Tuija Männistö, Christian Delles, Ulla Feldt-Rasmussen, Erik K Alexander, Scott M Nelson, Loyal Chaker, Elizabeth N Pearce, Mònica Guxens, Eric A P Steegers, John P

- Walsh, Tim I M Korevaar. 2020. Association of maternal thyroid function with birthweight: a systematic review and individual-participant data meta-analysis. *The Lancet Diabetes & Endocrinology* 8:6, 501-510. [[Crossref](#)]
753. Rima K Dhillon-Smith. 2020. Maternal thyroid function, levothyroxine, and birthweight—a balancing act. *The Lancet Diabetes & Endocrinology* 8:6, 461-462. [[Crossref](#)]
754. Rebecca M. Lebeaux, Brett T. Doherty, Lisa G. Gallagher, R. Thomas Zoeller, Andrew N. Hoofnagle, Antonia M. Calafat, Margaret R. Karagas, Kimberly Yolton, Aimin Chen, Bruce P. Lanphear, Joseph M. Braun, Megan E. Romano. 2020. Maternal serum perfluoroalkyl substance mixtures and thyroid hormone concentrations in maternal and cord sera: The HOME Study. *Environmental Research* 185, 109395. [[Crossref](#)]
755. Su Liu, Fen Xu, Hongxia Wei, Chunyu Huang, Xian Chen, Ruochun Lian, Yong Zeng. 2020. The correlation of thyroid autoimmunity and peripheral and uterine immune status in women with recurrent miscarriage. *Journal of Reproductive Immunology* 139, 103118. [[Crossref](#)]
756. J.A. Rubio García, S. Lallena Pérez, M. Ontañón Nasarre, C. Blanco Carrera. 2020. Protocolo diagnóstico y tratamiento de la patología tiroidea en la mujer embarazada y en el posparto. *Medicine - Programa de Formación Médica Continuada Acreditado* 13:13, 752-758. [[Crossref](#)]
757. Deborah Levie, Sarah C Bath, Mònica Guxens, Tim I M Korevaar, Mariana Dineva, Eduardo Fano, Jesús M Ibarluzea, Sabrina Llop, Mario Murcia, Margaret P Rayman, Jordi Sunyer, Robin P Peeters, Henning Tiemeier. 2020. Maternal Iodine Status During Pregnancy Is Not Consistently Associated with Attention-Deficit Hyperactivity Disorder or Autistic Traits in Children. *The Journal of Nutrition* 150:6, 1516-1528. [[Crossref](#)]
758. 2020. Thyroid Disease in Pregnancy. *Obstetrics & Gynecology* 135:6, e261-e274. [[Crossref](#)]
759. Jilai Xie, Lihong Jiang, Annapurna Sadhukhan, Songqing Yang, Qiuping Yao, Ping Zhou, Jinpeng Rao, Min Jin. 2020. Effect of antithyroid antibodies on women with recurrent miscarriage: A meta-analysis. *American Journal of Reproductive Immunology* 83:6. . [[Crossref](#)]
760. Maiko Kakita-Kobayashi, Hiromi Murata, Akemi Nishigaki, Yoshiko Hashimoto, Shinnosuke Komiya, Hiroaki Tsubokura, Takeharu Kido, Naoko Kida, Tomoko Tsuzuki-Nakao, Yoshiyuki Matsuo, Hidemasa Bono, Kiichi Hirota, Hidetaka Okada. 2020. Thyroid Hormone Facilitates in vitro Decidualization of Human Endometrial Stromal Cells via Thyroid Hormone Receptors. *Endocrinology* 161:6. . [[Crossref](#)]
761. Yanan Zhang, Xinmei Huang, Zaoping Chen, Qian Yang, Xiaoying Li, Rui Zhang, Zhiyan Yu, Qian Li, Jun Liu, Bingbing Zha. 2020. IRON DEFICIENCY, A RISK FACTOR FOR THYROID AUTOIMMUNITY DURING SECOND TRIMESTER OF PREGNANCY IN CHINA. *Endocrine Practice* 26:6, 595-603. [[Crossref](#)]
762. Christopher Alan Muir, Ashish Munsif, Kenrick Blaker, Yvonne Feng, Mario D'Souza, Shailja Tewari. 2020. Antenatal Thyroid Function Does Not Increase Risk of Gestational Diabetes Mellitus in a Multi-Ethnic Pregnancy Cohort. *International Journal of Thyroidology* 13:1, 13-18. [[Crossref](#)]
763. Siew Cheng Foong, May Loong Tan, Wai Cheng Foong, Lisa A Marasco, Jacqueline J Ho, Joo Howe Ong. 2020. Oral galactagogues (natural therapies or drugs) for increasing breast milk production in mothers of non-hospitalised term infants. *Cochrane Database of Systematic Reviews* 143. . [[Crossref](#)]
764. Chun-Jui Huang, Chi-Lung Tseng, Harn-Shen Chen, Chii-Min Hwu, Kam-Tsun Tang, Justin Ging-Shing Won, Chiao-Wei Shih, Chang-Ching Yeh, Chen-Chang Yang, Fan-Fen Wang. 2020. Iodine nutritional status of pregnant women in an urban area of northern Taiwan in 2018. *PLOS ONE* 15:5, e0233162. [[Crossref](#)]
765. Anna-Simone J. Frank, David S. Matteson, Hiroko K. Solvang, Angela Lupattelli, Hedvig Nordeng. 2020. Extending balance assessment for the generalized propensity score under multiple imputation. *Epidemiologic Methods* 9:1. . [[Crossref](#)]
766. Leqi He, Xiuju Zhu, Qian Yang, Xiaoying Li, Xinmei Huang, Chunmei Shen, Jun Liu, Bingbing Zha. 2020. Low Serum IL-17A in Pregnancy During Second Trimester Is Associated With an Increased Risk of Subclinical Hypothyroidism. *Frontiers in Endocrinology* 11. . [[Crossref](#)]
767. Chutintorn Sriphrapradang. 2020. Diagnosis and Management of Graves' Disease in Thailand: A Survey of Current Practice. *Journal of Thyroid Research* 2020, 1-8. [[Crossref](#)]
768. Elly Morros González, Leonardo Javier Rojas Melo, Viviana Cruz Ramírez, Angélica Imitola. 2020. Tiroidectomía en paciente embarazada con enfermedad de Graves sin respuesta a tratamiento médico: reporte de caso. *Universitas Médica* 61:3. . [[Crossref](#)]
769. Oktay Bulur, Zeliha Atak, Derun Taner Ertugrul, Esin Beyan, Emre Gunakan, Serdar Karakaya, Kubilay Sahin, Kursat Dal. 2020. Trimester-specific reference intervals of thyroid function tests in Turkish pregnant. *Gynecological Endocrinology* 36:5, 413-416. [[Crossref](#)]
770. Thanuya Francis, Niroshan Francis, John H. Lazarus, Onyebuchi E. Okosieme. 2020. Safety of antithyroid drugs in pregnancy: update and therapy implications. *Expert Opinion on Drug Safety* 19:5, 565-576. [[Crossref](#)]

771. Niroshan Francis, Thanuya Francis, John H Lazarus, Onyebuchi E Okosieme. 2020. Current controversies in the management of Graves' hyperthyroidism. *Expert Review of Endocrinology & Metabolism* **15**:3, 159-169. [[Crossref](#)]
772. Gilbert H. Daniels,. 2020. Should the Toxicity of Potassium Iodide Treatment for Hyperthyroidism in Pregnancy Be Taken with a "Grain of Salt"? *Clinical Thyroidology* **32**:5, 245-250. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
773. Ursula Rochau, Vjollca Qerimi Rushaj, Monika Schaffner, Marie Schönhensch, Igor Stojkov, Beate Jahn, Alicja Hubalewska-Dydejczyk, Iris Erlund, Betina H. Thuesen, Michael Zimmermann, Rodrigo Moreno-Reyes, John H. Lazarus, Henry Völzke, Uwe Siebert. 2020. Decision-Analytic Modeling Studies in Prevention and Treatment of Iodine Deficiency and Thyroid Disorders: A Systematic Overview. *Thyroid* **30**:5, 746-758. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
774. Laura Alessandrini, Pietro Locantore, Alfredo Pontecorvi. 2020. Quando "trattare" il TSH nella donna delle coppie infertili?. *L'Endocrinologo* **21**:S1, 19-20. [[Crossref](#)]
775. Francesco Trimarchi, Luigi Bartalena, Paola Fierabracci, Francesco Vermiglio, Paolo Vitti, Luca Chiovato. 2020. Funzione tiroidea, concepimento e gravidanza: un dibattito aperto. *L'Endocrinologo* **21**:S1, 75-77. [[Crossref](#)]
776. Chao Wang, Jiansheng Zhu, Zhan Zhang, Hang Chen, Minghui Ji, Chao Chen, Yuhuan Hu, Yongquan Yu, Rong Xia, Jiemiao Shen, Xing Gong, Shou-Lin Wang. 2020. Rno-miR-224-5p contributes to 2,2',4,4'-tetrabromodiphenyl ether-induced low triiodothyronine in rats by targeting deiodinases. *Chemosphere* **246**, 125774. [[Crossref](#)]
777. Ran S. Rotem, Gabriel Chodick, Varda Shalev, Michael Davidovitch, Gideon Koren, Russ Hauser, Brent A. Coull, Ellen W. Seely, Vy T. Nguyen, Marc G. Weisskopf. 2020. Maternal Thyroid Disorders and Risk of Autism Spectrum Disorder in Progeny. *Epidemiology* **31**:3, 409-417. [[Crossref](#)]
778. Sun Y Lee, Howard J Cabral, Ann Aschengrau, Elizabeth N Pearce. 2020. Associations Between Maternal Thyroid Function in Pregnancy and Obstetric and Perinatal Outcomes. *The Journal of Clinical Endocrinology & Metabolism* **105**:5, e2015-e2023. [[Crossref](#)]
779. Scrinic Olesia, Delia Corina Elena, Toma Geanina Mirela, Circo Eduard. 2020. Indicators of Iodine Status in Pregnancy and Postpartum in a Group of Pregnant Women from Perimarine Area of Romania. *ARS Medica Tomitana* **26**:2, 63-69. [[Crossref](#)]
780. Maria Teresa Murillo-Llorente, Carmen Fajardo-Montañana, Marcelino Pérez-Bermejo. 2020. Reference Values of Thyroid Hormones During the First Trimester of Pregnancy in Valencian Community (Spain) and Their Relationship with Iodine Intake. *Nutrients* **12**:5, 1433. [[Crossref](#)]
781. Panagiotis Tsikouras, Theodora Deftereou, Xanthoula Anthoulaki, Anastasia Bothou, Anna Chalkidou, Fotini Gaitatzi, Ioannis Tsirkas, Arsou Chalil Bourazan, Eirini Bampageorgaka, George Stanulov, Eleftherios Chatzimichael, Spyridon Michalopoulos, Panagiotis Petsidis, Georgios Iatrakis, Stefanos Zervoudis, Maria Lambropoulou, Werner Rath, Georgios Galazios. Abortions in First Trimester Pregnancy, Management, Treatment . [[Crossref](#)]
782. Salvatore Benvenga. 2020. Targeted Antenatal Screening for Predicting Postpartum Thyroiditis and Its Evolution Into Permanent Hypothyroidism. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
783. Yang Yang, Yuanyuan Hou, Huiru Wang, Xiaotong Gao, Xichang Wang, Jiashu Li, Weiping Teng, Zhongyan Shan. 2020. Maternal Thyroid Dysfunction and Gestational Anemia Risk: Meta-Analysis and New Data. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
784. Prakruti Dash, Rajlaxmi Tiwari. Autoimmune Basis of Sub Clinical Hypothyroidism in Pregnancy . [[Crossref](#)]
785. Soledad Velasco, Francisca Illanes. 2020. Unmasking Thyroid Disease in the Postpartum Period. *Journal of Women's Health* **29**:4, 473-474. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
786. Hanni Ke, Jingmei Hu, Lijuan Zhao, Lingling Ding, Xue Jiao, Yingying Qin. 2020. Impact of Thyroid Autoimmunity on Ovarian Reserve, Pregnancy Outcomes, and Offspring Health in Euthyroid Women Following In Vitro Fertilization/Intracytoplasmic Sperm Injection. *Thyroid* **30**:4, 588-597. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
787. Lai Yuk Yuen, Michael Ho Ming Chan, Daljit Singh Sahota, Lydia Choi Wan Lit, Chung Shun Ho, Ronald Ching Wan Ma, Wing Hung Tam. 2020. Development of Gestational Age-Specific Thyroid Function Test Reference Intervals in Four Analytic Platforms Through Multilevel Modeling. *Thyroid* **30**:4, 598-608. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
788. Chrysoula Dosiou. 2020. Thyroid and Fertility: Recent Advances. *Thyroid* **30**:4, 479-486. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
789. Samantha J. Lain, Sheryl L. Rifas-Shiman, Elizabeth N. Pearce, Natasha Nassar, Emily Oken. 2020. Neonatal Thyroxine, Maternal Thyroid Function, and Cognition in Mid-childhood in a US Cohort. *Maternal and Child Health Journal* **24**:4, 503-513. [[Crossref](#)]
790. Sima Nazarpour, Fahimeh Ramezani Tehrani, Mina Amiri, Masoumeh Simbar, Maryam Tohidi, Razieh Bidhendi Yarandi, Fereidoun Azizi. 2020. Maternal Urinary Iodine Concentration and Pregnancy Outcomes: Tehran Thyroid and Pregnancy Study. *Biological Trace Element Research* **194**:2, 348-359. [[Crossref](#)]

791. Stine Linding Andersen. 2020. Farmaci antitiroidei in gravidanza e malformazioni congenite: un aggiornamento. *L'Endocrinologo* 21:2, 76-79. [[Crossref](#)]
792. Gro D. Villanger, Samantha S.M. Drover, Rachel C. Nethery, Cathrine Thomsen, Amrit K. Sakhi, Kristin R. Øvergaard, Pal Zeiner, Jane A. Hoppin, Ted Reichborn-Kjennerud, Heidi Aase, Stephanie M. Engel. 2020. Associations between urine phthalate metabolites and thyroid function in pregnant women and the influence of iodine status. *Environment International* 137, 105509. [[Crossref](#)]
793. Andrew T. Day, David J. Sher, Rebecca C. Lee, John M. Truelson, Larry L. Myers, Baran D. Sumer, Lenka Stankova, Brittny N. Tillman, Randall S. Hughes, Saad A. Khan, Eli A. Gordin. 2020. Head and neck oncology during the COVID-19 pandemic: Reconsidering traditional treatment paradigms in light of new surgical and other multilevel risks. *Oral Oncology* 104684. [[Crossref](#)]
794. Geneviève Leduc-Robert, Mahmoud Iews, Amr O Abdelkareem, Christina Williams, Dena Bloomenthal, Faten Abdelhafez, Mohamed A Bedaiwy. 2020. Prevalence of thyroid autoimmunity and effect of levothyroxine treatment in a cohort of 1064 patients with recurrent pregnancy loss. *Reproductive BioMedicine Online* 40:4, 582-592. [[Crossref](#)]
795. Jon T. Andersen, Tina Bergmann Futtrup. 2020. Drugs in pregnancy. *Adverse Drug Reaction Bulletin* 321:1, 1243-1246. [[Crossref](#)]
796. Heinz G. Bohnet. 2020. Monitoring of Thyroid Malfunction and Therapies in Pregnancy and the Postpartum Period: A Systematic Updated Critical Review of the Literature. *Therapeutic Drug Monitoring* 42:2, 222-228. [[Crossref](#)]
797. Siew Mei Yap, Mary Dillon, Rachel K Crowley, Christopher McGuigan. 2020. Alemtuzumab-related thyroid disease in people with multiple sclerosis is associated with age and brainstem phenotype at disease onset. *Multiple Sclerosis Journal - Experimental, Translational and Clinical* 6:2, 205521732093392. [[Crossref](#)]
798. Xiaodong Sun, Ningning Hou, Hongsheng Wang, Lin Ma, Jinhong Sun, Yongping Liu. 2020. A Meta-Analysis of Pregnancy Outcomes With Levothyroxine Treatment in Euthyroid Women With Thyroid Autoimmunity. *The Journal of Clinical Endocrinology & Metabolism* 105:4, 1009-1019. [[Crossref](#)]
799. Andreea Borlea, Florin Borcan, Ioan Sporea, Cristina Dehelean, Romeo Negrea, Laura Cotoi, Dana Stoian. 2020. TI-RADS Diagnostic Performance: Which Algorithm Is Superior and How Elastography and 4D Vascularity Improve the Malignancy Risk Assessment. *Diagnostics* 10:4, 180. [[Crossref](#)]
800. Xuelian Fu, Huifang Cong, Shuyu Zhao, Yan Li, Tianyi Liu, Yuhong Sun, Nan Lv. 2020. Construction of Glycometabolism- and Hormone-Related lncRNA-Mediated Feedforward Loop Networks Reveals Global Patterns of lncRNAs and Drug Repurposing in Gestational Diabetes. *Frontiers in Endocrinology* 11. . [[Crossref](#)]
801. Elizabeth N. Pearce. 2020. Substituting Potassium Iodide For Methimazole In First-Trimester Pregnant Women With Graves' Disease May Unpredictably Worsen Hyperthyroidism. *Clinical Thyroidology* 32:3, 117-119. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
802. Ai Yoshihara, Jaeduk Yoshimura Noh, Natsuko Watanabe, Yo Kunii, Miho Suzuki, Masako Matsumoto, Nami Suzuki, Marino Hiruma, Mitsuha Morisaki, Akiko Kawaguchi, Kiminori Sugino, Koichi Ito. 2020. Characteristics of Patients with Graves' Disease Whose Thyroid Hormone Levels Increase After Substituting Potassium Iodide for Methimazole in the First Trimester of Pregnancy. *Thyroid* 30:3, 451-456. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
803. Miriam Cellini, Maria Giulia Santaguida, Iliara Stramazzo, Silvia Capriello, Nunzia Brusca, Alessandro Antonelli, Poupak Fallahi, Lucilla Gargano, Marco Centanni, Camilla Virili. 2020. Recurrent Pregnancy Loss in Women with Hashimoto's Thyroiditis with Concurrent Non-Endocrine Autoimmune Disorders. *Thyroid* 30:3, 457-462. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
804. Maria Andersson, Sandra Hunziker, Ralph Fingerhut, Michael B. Zimmermann, Isabelle Herter-Aeberli. 2020. Effectiveness of increased salt iodine concentration on iodine status: trend analysis of cross-sectional national studies in Switzerland. *European Journal of Nutrition* 59:2, 581-593. [[Crossref](#)]
805. Deniz Karcaaltincaba, Murat Aykut Ozek, Nagehan Ocal, Pinar Calis, Melis Altug Inan, Merih Bayram. 2020. Prevalences of subclinical and overt hypothyroidism with universal screening in early pregnancy. *Archives of Gynecology and Obstetrics* 301:3, 681-686. [[Crossref](#)]
806. E. Minaldi, S. D'Andrea, C. Castellini, A. Martorella, F. Francavilla, S. Francavilla, A. Barbonetti. 2020. Thyroid autoimmunity and risk of post-partum depression: a systematic review and meta-analysis of longitudinal studies. *Journal of Endocrinological Investigation* 43:3, 271-277. [[Crossref](#)]
807. Alicja Hubalewska-Dydejczyk, Leonidas Duntas, Aleksandra Gilis-Januszczywska. 2020. Pregnancy, thyroid, and the potential use of selenium. *Hormones* 19:1, 47-53. [[Crossref](#)]
808. Flavia Di Bari, Sarah Perelli, Angela Scilipoti, Malgorzata Wasniewska, Roberto Vita, Francesco Vermiglio, Salvatore Benvenaga, Mariacarla Moleti. 2020. Stress-Triggered Graves' Disease with Multiple Exacerbations in a Pregnant Woman with High Levels of Thyrotropin Receptor Antibodies and No Complicated Delivery: a Case Report. *SN Comprehensive Clinical Medicine* 2:3, 355-360. [[Crossref](#)]

809. Allan C. Dong, Jessica Morgan, Monica Kane, Alex Stagnaro-Green, Mary D. Stephenson. 2020. Subclinical hypothyroidism and thyroid autoimmunity in recurrent pregnancy loss: a systematic review and meta-analysis. *Fertility and Sterility* **113**:3, 587-600.e1. [[Crossref](#)]
810. Diana Alecsandru, Juan Antonio Garcia Velasco. 2020. Levothyroxine and thyroid peroxidase antibodies in women with recurrent pregnancy loss. *Fertility and Sterility* **113**:3, 546. [[Crossref](#)]
811. Kristian Hillert Winther, Margaret Philomena Rayman, Steen Joop Bonnema, Laszlo Hegedüs. 2020. Selenium in thyroid disorders — essential knowledge for clinicians. *Nature Reviews Endocrinology* **16**:3, 165-176. [[Crossref](#)]
812. Shinyoung Jun, Jaime J. Gahche, Nancy Potischman, Johanna T. Dwyer, Patricia M. Guenther, Katherine A. Sauder, Regan L. Bailey. 2020. Dietary Supplement Use and Its Micronutrient Contribution During Pregnancy and Lactation in the United States. *Obstetrics & Gynecology* **135**:3, 623-633. [[Crossref](#)]
813. Kepal N. Patel, Linwah Yip, Carrie C. Lubitz, Elizabeth G. Grubbs, Barbra S. Miller, Wen Shen, Peter Angelos, Herbert Chen, Gerard M. Doherty, Thomas J. Fahey, Electron Kebebew, Virginia A. Livolsi, Nancy D. Perrier, Jennifer A. Sipos, Julie A. Sosa, David Steward, Ralph P. Tufano, Christopher R. McHenry, Sally E. Carty. 2020. The American Association of Endocrine Surgeons Guidelines for the Definitive Surgical Management of Thyroid Disease in Adults. *Annals of Surgery* **271**:3, e21-e93. [[Crossref](#)]
814. P. Lakshmi, D. Ramyachitra. Comparative analysis of Classification algorithms for hyperthyroidism 1241-1244. [[Crossref](#)]
815. Charlotte Hales, Peter N Taylor, Sue Channon, Kirsten McEwan, Anita Thapar, Kate Langley, Iliaria Muller, Mohd S Draman, Colin Dayan, John W Gregory, Onyebuchi Okosieme, John H Lazarus, D Aled Rees, Marian Ludgate. 2020. Controlled Antenatal Thyroid Screening II: Effect of Treating Maternal Suboptimal Thyroid Function on Child Behavior. *The Journal of Clinical Endocrinology & Metabolism* **105**:3, e417-e427. [[Crossref](#)]
816. Greisa Vila, Maria Fleseriu. 2020. Fertility and Pregnancy in Women With Hypopituitarism: A Systematic Literature Review. *The Journal of Clinical Endocrinology & Metabolism* **105**:3, e53-e65. [[Crossref](#)]
817. Christina Xiao, Philippe Grandjean, Damaskini Valvi, Flemming Nielsen, Tina Kold Jensen, Pal Weihe, Youssef Oulhote. 2020. Associations of Exposure to Perfluoroalkyl Substances With Thyroid Hormone Concentrations and Birth Size. *The Journal of Clinical Endocrinology & Metabolism* **105**:3, 735-745. [[Crossref](#)]
818. Yang Zhang, Weijie Sun, Sainan Zhu, Youyuan Huang, Yu Huang, Ying Gao, Junqing Zhang, Huixia Yang, Xiaohui Guo. 2020. The Impact of Thyroid Function and TPOAb in the First Trimester on Pregnancy Outcomes: A Retrospective Study in Peking. *The Journal of Clinical Endocrinology & Metabolism* **105**:3, e368-e380. [[Crossref](#)]
819. Małgorzata Karbownik-Lewińska, Jan Stępiak, Anna Żurawska, Andrzej Lewiński. 2020. Less Favorable Lipid Profile and Higher Prevalence of Thyroid Antibodies in Women of Reproductive Age with High-Normal TSH—Retrospective Study. *International Journal of Environmental Research and Public Health* **17**:6, 2122. [[Crossref](#)]
820. Ali Etemadi, Atieh Amouzegar, Ladan Mehran, Maryam Tohidi, Fereidoun Azizi, Kamyar Moradi, Hossein Delshad. 2020. Isolated Hypothyroxinemia in Iranian Pregnant Women, the Role of Iodine Deficiency: A Population-Based Cross-Sectional Study. *Thyroid* **30**:2, 262-269. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
821. Magdy Mohamed Allam, Hanaa Tarek El-Zawawy, Sherine Samir Barakat, Soha Magdy Ahmed, Rasha Noureldin M. Saleh. 2020. A hidden cause of infertility in hypothyroid patients. *Clinical Case Reports* **8**:2, 374-378. [[Crossref](#)]
822. Miriam Promintzer-Schifferl, Michael Krebs. 2020. Schilddrüse und Schwangerschaft: Schein und Sein. *Wiener Medizinische Wochenschrift* **170**:1-2, 35-40. [[Crossref](#)]
823. Eva Petnehazy, Wolfgang Buchinger. 2020. Hashimoto Thyreoiditis, therapeutische Optionen und extrathyreoidale Assoziationen – ein aktueller Überblick. *Wiener Medizinische Wochenschrift* **170**:1-2, 26-34. [[Crossref](#)]
824. Qiwei Xiao, Raul Cetto, Denis J. Doorly, Alister J. Bates, Jan N. Rose, Charlotte McIntyre, Andrew Comerford, Gitta Madani, Neil S. Tolley, Robert Schroter. 2020. Assessing Changes in Airflow and Energy Loss in a Progressive Tracheal Compression Before and After Surgical Correction. *Annals of Biomedical Engineering* **48**:2, 822-833. [[Crossref](#)]
825. Marco Castellana, Carlo Castellana, Luca Giovanella, Pierpaolo Trimboli. 2020. Prevalence of gastrointestinal disorders having an impact on tablet levothyroxine absorption: should this formulation still be considered as the first-line therapy?. *Endocrine* **67**:2, 281-290. [[Crossref](#)]
826. Maureen J. Lage, Jamie Vora, Zsolt Hepp, Ramon Espallat. 2020. Levothyroxine Treatment of Pregnant Women with Hypothyroidism: Retrospective Analysis of a US Claims Database. *Advances in Therapy* **37**:2, 933-945. [[Crossref](#)]
827. M. Tonacchera, L. Chiovato, L. Bartalena, A. F. Cavaliere, P. Vitti. 2020. Treatment of Graves' hyperthyroidism with thionamides: a position paper on indications and safety in pregnancy. *Journal of Endocrinological Investigation* **43**:2, 257-265. [[Crossref](#)]

828. Dagnachew Muluye Fetene, Kim S. Betts, Rosa Alati. 2020. The role of maternal prenatal thyroid function on offspring depression: Findings from the ALSPAC cohort. *Development and Psychopathology* 32:1, 189-196. [[Crossref](#)]
829. Hua Lai, Zheng-Yu Zhan, Huai Liu. 2020. Association between thyroid hormone parameters during early pregnancy and gestational hypertension: a prospective cohort study. *Journal of International Medical Research* 48:2, 030006052090481. [[Crossref](#)]
830. Bevin E. Blake, Harlie A. Cope, Samantha M. Hall, Robert D. Keys, Beth W. Mahler, James McCord, Brittany Scott, Heather M. Stapleton, Mark J. Strynar, Susan A. Elmore, Suzanne E. Fenton. 2020. Evaluation of Maternal, Embryo, and Placental Effects in CD-1 Mice following Gestational Exposure to Perfluorooctanoic Acid (PFOA) or Hexafluoropropylene Oxide Dimer Acid (HFPO-DA or GenX). *Environmental Health Perspectives* 128:2, 027006. [[Crossref](#)]
831. Daisuke Katoh, Hiroshi Yoshino, Kayoko Ikehara, Naoki Kumashiro, Hiroshi Uchino, Kumiko Tsuboi, Takahisa Hirose. 2020. Successful Treatment of Amiodarone-induced Thyrotoxicosis Type 1 in Combination with Methimazole and Potassium Iodide in a Patient with Hashimoto's Thyroiditis. *Internal Medicine* 59:3, 383-388. [[Crossref](#)]
832. Margarita Alvarez de la Rosa, Olga Rosales Aedo, Ricardo Darias Garzón, Ana Isabel Padilla Pérez, Juan Mario Troyano Luque. 2020. Iatrogenic fetal goiter. Conservative management and spontaneous resolution. *Case Reports in Perinatal Medicine* 9:1. . [[Crossref](#)]
833. Naoko Hirahara, Hiroko Misawa Nakamura, Shigekazu Sasaki, Akio Matsushita, Kenji Ohba, Go Kuroda, Yuki Sakai, Shinsuke Shinkai, Hiroshi Haeno, Takuhiro Nishio, Shuichi Yoshida, Yutaka Oki, Takafumi Suda. 2020. Liganded T3 receptor $\beta 2$ inhibits the positive feedback autoregulation of the gene for GATA2, a transcription factor critical for thyrotropin production. *PLOS ONE* 15:1, e0227646. [[Crossref](#)]
834. Pianpian Fan, Zhong-Cheng Luo, Ning Tang, Weiye Wang, Zhiwei Liu, Jun Zhang, Fengxiu Ouyang. 2020. Advanced Maternal Age, Mode of Delivery, and Thyroid Hormone Levels in Chinese Newborns. *Frontiers in Endocrinology* 10. . [[Crossref](#)]
835. Tim I.M. Korevaar. 2020. Levothyroxine Overtreatment During Pregnancy Is Associated with a Higher Risk of Adverse Child Mental Health Outcomes. *Clinical Thyroidology* 32:1, 20-23. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
836. Georgiana Sitoris, Flora Veltri, Pierre Kleynen, Alexandra Cogan, Julie Belhomme, Serge Rozenberg, Thierry Pepersack, Kris Poppe. 2020. The Impact of Thyroid Disorders on Clinical Pregnancy Outcomes in a Real-World Study Setting. *Thyroid* 30:1, 106-115. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
837. Jerome M. Hershman, Terry F. Davies, Charles H. Emerson, Peter A. Kopp. 2020. Retrospective View of Thyroid by Its Former Editors. *Thyroid* 30:1, 2-7. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
838. Hye Ok Kim, Kyungjong Lee, Sang Moo Lee, Gi Hyeon Seo. 2020. Association Between Pregnancy Outcomes and Radioactive Iodine Treatment After Thyroidectomy Among Women With Thyroid Cancer. *JAMA Internal Medicine* 180:1, 54. [[Crossref](#)]
839. Anna S. Frank, Angela Lupattelli, David S. Matteson, Helle Margrete Meltzer, Hedvig Nordeng. 2020. Thyroid hormone replacement therapy patterns in pregnant women and perinatal outcomes in the offspring. *Pharmacoepidemiology and Drug Safety* 29:1, 111-121. [[Crossref](#)]
840. Shuhei So, Wakasa Yamaguchi, Nao Murabayashi, Naomi Miyano, Fumiko Tawara. 2020. Effect of moderately increased thyroid-stimulating hormone levels and presence of thyroid antibodies on pregnancy among infertile women. *Reproductive Medicine and Biology* 19:1, 82-88. [[Crossref](#)]
841. Lorraine Lau, Ralf Paschke. Antithyroid Drugs 1-7. [[Crossref](#)]
842. Lubaina Presswala, Alyson K. Myers. Endocrine Considerations in Pregnant Women with Chronic Kidney Disease 129-145. [[Crossref](#)]
843. Görker Sel. Pregnancy and Thyroid Diseases 157-159. [[Crossref](#)]
844. Eyas Alkhalili, Jason D. Prescott. The Role of Ultrasound in Transoral Endocrine Surgery 49-61. [[Crossref](#)]
845. Karl Neff, Kate Hunt, Jill Shawe. Chronic Medical Conditions 175-210. [[Crossref](#)]
846. Jeena Sandeep, James V. Hennessey. Thyroid Problems Encountered Specifically in Inpatients with Cardiac Disease 101-113. [[Crossref](#)]
847. Lisa B. Bernstein, Michael P. Carson, Meredith O. Cruz, Rachel K. Harrison, Amanda Johnson, Anna Kho, Beth Lewis, Sarah A. Tilstra. Obstetric Medicine 583-616. [[Crossref](#)]
848. Catharina Bullmann. Schilddrüsenfunktionsstörungen in der Reproduktionsmedizin 39-47. [[Crossref](#)]
849. Wolfgang Heyl, Jürgen Wacker. Schwangerenvorsorge 43-62. [[Crossref](#)]
850. Merel van Poelje. Schildklierandoeningen 243-247. [[Crossref](#)]

851. Chunlei Ji, Ye Bu, Chunyuan Tian, Lijun Fan, Shoujun Liu, Ying Liu, Dianjun Sun. 2020. Determination of Reference Intervals of Ratios of Concentrations of Urinary Iodine to Creatinine and Thyroid Hormone Concentrations in Pregnant Women Consuming Adequate Iodine in Harbin, Heilongjiang Province. *Biological Trace Element Research* **193**:1, 36-43. [[Crossref](#)]
852. Anthony P. Weetman. Thyroid Disease 749-767. [[Crossref](#)]
853. Kirsten E. Salmeen, Ingrid J. Block-Kurbisch. Thyroid Physiology During Pregnancy, Postpartum, and Lactation 53-60. [[Crossref](#)]
854. Zoe E. Quandt, Kirsten E. Salmeen, Ingrid J. Block-Kurbisch. Thyroid Disorders During Pregnancy, Postpartum, and Lactation 287-315. [[Crossref](#)]
855. Christopher W. Rowe, Kristien Boelaert, Roger Smith. Thyroid Cancer During Pregnancy and Lactation 317-327. [[Crossref](#)]
856. Sarah Elizabeth Lawrence, Julia Elisabeth von Oettingen, Johnny Deladoëy. Fetal and Postnatal Disorders of Thyroid Function 735-754. [[Crossref](#)]
857. David C. Lin, Joely A. Straseski. Laboratory evaluation of thyroid function 703-727. [[Crossref](#)]
858. Michael Scott, Lindsey Stuart. Preconception and Pregnancy Health 1714-1738.e8. [[Crossref](#)]
859. Michael B. Zimmermann. Iodine and the iodine deficiency disorders 429-441. [[Crossref](#)]
860. N. Jung, W.-J. Mayet, P.R. Mertens, J. Meyer (federführend), O.-A. Müller, M. Pfeifer, P. Schellinger, N. Weiss, C.-M. Wendtner. ERKRANKUNGEN DER ENDOKRINEN ORGANE UND DES STOFFWECHSELS H-1-H9-9. [[Crossref](#)]
861. Sepide Talebi, Mozhgan Karimifar, Zahra Heidari, Hamed Mohammadi, Gholamreza Askari. 2020. The effects of synbiotic supplementation on thyroid function and inflammation in hypothyroid patients: A randomized, double-blind, placebo-controlled trial. *Complementary Therapies in Medicine* **48**, 102234. [[Crossref](#)]
862. Sergio Donnay, Carmen Fajardo, José Carlos Fernández-García, Teresa Torres, Orosia Bandrés, José Ramón Domínguez, Edelmiro Menéndez, Joaquín Serrano, Sara Torrejón, Irela López, José Javier Pineda, José Muñoz, Anna Lucas, Frederic Tortosa, Gracia Moll, Francisca Vich. 2020. Diagnosis, treatment, and management of gestational hypothyroidism. The TIROGEST study. *Endocrinología, Diabetes y Nutrición (English ed.)* **67**:1, 36-42. [[Crossref](#)]
863. Sergio Donnay, Carmen Fajardo, José Carlos Fernández-García, Teresa Torres, Orosia Bandrés, José Ramón Domínguez, Edelmiro Menéndez, Joaquín Serrano, Sara Torrejón, Irela López, José Javier Pineda, José Muñoz, Anna Lucas, Frederic Tortosa, Gracia Moll, Francisca Vich. 2020. Diagnosis, treatment, and management of gestational hypothyroidism. The TIROGEST study. *Endocrinología, Diabetes y Nutrición* **67**:1, 36-42. [[Crossref](#)]
864. Fangyuan Li, Zhang Qi, Lu Hua, Xinxin Wang, Mi Ling, Du Juan. 2020. The efficacy of acupuncture for the treatment and the fertility improvement in child-bearing period female with Hashimoto Disease. *Medicine* **99**:27, e20909. [[Crossref](#)]
865. Silvia D'Ippolito, Carlo Ticconi, Chiara Tersigni, Serafina Garofalo, Carmelinda Martino, Antonio Lanzzone, Giovanni Scambia, Nicoletta Di Simone. 2020. The pathogenic role of autoantibodies in recurrent pregnancy loss. *American Journal of Reproductive Immunology* **83**:1. . [[Crossref](#)]
866. Stine Linding Andersen, Stig Andersen. 2020. Turning to Thyroid Disease in Pregnant Women. *European Thyroid Journal* **9**:5, 225-233. [[Crossref](#)]
867. Emilia Sbardella, Marianna Minnetti, Riccardo Pofi, Alessia Cozzolino, Ermanno Greco, Daniele Gianfrilli, Andrea M. Isidori. 2020. Late Effects of Parasellar Lesion Treatment: Hypogonadism and Infertility. *Neuroendocrinology* **110**:9-10, 868-881. [[Crossref](#)]
868. Samantha Anandappa, Mamta Joshi, Lukasz Polanski, Paul V. Carroll. 2020. Thyroid disorders in subfertility and early pregnancy. *Therapeutic Advances in Endocrinology and Metabolism* **11**, 204201882094585. [[Crossref](#)]
869. ## #. 2020. Establishment of Reference Range of Thyroid Hormone Levels of Pregnant Women in Yan'an Area and Analysis of Thyroid Function. *Advances in Clinical Medicine* **10**:12, 2991-2998. [[Crossref](#)]
870. Yuko Inagaki, Ken Takeshima, Masahiro Nishi, Hiroyuki Ariyasu, Asako Doi, Chiaki Kurimoto, Shinsuke Uraki, Shuhei Morita, Yasushi Furukawa, Hidefumi Inaba, Hiroshi Iwakura, Toshio Shimokawa, Tomoko Utsunomiya, Takashi Akamizu. 2020. The influence of thyroid autoimmunity on pregnancy outcome in infertile women: a prospective study. *Endocrine Journal* **67**:8, 859-868. [[Crossref](#)]
871. Nami Suzuki, Ai Yoshihara, Jaeduk Yoshimura Noh, Kazuya Kinoshita, Junki Ohnishi, Maki Saito, Kiminori Sugino, Kochi Ito. 2020. TRAb elevations occurred even in the third trimester; a case of a mother of a child with neonatal thyroid dysfunction, who received radioactive iodine therapy for Graves' disease. *Endocrine Journal* **67**:10, 1019-1022. [[Crossref](#)]
872. Liane Eng, Leslie Lam. 2020. Thyroid Function During the Fetal and Neonatal Periods. *NeoReviews* **21**:1, e30-e36. [[Crossref](#)]

873. Maria Teresa Murillo-Llorente, Carmen Fajardo-Montañana, Marcelino Perez-Bermejo. 2020. Artificial Neural Network for Predicting Iodine Deficiency in the First Trimester of Pregnancy in Healthy Women. *The Tohoku Journal of Experimental Medicine* **252**:3, 185-191. [[Crossref](#)]
874. N. M. Platonova, N. P. Makolina, A. A. Rybakova, E. A. Troshina. 2020. Autoimmune thyroiditis and pregnancy: changes in the modern diagnostic and therapeutic paradigms. *Problemy reproduksii* **26**:1, 29. [[Crossref](#)]
875. K.V. Krasnopol'skaja, F.F. Burumkulova, E.A. Sokolova, K. M. Isakova, Ch.G. Gutueva. 2020. IVF in infertile women with different forms of thyroid pathology. *Problemy reproduksii* **26**:3, 46. [[Crossref](#)]
876. S.S. Paskar, A.S. Kalugina. 2020. Screening assessment of thyroid-stimulating hormone and prolactin levels and their effect on ART outcomes in women of young reproductive age. *Problemy reproduksii* **26**:6, 70. [[Crossref](#)]
877. K.V. Krasnopol'skaya, F.F. Burumkulova, Ch.G. Gutueva, E.A. Sokolova, K.M. Isakova. 2020. Effectiveness of in vitro fertilization programs in patients with thyroid disease. *Rossiiskii vestnik akushera-ginekologa* **20**:2, 64. [[Crossref](#)]
878. Tatiana Martins Benvenuto Louro Berbara, Nathalie Silva de Morais, Débora Ayres Saraiva, Carolina Martins Corcino, Annie Schtscherbyna, Karina Lúcia Moreira, Patrícia de Fátima dos Santos Teixeira, Mario Vaisman. 2020. Selective case finding versus universal screening for detecting hypothyroidism in the first trimester of pregnancy: a comparative evaluation of a group of pregnant women from Rio de Janeiro. *Archives of Endocrinology and Metabolism* . [[Crossref](#)]
879. 2020. 14. Management of Diabetes in Pregnancy: Standards of Medical Care in Diabetes—2020. *Diabetes Care* **43**:Supplement_1, S183-S192. [[Crossref](#)]
880. Renato Tozzoli, Francesca Di Serio, Paolo Metus, Vincenzo Brescia. 2020. Intervalli di riferimento del TSH in Italia: uno studio geoeidemiologico a due siti. *La Rivista Italiana della Medicina di Laboratorio* **16**:2. . [[Crossref](#)]
881. Teodora Turtă, Melinda Matyas, Lucia Procopciuc, Daniel Mureșan, Gabriela Caracostea. 2020. Updates in the management of thyroid diseases in women's reproductive health. *Obstetrica și Ginecologia* **1**:68, 16. [[Crossref](#)]
882. Allan C. Dong, Mary D. Stephenson, Alex Stewart Stagnaro-Green. 2020. The Need for Dynamic Clinical Guidelines: A Systematic Review of New Research Published After Release of the 2017 ATA Guidelines on Thyroid Disease During Pregnancy and the Postpartum. *Frontiers in Endocrinology* **11**. . [[Crossref](#)]
883. Jan Calissendorff, Henrik Falhammar. 2020. To Treat or Not to Treat Subclinical Hypothyroidism, What Is the Evidence?. *Medicina* **56**:1, 40. [[Crossref](#)]
884. Mu-Hong Chen, Kai-Lin Huang, Ju-Wei Hsu, Ya-Mei Bai, Tzeng-Ji Chen, Shih-Jen Tsai. 2020. Maternal hyperthyroidism during pregnancy and offspring risks of attention-deficit/hyperactivity disorder and autism spectrum disorder: A nationwide study. *Taiwanese Journal of Psychiatry* **34**:3, 115. [[Crossref](#)]
885. Subhadip Pramanik, Pradip Mukhopadhyay, Kingshuk Bhattacharjee, Rana Bhattacharjee, Bidisha Mukherjee, SamimAli Mondal, Sandip Bandhopadhyay, Subhas Biswas, Subhankar Chowdhury, Sujoy Ghosh. 2020. Trimester-specific reference intervals for thyroid function parameters in Indian pregnant women during final phase of transition to iodine sufficiency. *Indian Journal of Endocrinology and Metabolism* **24**:2, 160. [[Crossref](#)]
886. Leena Wadhwa, KMonica Marghret, Sarika Arora. 2020. Evaluation of Reproductive Outcome in Infertile Hypothyroid Women on Thyroxine Therapy. *Journal of Human Reproductive Sciences* **13**:4, 272. [[Crossref](#)]
887. George J. Kahaly, Tanja Diana, Paul D. Olivo. 2020. TSH RECEPTOR ANTIBODIES: RELEVANCE & UTILITY. *Endocrine Practice* **26**:1, 97-106. [[Crossref](#)]
888. Rishi Raj, Edilfavia Mae Uy, Matthew Hager, Kamyar Asadipooya. 2019. Delivery of Euthyroid Baby following Hyperthyroidism in Twin Gestation with Coexisting Complete Hydatidiform Mole. *Case Reports in Endocrinology* **2019**, 1-6. [[Crossref](#)]
889. Laura Croce, Giulia Di Dalmazi, Francesca Orsolini, Camilla Virili, Giulia Brigante, Elena Gianetti, Mariacarla Moleti, Giorgio Napolitano, Massimo Tonacchera, Mario Rotondi. 2019. Graves' Disease and the Post-partum Period: An Intriguing Relationship. *Frontiers in Endocrinology* **10**. . [[Crossref](#)]
890. Lee Sun Y.. 2019. No Increased Risks of Abortion, Preterm Delivery, or Congenital Malformation at 6 Months After Radioactive Iodine Treatment for Thyroid Cancer. *Clinical Thyroidology* **31**:12, 533-536. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
891. Sanjay Kalra, Sameer Aggarwal, Deepak Khandelwal. 2019. Thyroid Dysfunction and Type 2 Diabetes Mellitus: Screening Strategies and Implications for Management. *Diabetes Therapy* **10**:6, 2035-2044. [[Crossref](#)]
892. Dyah Purnamasari, Susie Setyowati, Em Yunir, Anandhara Indriani Khumaedi, Dante Saksono Harbuwono. 2019. Gastroschisis Following Treatment with High-Dose Methimazole in Pregnancy: A Case Report. *Drug Safety - Case Reports* **6**:1. . [[Crossref](#)]
893. Anca Zimmermann. 2019. Schilddrüsenerkrankungen in der Schwangerschaft. *Journal für Gynäkologische Endokrinologie/Österreich* **29**:4, 122-132. [[Crossref](#)]

894. Victoria Alcázar Lázaro, Teresa López del Val, Concepción García Lacalle, Beatriz Torres Moreno, Gabriela Castillo Carvajal, Lucrecia Vergara Fernández, Lina Benfdil, Covadonga Torre Carrera, María Carmen Orizales Lago, Leonardo Ramos Zuñiga. 2019. Slightly elevated thyrotropin levels in pregnancy in our clinical practice. *Endocrinología, Diabetes y Nutrición (English ed.)* **66**:10, 620-627. [[Crossref](#)]
895. Victoria Alcázar Lázaro, Teresa López del Val, Concepción García Lacalle, Beatriz Torres Moreno, Gabriela Castillo Carvajal, Lucrecia Vergara Fernández, Lina Benfdil, Covadonga Torre Carrera, María Carmen Orizales Lago, Leonardo Ramos Zuñiga. 2019. Slightly elevated thyrotropin levels in pregnancy in our clinical practice. *Endocrinología, Diabetes y Nutrición* **66**:10, 620-627. [[Crossref](#)]
896. Salvatore Benvenga, Roberto Vita, Flavia Di Bari, Carmela Lo Re, Angela Scilipoti, Grazia Giorgianni, Loredana Grasso, Marina Raffaella Galletti, Mattia Grazia Mandolino, Maria Le Donne. 2019. Assessment of serum thyroid hormone autoantibodies in the first trimester of gestation as predictors of postpartum thyroiditis. *Journal of Clinical & Translational Endocrinology* **18**, 100201. [[Crossref](#)]
897. Marlena Godlewska, Damian Gawel, Ashley M. Buckle, J. Paul Banga. 2019. Thyroid Peroxidase Revisited – What’s New?. *Hormone and Metabolic Research* **51**:12, 765-769. [[Crossref](#)]
898. Victor J. Pop, Vlad Ormindean, Andreia Mocan, Margreet Meems, Maarten Broeren, Johan K. Denollet, Wilmar M. Wiersinga, Adomas Bunevicius. 2019. Maternal cognitive function during pregnancy in relation to hypo- and hyperthyroxinemia. *Clinical Endocrinology* **91**:6, 824-833. [[Crossref](#)]
899. J. Leng, W. Li, L. Wang, S. Zhang, H. Liu, W. Li, S. Wang, P. Shao, L. Pan, S. Wang, E. Liu. 2019. Higher thyroid-stimulating hormone levels in the first trimester are associated with gestational diabetes in a Chinese population. *Diabetic Medicine* **36**:12, 1679-1685. [[Crossref](#)]
900. Zareen Kiran, Aisha Sheikh, Sarwar Malik, Areeba Meraj, Maha Masood, Safana Ismail, Muhammad Owais Rashid, Quratulain Shaikh, Numan Majeed, Luman Sheikh, Najmul Islam. 2019. Maternal characteristics and outcomes affected by hypothyroidism during pregnancy (maternal hypothyroidism on pregnancy outcomes, MHPO-1). *BMC Pregnancy and Childbirth* **19**:1. . [[Crossref](#)]
901. Chrysoula Dosiou, Alex Stagnaro-Green. 2019. The TABLET trial: limitations and implications. *BMC Medicine* **17**:1. . [[Crossref](#)]
902. Luisa Torres-Sánchez, Ricardo Gamboa, Susana Bassol-Mayagoitia, Claudia Huesca-Gómez, Martha Patricia Nava, Jennifer Illian Vázquez-Potisek, Leticia Yáñez-Estrada, Rebeca Mejía-Saucedo, Julia Blanco-Muñoz. 2019. Para-occupational exposure to pesticides, PON1 polymorphisms and hypothyroxinemia during the first half of pregnancy in women living in a Mexican floricultural area. *Environmental Health* **18**:1. . [[Crossref](#)]
903. Sarah W. Curtis, Metrecia L. Terrell, Melanie H. Jacobson, Dawayland O. Cobb, Victoria S. Jiang, Michael F. Neblett, Sabrina A. Gerkowicz, Jessica B. Spencer, M. Elizabeth Marder, Dana Boyd Barr, Karen N. Conneely, Alicia K. Smith, Michele Marcus. 2019. Thyroid hormone levels associate with exposure to polychlorinated biphenyls and polybrominated biphenyls in adults exposed as children. *Environmental Health* **18**:1. . [[Crossref](#)]
904. Maria P. Velez, Candyce Hamel, Brian Hutton, Laura Gaudet, Mark Walker, Micere Thuku, Kelly D. Cobey, Misty Pratt, Becky Skidmore, Graeme N. Smith. 2019. Care plans for women pregnant using assisted reproductive technologies: a systematic review. *Reproductive Health* **16**:1. . [[Crossref](#)]
905. Roberto Negro. 2019. Levothyroxine before conception in women with thyroid antibodies: a step forward in the management of thyroid disease in pregnancy. *Thyroid Research* **12**:1. . [[Crossref](#)]
906. Deborah Levie, Tim I M Korevaar, Sarah C Bath, Mario Murcia, Mariana Dineva, Sabrina Llop, Mercedes Espada, Antonius E van Herwaarden, Yolanda B de Rijke, Jesús M Ibarluzea, Jordi Sunyer, Henning Tiemeier, Margaret P Rayman, Mònica Guxens, Robin P Peeters. 2019. Association of Maternal Iodine Status With Child IQ: A Meta-Analysis of Individual Participant Data. *The Journal of Clinical Endocrinology & Metabolism* **104**:12, 5957-5967. [[Crossref](#)]
907. Chen Zhang, Xi Yang, Yong Zhang, Fei Guo, Shuai Yang, Robin P Peeters, Tim I M Korevaar, Jianxia Fan, He-Feng Huang. 2019. Association Between Maternal Thyroid Hormones and Birth Weight at Early and Late Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* **104**:12, 5853-5863. [[Crossref](#)]
908. Stine Linding Andersen, Louise Knøsgaard, Jørn Olsen, Peter Vestergaard, Stig Andersen. 2019. Maternal Thyroid Function, Use of Antithyroid Drugs in Early Pregnancy, and Birth Defects. *The Journal of Clinical Endocrinology & Metabolism* **104**:12, 6040-6048. [[Crossref](#)]
909. Shoshana Sztal-Mazer. 2019. Letter to the Editor: “Association of Maternal Iodine Status With Child IQ: A Meta-Analysis of Individual-Participant Data”. *The Journal of Clinical Endocrinology & Metabolism* **104**:12, 6116-6117. [[Crossref](#)]
910. Levie Deborah, Derakhshan Arash, Shu Huan, Broeren Maarten A.C., de Poortere Ralph A., Peeters Robin P., Bornehag Carl-Gustaf, Demeneix Barbara, Korevaar Tim I.M.. 2019. The Association of Maternal Iodine Status in Early Pregnancy with Thyroid

- Function in the Swedish Environmental Longitudinal, Mother and Child, Asthma and Allergy Study. *Thyroid* **29**:11, 1660-1668. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
911. Hsu Phern Chong, Halimah Alazzani, Kristien Boelaert. 2019. Endocrine disorders in pregnancy. *Obstetrics, Gynaecology & Reproductive Medicine* **29**:11, 301-305. [[Crossref](#)]
 912. Eunice López-Muñoz, Leovigildo Mateos-Sánchez, Gabriel Enrique Mejía-Terrazas, Sharon Esperanza Bedwell-Cordero. 2019. Hypothyroidism and isolated hypothyroxinemia in pregnancy, from physiology to the clinic. *Taiwanese Journal of Obstetrics and Gynecology* **58**:6, 757-763. [[Crossref](#)]
 913. Kosuke Inoue, Beate Ritz, Stine Linding Andersen, Cecilia Høst Ramlau-Hansen, Birgit Bjerre Høyer, Bodil Hammer Bech, Tine Brink Henriksen, Eva Cecilie Bonefeld-Jørgensen, Jørn Olsen, Zeyan Liew. 2019. Perfluoroalkyl Substances and Maternal Thyroid Hormones in Early Pregnancy; Findings in the Danish National Birth Cohort. *Environmental Health Perspectives* **127**:11, 117002. [[Crossref](#)]
 914. Sehar Iqbal, Petra Rust, Lisbeth Weitensfelder, Inayat Ali, Michael Kundi, Hanns Moshhammer, Cem Ekmekcioglu. 2019. Iron and Iodine Status in Pregnant Women from A Developing Country and Its Relation to Pregnancy Outcomes. *International Journal of Environmental Research and Public Health* **16**:22, 4414. [[Crossref](#)]
 915. Simona Censi, Sara Watutantrige-Fernando, Giulia Groccia, Jacopo Manso, Mario Plebani, Diego Faggian, Monica Maria Mion, Roberta Venturini, Alessandra Andrisani, Anna Casaro, Pietro Vita, Alessandra Avogadro, Marta Camilot, Carla Scaroni, Loris Bertazza, Susi Barollo, Caterina Mian. 2019. The Effects of Iodine Supplementation in Pregnancy on Iodine Status, Thyroglobulin Levels and Thyroid Function Parameters: Results from a Randomized Controlled Clinical Trial in a Mild-to-Moderate Iodine Deficiency Area. *Nutrients* **11**:11, 2639. [[Crossref](#)]
 916. Hang Zhou, Zheng Feei Ma, Yiming Lu, Binyu Pan, Jian Shao, Liya Wang, Yanyan Du, Qihua Zhao. 2019. Assessment of Iodine Status among Pregnant Women and Neonates Using Neonatal Thyrotropin (TSH) in Mainland China after the Introduction of New Revised Universal Salt Iodisation (USI) in 2012: A Re-Emergence of Iodine Deficiency?. *International Journal of Endocrinology* **2019**, 1-7. [[Crossref](#)]
 917. Galina Kh. Safaryan, Alexander M. Gzgzryan, Lyailya Kh. Dzhemlikhanova, Dariko A. Niauri. 2019. The efficiency of IVF/ICSI protocols in female subclinical hypothyroidism and thyroid autoimmunity. *Journal of obstetrics and women's diseases* **68**:4, 83-94. [[Crossref](#)]
 918. Roberta Venturella, Alberto Vaiarelli, Danilo Cimadomo, Sara Pedri, Daniela Lico, Rossella Mazzilli, Rita Mocciaro, Laura Rienzi, Costantino Di Carlo, Filippo Maria Ubaldi, Fulvio Zullo. 2019. State of the art and emerging drug therapies for female infertility. *Gynecological Endocrinology* **35**:10, 835-841. [[Crossref](#)]
 919. Anna S. Frank, Angela Lupattelli, Ragnhild E. Brandlistuen, Hedvig Nordeng. 2019. Maternal Thyroid Hormone Replacement Therapy Exposure and Language and Communication Skills of Offspring at 8 Years of Age. *JAMA Network Open* **2**:10, e1912424. [[Crossref](#)]
 920. Akhgar Ghassabian, Livia Pierotti, Mikel Basterrechea, Leda Chatzi, Marisa Estarlich, Ana Fernández-Somoano, Abby F. Fleisch, Diane R. Gold, Jordi Julvez, Polyxeni Karakosta, Aitana Lertxundi, Maria-Jose Lopez-Espinosa, Tessa A. Mulder, Tim I. M. Korevaar, Emily Oken, Robin P. Peeters, Sheryl Rifas-Shiman, Euripides Stephanou, Adonina Tardón, Henning Tiemeier, Martine Vrijheid, Tanja G. M. Vrijkotte, Jordi Sunyer, Mònica Guxens. 2019. Association of Exposure to Ambient Air Pollution With Thyroid Function During Pregnancy. *JAMA Network Open* **2**:10, e1912902. [[Crossref](#)]
 921. Jackson Kristen, Cooper David S.. 2019. Subclinical Hypothyroidism and Thyroid Autoimmunity Are Associated with Preterm Delivery in an Individual Participant Meta-Analysis. *Clinical Thyroidology* **31**:10, 410-416. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
 922. Lee Sun Y.. 2019. Birth Rates Increase After Levothyroxine Treatment in Women with Serum Thyroid Autoimmunity and Recurrent Pregnancy Loss. *Clinical Thyroidology* **31**:10, 417-420. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
 923. Toft Daniel J.. 2019. Maternal Use of Methimazole, of Propylthiouracil, and Possibly Maternal Hypothyroidism Are Associated with an Increased Risk of Birth Defects. *Clinical Thyroidology* **31**:10, 421-423. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
 924. Bliddal Sofie, Feldt-Rasmussen Ulla, Rasmussen Åse Krogh, Kolte Astrid Marie, Hilsted Linda Marie, Christiansen Ole Bjarne, Nielsen Claus Henrik, Nielsen Henriette Svarre. 2019. Thyroid Peroxidase Antibodies and Prospective Live Birth Rate: A Cohort Study of Women with Recurrent Pregnancy Loss. *Thyroid* **29**:10, 1465-1474. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
 925. Fan Jianxia, Zhang Yong, Zhang Chen, Barjaktarovic Mirjana, Yang Xi, Peeters Robin P., Huang He-Feng, Korevaar Tim I.M.. 2019. Persistency of Thyroid Dysfunction from Early to Late Pregnancy. *Thyroid* **29**:10, 1475-1484. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
 926. Timo Minnemann, Catharina Bullmann. 2019. Schilddrüsenhormonsubstitution in der Schwangerschaft. *Der Gynäkologe* **52**:10, 745-755. [[Crossref](#)]

927. Sima Nazarpour, Fahimeh Ramezani Tehrani, Mina Amiri, Razieh Bidhendi Yarandi, Fereidoun Azizi. 2019. Levothyroxine treatment and pregnancy outcomes in women with subclinical hypothyroidism: a systematic review and meta-analysis. *Archives of Gynecology and Obstetrics* **300**:4, 805-819. [[Crossref](#)]
928. Bernadette Biondi, David S. Cooper. 2019. Thyroid hormone therapy for hypothyroidism. *Endocrine* **66**:1, 18-26. [[Crossref](#)]
929. Flavia Magri, Luca Chiovato, Laura Croce, Mario Rotondi. 2019. Thyroid hormone therapy for subclinical hypothyroidism. *Endocrine* **66**:1, 27-34. [[Crossref](#)]
930. Zhongyan Shan, Weiping Teng. 2019. Thyroid hormone therapy of hypothyroidism in pregnancy. *Endocrine* **66**:1, 35-42. [[Crossref](#)]
931. Suvarna Khadilkar. 2019. Thyroid-Stimulating Hormone Values in Pregnancy: Cutoff Controversy Continues?. *The Journal of Obstetrics and Gynecology of India* **69**:5, 389-394. [[Crossref](#)]
932. Jean-Philippe Rousseau, Anabel Buteau-Poulin, Richard Kinkead. 2019. Maternal thyroid hormone deficiency and cardiorespiratory disorder in rat pups. *Experimental Neurology* **320**, 112960. [[Crossref](#)]
933. Freddy J.K. Toloza, Sanaz Abedzadeh-Anaraki, Spyridoula Maraka. 2019. Subclinical hypothyroidism in pregnancy. *Current Opinion in Endocrinology, Diabetes & Obesity* **26**:5, 225-231. [[Crossref](#)]
934. Caroline T. Nguyen, Jorge H. Mestman. 2019. Graves' hyperthyroidism in pregnancy. *Current Opinion in Endocrinology & Diabetes and Obesity* **26**:5, 232-240. [[Crossref](#)]
935. Freddy J K Toloza, Naykky M Singh Ospina, Rene Rodriguez-Gutierrez, Derek T O'Keeffe, Juan P Brito, Victor M Montori, Spyridoula Maraka. 2019. Practice Variation in the Care of Subclinical Hypothyroidism During Pregnancy: A National Survey of Physicians in the United States. *Journal of the Endocrine Society* **3**:10, 1892-1906. [[Crossref](#)]
936. Creswell J Eastman, Gary Ma, Mu Li. 2019. Optimal Assessment and Quantification of Iodine Nutrition in Pregnancy and Lactation: Laboratory and Clinical Methods, Controversies and Future Directions. *Nutrients* **11**:10, 2378. [[Crossref](#)]
937. Efrat Spiegel, Andrea R. Spence, Nicholas Czuzoj-Shulman, Haim Arie Abenhaim. 2019. Pregnancy outcomes after thyroid cancer. *Journal of Perinatal Medicine* **47**:7, 710-716. [[Crossref](#)]
938. Tomas Salek, Ishraq Dhaifalah, Dagmar Langova, Jana Havalova. 2019. The prevalence of maternal hypothyroidism in first trimester screening from 11 to 14 weeks of gestation. *Biomedical Papers* **163**:3, 265-268. [[Crossref](#)]
939. Salman Razvi, Sindeeph Bhana, Sanaa Mrabeti. 2019. Challenges in Interpreting Thyroid Stimulating Hormone Results in the Diagnosis of Thyroid Dysfunction. *Journal of Thyroid Research* **2019**, 1-8. [[Crossref](#)]
940. Chutintorn Sriphrapadang, Sittichai Pinyopodjanard, Onnicha Suntornlohanakul, Hataikarn Nimitphong, Naricha Chirakalwasan, Sunee Saetang, Thunyarat Anothaisintawee, Nantaporn Siwasaranond, Areesa Manodpitipong, La-Or Chailurkit, Sirimon Reutrakul. 2019. Lack of associations between thyroid function and obstructive sleep apnea severity in adults with prediabetes and diabetes mellitus. *Sleep and Breathing* **23**:3, 963-967. [[Crossref](#)]
941. Leonidas H. Duntas, Jacqueline Jonklaas. 2019. Levothyroxine Dose Adjustment to Optimise Therapy Throughout a Patient's Lifetime. *Advances in Therapy* **36**:S2, 30-46. [[Crossref](#)]
942. Luca Chiovato, Flavia Magri, Allan Carlé. 2019. Hypothyroidism in Context: Where We've Been and Where We're Going. *Advances in Therapy* **36**:S2, 47-58. [[Crossref](#)]
943. Kristen Kobaly, Susan J. Mandel. 2019. Hyperthyroidism and Pregnancy. *Endocrinology and Metabolism Clinics of North America* **48**:3, 533-545. [[Crossref](#)]
944. Trevor E. Angell, Erik K. Alexander. 2019. Thyroid Nodules and Thyroid Cancer in the Pregnant Woman. *Endocrinology and Metabolism Clinics of North America* **48**:3, 557-567. [[Crossref](#)]
945. Peter N. Taylor, John H. Lazarus. 2019. Hypothyroidism in Pregnancy. *Endocrinology and Metabolism Clinics of North America* **48**:3, 547-556. [[Crossref](#)]
946. Batuhan Turgay, Yavuz Emre Şükür, Hasan Ulubaşoğlu, Murat Sönmezer, Bülent Berker, Cem Atabekoğlu, Ruşen Aytac, Batuhan Özmen. 2019. The association of thyroid stimulating hormone levels and intrauterine insemination outcomes of euthyroid unexplained subfertile couples. *European Journal of Obstetrics & Gynecology and Reproductive Biology* **240**, 99-102. [[Crossref](#)]
947. Layla Thurston, Ali Abbara, Waljit S Dhillon. 2019. Investigation and management of subfertility. *Journal of Clinical Pathology* **72**:9, 579-587. [[Crossref](#)]
948. Y. Yonekura, S. Mattsson, G. Flux, W.E. Bolch, L.T. Dauer, D.R. Fisher, M. Lassmann, S. Palm, M. Hosono, M. Doruff, C. Divgi, P. Zanzonico. 2019. ICRP Publication 140: Radiological Protection in Therapy with Radiopharmaceuticals. *Annals of the ICRP* **48**:1, 5-95. [[Crossref](#)]
949. Sandra Lowe. 2019. Diagnostic imaging in pregnancy: Making informed decisions. *Obstetric Medicine* **12**:3, 116-122. [[Crossref](#)]

950. Salvatore Benvenaga, Ulla Feldt-Rasmussen, Daniela Bonofiglio, Ernest Asamoah. 2019. Nutraceutical Supplements in the Thyroid Setting: Health Benefits beyond Basic Nutrition. *Nutrients* **11**:9, 2214. [[Crossref](#)]
951. Anne R. Cappola, Brian M. Casey. 2019. Thyroid Function Test Abnormalities During Pregnancy. *JAMA* **322**:7, 617. [[Crossref](#)]
952. Tim I. M. Korevaar, Arash Derakhshan, Peter N. Taylor, Marcel Meima, Liangmiao Chen, Sofie Bliddal, David M. Carty, Margreet Meems, Bijay Vaidya, Beverley Shields, Farkhanda Ghafoor, Polina V. Popova, Lorena Mosso, Emily Oken, Eila Suvanto, Aya Hisada, Jun Yoshinaga, Suzanne J. Brown, Judit Bassols, Juha Auvinen, Wichor M. Bramer, Abel López-Bermejo, Colin Dayan, Laura Boucai, Marina Vafeiadi, Elena N. Grineva, Alexandra S. Tkachuck, Victor J. M. Pop, Tanja G. Vrijkotte, Mònica Guxens, Leda Chatzi, Jordi Sunyer, Ana Jiménez-Zabala, Isolina Riaño, Mario Murcia, Xuemian Lu, Shafqat Mukhtar, Christian Delles, Ulla Feldt-Rasmussen, Scott M. Nelson, Erik K. Alexander, Layal Chaker, Tuija Männistö, John P. Walsh, Elizabeth N. Pearce, Eric A. P. Steegers, Robin P. Peeters. 2019. Association of Thyroid Function Test Abnormalities and Thyroid Autoimmunity With Preterm Birth. *JAMA* **322**:7, 632. [[Crossref](#)]
953. An Eerdekens, Lies Langouche, Fabian Güiza, Johan Verhaeghe, Gunnar Naulaers, Christine Vanhole, Greet Van den Berghe. 2019. Maternal and placental responses before preterm birth: adaptations to increase fetal thyroid hormone availability?. *The Journal of Maternal-Fetal & Neonatal Medicine* **32**:16, 2746-2757. [[Crossref](#)]
954. Mei-Qin Wu, Jin Liu, Ya-Qian Wang, Ying Yang, Chong-Huai Yan, Jing Hua. 2019. The Impact of Subclinical Hypothyroidism on Adverse Perinatal Outcomes and the Role of Thyroid Screening in Pregnancy. *Frontiers in Endocrinology* **10**. [[Crossref](#)]
955. Korevaar Tim I.M.. 2019. Initiation of Levothyroxine Therapy for Subclinical Hypothyroidism during Pregnancy in the United States. *Clinical Thyroidology* **31**:8, 318-321. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
956. Yesica María Rodríguez Cortés, Araceli Amaya Chávez, Magdalena García Fabila, Silvia Jiménez Morales, Carlos Jhovani Pérez Amado, Hugo Mendieta Zerón. 2019. TSH β X1 Splice Variant Expression and D2 Thr92Ala Polymorphism Analysis in Pregnant Women with Thyroid Diseases. *SN Comprehensive Clinical Medicine* **1**:8, 590-596. [[Crossref](#)]
957. Francesco Vermiglio, Mariacarla Moleti. 2019. Maternal thyroid function and brain development: time for preconception screening?. *The Lancet Diabetes & Endocrinology* **7**:8, 589-590. [[Crossref](#)]
958. Peter S. Hamblin, Penelope M. Sheehan, Carolyn Allan, Christine A. Houlihan, Zhong X. Lu, Simon P. Forehan, Duncan J. Topliss, Christopher Gilfillan, Bala Krishnamurthy, Debra Renouf, Shoshana Sztal-Mazer, Suresh Varadarajan. 2019. Subclinical hypothyroidism during pregnancy: the Melbourne public hospitals consensus. *Internal Medicine Journal* **49**:8, 994-1000. [[Crossref](#)]
959. Catarina Martins Machado, Jorge Manuel Castro, Rosa Arménia Campos, Maria João Oliveira. 2019. Graves' disease complicated by fetal goitrous hypothyroidism treated with intra-amniotic administration of levothyroxine. *BMJ Case Reports* **12**:8, e230457. [[Crossref](#)]
960. Tim I M Korevaar, Robin P Peeters. 2019. Improving Risk Stratification Strategies for Thyroid Disease During Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* **104**:8, 3262-3263. [[Crossref](#)]
961. María Dolores Martín-Escalante, Jairo Luque-del Pino, Josefa A. Aguilar-García, Victoria Agustín-Bandera, Javier García-Alegria. 2019. Abortos de repetición en paciente con anticuerpos antitiroideos positivos. *Revista Española de Casos Clínicos en Medicina Interna* **4**:2, 93-95. [[Crossref](#)]
962. Galina Khachikovna Safarian, Alexander Mkrtichevich Gzgzyan, Kharryasovna Dzhemlikhanova Lyailya, Dariko Alexandrovna Niauri. 2019. Does subclinical hypothyroidism and/or thyroid autoimmunity influence the IVF/ICSI outcome? Review of the literature. *Gynecological Endocrinology* **35**:sup1, 56-59. [[Crossref](#)]
963. Jenna Van Beck, Sobia F. Khaja. 2019. Thyroglossal Duct Cyst Carcinoma in a Young Female: Case Report and Review of Literature. *Case Reports in Otolaryngology* **2019**, 1-3. [[Crossref](#)]
964. Amit Akirov, Rouhi Fazelzad, Shereen Ezzat, Lehana Thabane, Anna M. Sawka. 2019. A Systematic Review and Meta-Analysis of Patient Preferences for Combination Thyroid Hormone Treatment for Hypothyroidism. *Frontiers in Endocrinology* **10**. [[Crossref](#)]
965. 2019. Levothyroxine in Women with Thyroid Peroxidase Antibodies before Conception. *New England Journal of Medicine* **381**:2, 190-191. [[Crossref](#)]
966. Angela X. Chen, Angela M. Leung, Tim I.M. Korevaar. 2019. Thyroid Function and Conception. *New England Journal of Medicine* **381**:2, 178-181. [[Crossref](#)]
967. Bernadette Biondi, Anne R. Cappola, David S. Cooper. 2019. Subclinical Hypothyroidism. *JAMA* **322**:2, 153. [[Crossref](#)]
968. Min Zhou, Min Wang, Juming Li, Xiaohui Luo, Minxiang Lei. 2019. Effects of thyroid diseases on pregnancy outcomes. *Experimental and Therapeutic Medicine*. [[Crossref](#)]
969. Pearce Elizabeth N.. 2019. Selenium Supplementation in Pregnancy and the Postpartum Period Decreases Postpartum Antithyroid Antibody Titers, but Does Not Affect Thyroid Function or Health-Related Quality of Life. *Clinical Thyroidology* **31**:7, 299-301. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]

970. S. Gricourt, M. Peigné, D. Luton. 2019. Síndromes endocrinológicos del posparto. *EMC - Ginecología-Obstetricia* **55**:3, 1-9. [[Crossref](#)]
971. Hajime Ishinaga, Satoshi Nakamura, Kazuhiko Takeuchi. 2019. A Case of Aggressive Thyroid Cancer Detected in the Second Half of Pregnancy. *Nihon Kikan Shokudoka Gakkai Kaibo* **70**:3, 254-257. [[Crossref](#)]
972. Mitchell Katrina B., Fleming Margaret M., Anderson Philip O., Giesbrandt Jamie G., the Academy of Breastfeeding Medicine. 2019. ABM Clinical Protocol #31: Radiology and Nuclear Medicine Studies in Lactating Women. *Breastfeeding Medicine* **14**:5, 290-294. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
973. Cho Geum Joon, Kim So-youn, Lee Hoi Chang, Lee Kyu-Min, Han Sung Won, Oh Min-Jeong, Woodruff Teresa K.. 2019. Risk of Adverse Obstetric Outcomes and the Abnormal Growth of Offspring in Women with a History of Thyroid Cancer. *Thyroid* **29**:6, 879-885. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
974. M Ahsan Akhtar, Rina Agrawal, Julie Brown, Yasmin Sajjad, Laurentiu Craciunas. 2019. Thyroxine replacement for subfertile women with euthyroid autoimmune thyroid disease or subclinical hypothyroidism. *Cochrane Database of Systematic Reviews* **2019**:6. . [[Crossref](#)]
975. Paola Fierabracci. 2019. Terapia con levotiroxina prima del concepimento in donne con positività degli anticorpi anti-tireoperossidasi. *L'Endocrinologo* **20**:3, 183-184. [[Crossref](#)]
976. Carole Emile. 2019. Bilan biologique avant assistance médicale à la procréation. *Option/Bio* **30**:599-600, 19-20. [[Crossref](#)]
977. Jacqueline Jonklaas, Salman Razvi. 2019. Reference intervals in the diagnosis of thyroid dysfunction: treating patients not numbers. *The Lancet Diabetes & Endocrinology* **7**:6, 473-483. [[Crossref](#)]
978. Lía Nattero-Chávez, Manuel Luque-Ramírez, Héctor F. Escobar-Morreale. 2019. Systemic endocrinopathies (thyroid conditions and diabetes): impact on postnatal life of the offspring. *Fertility and Sterility* **111**:6, 1076-1091. [[Crossref](#)]
979. Mariacarla Moleti, Maria Di Mauro, Giacomo Sturniolo, Marco Russo, Francesco Vermiglio. 2019. Hyperthyroidism in the pregnant woman: Maternal and fetal aspects. *Journal of Clinical & Translational Endocrinology* **16**, 100190. [[Crossref](#)]
980. Anu Sharma, Marius N. Stan. 2019. Thyrotoxicosis: Diagnosis and Management. *Mayo Clinic Proceedings* **94**:6, 1048-1064. [[Crossref](#)]
981. SANJITA B. CHITTIMOJU, ELIZABETH N. PEARCE. 2019. Iodine Deficiency and Supplementation in Pregnancy. *Clinical Obstetrics & Gynecology* **62**:2, 330-338. [[Crossref](#)]
982. CAROLINE T. NGUYEN, JORGE H. MESTMAN. 2019. Postpartum Thyroiditis. *Clinical Obstetrics & Gynecology* **62**:2, 359-364. [[Crossref](#)]
983. SCOTT A. SULLIVAN. 2019. Thyroid Nodules and Thyroid Cancer in Pregnancy. *Clinical Obstetrics & Gynecology* **62**:2, 365-372. [[Crossref](#)]
984. SCOTT A. SULLIVAN. 2019. Hypothyroidism in Pregnancy. *Clinical Obstetrics & Gynecology* **62**:2, 308-319. [[Crossref](#)]
985. ALISON K. CHAPMAN, ZACHARY J. FARMER, LUCY D. MASTRANDREA, KRISTAL A. MATLOCK. 2019. Neonatal Thyroid Function and Disorders. *Clinical Obstetrics & Gynecology* **62**:2, 373-387. [[Crossref](#)]
986. RYAN D. CUFF. 2019. Hyperthyroidism During Pregnancy. *Clinical Obstetrics and Gynecology* **62**:2, 320-329. [[Crossref](#)]
987. MARY STERRETT. 2019. Maternal and Fetal Thyroid Physiology. *Clinical Obstetrics & Gynecology* **62**:2, 302-307. [[Crossref](#)]
988. SCOTT A. SULLIVAN, RYAN D. CUFF, CHRISTOPHER G. GOODIER, ALISON K. CHAPMAN, SANJITA B. CHITTIMOJU, MARY STERRETT. 2019. Case Studies in Thyroid Dysfunction and Pregnancy. *Clinical Obstetrics & Gynecology* **62**:2, 388-397. [[Crossref](#)]
989. Bernadette Biondi, George J Kahaly, R Paul Robertson. 2019. Thyroid Dysfunction and Diabetes Mellitus: Two Closely Associated Disorders. *Endocrine Reviews* **40**:3, 789-824. [[Crossref](#)]
990. Georgiana Sitoris, Flora Veltri, Pierre Kleynen, Julie Belhomme, Serge Rozenberg, Kris Poppe. 2019. Screening for Thyroid Dysfunction in Pregnancy With Targeted High-Risk Case Finding: Can It Be Improved?. *The Journal of Clinical Endocrinology & Metabolism* **104**:6, 2346-2354. [[Crossref](#)]
991. Anh Tran, Steve Hyer, Imran Rafi, Onyebuchi Okosieme. 2019. Thyroid hormone replacement in the preconception period and pregnancy. *British Journal of General Practice* **69**:683, 282-283. [[Crossref](#)]
992. Angelo Maria Patti, Rosaria Vincenza Giglio, Kalliopi Pafili, Manfredi Rizzo, Nikolaos Papanas. 2019. Advances in pharmacological treatment of type 1 diabetes during pregnancy. *Expert Opinion on Pharmacotherapy* **20**:8, 983-989. [[Crossref](#)]
993. Laura T. Dickens, Adam S. Cifu, Ronald N. Cohen. 2019. Diagnosis and Management of Thyroid Disease During Pregnancy and the Postpartum Period. *JAMA* **321**:19, 1928. [[Crossref](#)]

994. Anastasia-Stefania Alexopoulos, Rachel Blair, Anne L. Peters. 2019. Management of Preexisting Diabetes in Pregnancy. *JAMA* **321**:18, 1811. [[Crossref](#)]
995. G E Bekkering, T Agoritsas, L Lytvyn, A F Heen, M Feller, E Moutzouri, H Abdulazeem, B Aertgeerts, D Beecher, J P Brito, P D Farhoumand, N Singh Ospina, N Rodondi, M van Driel, E Wallace, M Snel, P M Okwen, R Siemieniuk, P O Vandvik, T Kuijpers, M Vermandere. 2019. Thyroid hormones treatment for subclinical hypothyroidism: a clinical practice guideline. *BMJ* **l2006**. [[Crossref](#)]
996. Pearce Elizabeth N.. 2019. Mild-to-Moderate Iodine Deficiency in Early Pregnancy Is Associated with Lower Verbal IQ in Children. *Clinical Thyroidology* **31**:5, 195-197. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
997. Grulet Hervé, Barraud Sara, Chikh Karim, Souchon Pierre François, Claris Olivier, Bouvier Raymonde, Trouillas Jacqueline, Gabriel René, Bory Jean Paul, Schwartz Claire, Orgiazzi Jacques, Delemer Brigitte. 2019. Three Consecutive Pregnancies in a Patient with Chronic Autoimmune Thyroid Disease Associated with Hypothyroidism and Extremely High Levels of Anti-Thyrotropin Receptor Antibodies. *Thyroid* **29**:5, 743-747. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
998. Alexandra Kautzky-Willer, Jürgen Harreiter, Yvonne Winhofer-Stöckl, Raimund Weitgasser, Monika Lechleitner. 2019. Gravidität bei vorbestehendem Diabetes (Update 2019). *Wiener klinische Wochenschrift* **131**:S1, 103-109. [[Crossref](#)]
999. X. Gong, A. Liu, Y. Li, H. Sun, Y. Li, C. Li, X. Yu, C. Fan, Z. Shan, W. Teng. 2019. The impact of isolated maternal hypothyroxinemia during the first and second trimester of gestation on pregnancy outcomes: an intervention and prospective cohort study in China. *Journal of Endocrinological Investigation* **42**:5, 599-607. [[Crossref](#)]
1000. M. Dolores Ollero, Javier Pineda, Juan Pablo Martínez de Esteban, Marta Toni, Mercedes Espada, Emma Anda. 2019. Optimization of the follow-up of pregnant women with autoimmune thyroid disease. *Endocrinología, Diabetes y Nutrición (English ed.)* **66**:5, 305-311. [[Crossref](#)]
1001. M. Dolores Ollero, Javier Pineda, Juan Pablo Martínez de Esteban, Marta Toni, Mercedes Espada, Emma Anda. 2019. Optimización del seguimiento de gestantes con enfermedad tiroidea autoinmune. *Endocrinología, Diabetes y Nutrición* **66**:5, 305-311. [[Crossref](#)]
1002. Meng Rao, Zhengyan Zeng, Fang Zhou, Huawei Wang, Jiang Liu, Rui Wang, Ya Wen, Zexing Yang, Cunmei Su, Zhenfang Su, Shuhua Zhao, Li Tang. 2019. Effect of levothyroxine supplementation on pregnancy loss and preterm birth in women with subclinical hypothyroidism and thyroid autoimmunity: a systematic review and meta-analysis. *Human Reproduction Update* **25**:3, 344-361. [[Crossref](#)]
1003. Ziyun Pan, Tingkai Cui, Wen Chen, Shu Gao, Elizabeth N. Pearce, Wei Wang, Yanting Chen, Wenxing Guo, Long Tan, Jun Shen, Wanqi Zhang. 2019. Serum iodine concentration in pregnant women and its association with urinary iodine concentration and thyroid function. *Clinical Endocrinology* **90**:5, 711-718. [[Crossref](#)]
1004. Maryam Kianpour, Ashraf Aminorroaya, Massoud Amini, Awat Feizi, Sima Aminorroaya Yamini, Mohsen Janghorbani. 2019. Thyroid-stimulating hormone (TSH) serum levels and risk of spontaneous abortion: A prospective population-based cohort study. *Clinical Endocrinology* **62**. . [[Crossref](#)]
1005. Jie Cai, Yi Zhang, Yuying Wang, Shengxian Li, Lihua Wang, Jun Zheng, Yihong Jiang, Ying Dong, Huan Zhou, Yaomin Hu, Jing Ma, Wei Liu, Tao Tao. 2019. High Thyroid Stimulating Hormone Level Is Associated With Hyperandrogenism in Euthyroid Polycystic Ovary Syndrome (PCOS) Women, Independent of Age, BMI, and Thyroid Autoimmunity: A Cross-Sectional Analysis. *Frontiers in Endocrinology* **10**. . [[Crossref](#)]
1006. Rima K. Dhillon-Smith, Lee J. Middleton, Kirandeep K. Sunner, Versha Cheed, Krys Baker, Samantha Farrell-Carver, Ruth Bender-Atik, Rina Agrawal, Kalsang Bhatia, Edmond Edi-Osagie, Tarek Ghobara, Pratima Gupta, Davor Jurkovic, Yacoub Khalaf, Marjory MacLean, Christopher McCabe, Khashia Mulbagal, Natalie Nunes, Caroline Overton, Siobhan Quenby, Raj Rai, Nick Raine-Fenning, Lynne Robinson, Jackie Ross, Andrew Sizer, Rachel Small, Alex Tan, Martyn Underwood, Mark D. Kilby, Kristien Boelaert, Jane Daniels, Shakila Thangaratinam, Shiao Y. Chan, Arri Coomarasamy. 2019. Levothyroxine in Women with Thyroid Peroxidase Antibodies before Conception. *New England Journal of Medicine* **380**:14, 1316-1325. [[Crossref](#)]
1007. Mario Rotondi, Valentina Capelli, Luca Chiovato, Rossella E. Nappi. 2019. 2017 ATA guidelines on the management of thyroid dysfunctions in pregnancy: what do OB/GYNs need to know?. *Gynecological Endocrinology* **35**:4, 276-279. [[Crossref](#)]
1008. M. C. Черенько, С. М. Черенько. 2019. Management of thyroid disorders during and after pregnancy. *Reproductive Endocrinology* **0**:45, 14-18. [[Crossref](#)]
1009. Hou Yuanyuan, Liu Aihua, Li Jiashu, Wang Haoyu, Yang Yang, Li Yongze, Fan Chenling, Zhang Hongmei, Wang Hong, Ding Shuangning, Teng Weiping, Shan Zhongyan. 2019. Different Thyroidal Responses to Human Chorionic Gonadotropin Under Different Thyroid Peroxidase Antibody and/or Thyroglobulin Antibody Positivity Conditions During the First Half of Pregnancy. *Thyroid* **29**:4, 577-585. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]

1010. Gil Gutvirtz, Asnat Walfisch, Tamar Wainstock, Daniella Landau, Eyal Sheiner. 2019. Maternal hypothyroidism and future pediatric neurological morbidity of the offspring. *Archives of Gynecology and Obstetrics* **299**:4, 975-981. [[Crossref](#)]
1011. Stephanie Allelein, Matthias Schott. 2019. Schilddrüse und Schwangerschaft. *gynäkologie + geburtshilfe* **24**:2, 34-42. [[Crossref](#)]
1012. Harvey J. Clewell, P. Robinan Gentry, C. Eric Hack, Tracy Greene, Rebecca A. Clewell. 2019. An evaluation of the USEPA Proposed Approaches for applying a biologically based dose-response model in a risk assessment for perchlorate in drinking water. *Regulatory Toxicology and Pharmacology* **103**, 237-252. [[Crossref](#)]
1013. Annalisa Schiepatti, Elisa Sprio, David S. Sanders, Elisabetta Lovati, Federico Biagi. 2019. Coeliac disease and obstetric and gynaecological disorders: where are we now?. *European Journal of Gastroenterology & Hepatology* **31**:4, 425-433. [[Crossref](#)]
1014. Rosa Cervelli, Salvatore Mazzeo, Giuseppe Boni, Antonio Boccuzzi, Francesca Bianchi, Federica Brozzi, Pierina Santini, Paolo Vitti, Roberto Cioni, Davide Caramella. 2019. Comparison between radioiodine therapy and single-session radiofrequency ablation of autonomously functioning thyroid nodules: A retrospective study. *Clinical Endocrinology* **90**:4, 608-616. [[Crossref](#)]
1015. Xiujuan Su, Yan Zhao, Zhijuan Cao, Yingying Yang, Tony Duan, Jing Hua. 2019. Association between isolated hypothyroxinaemia in early pregnancy and perinatal outcomes. *Endocrine Connections* **8**:4, 435-441. [[Crossref](#)]
1016. Ollero M. Dolores, Toni Marta, Pineda José Javier, Martínez Juan Pablo, Espada Mercedes, Anda Emma. 2019. Thyroid Function Reference Values in Healthy Iodine-Sufficient Pregnant Women and Influence of Thyroid Nodules on Thyrotropin and Free Thyroxine Values. *Thyroid* **29**:3, 421-429. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1017. Donovan Lois E., Metcalfe Amy, Chin Alex, Yamamoto Jennifer M., Virtanen Heidi, Johnson Jo-Ann, Krause Richard. 2019. A Practical Approach for the Verification and Determination of Site- and Trimester-Specific Reference Intervals for Thyroid Function Tests in Pregnancy. *Thyroid* **29**:3, 412-420. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1018. Andersen Stine Linding, Andersen Stig, Carlé Allan, Christensen Peter Astrup, Handberg Aase, Karmisholt Jesper, Knøsgaard Louise, Kristensen Søren Risom, Bülow Pedersen Inge, Vestergaard Peter. 2019. Pregnancy Week-Specific Reference Ranges for Thyrotropin and Free Thyroxine in the North Denmark Region Pregnancy Cohort. *Thyroid* **29**:3, 430-438. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1019. Carolina Martins Corcino, Tatiana Martins Benvenuto Louro Barbara, Débora Ayres Saraiva, Nathalie Anne de Oliveira e Silva de Moraes, Annie Schtscherbyna, Larissa Nascimento Gertrudes, Patrícia de Fátima dos Santos Teixeira, Mario Vaisman. 2019. Variation of iodine status during pregnancy and its associations with thyroid function in women from Rio de Janeiro, Brazil. *Public Health Nutrition* **67**, 1-9. [[Crossref](#)]
1020. Minghan Jia, Yanxin Wu, Bo Lin, Yawei Shi, Qi Zhang, Ying Lin, Shenming Wang, Yunjian Zhang. 2019. Meta-analysis of the association between maternal subclinical hypothyroidism and gestational diabetes mellitus. *International Journal of Gynecology & Obstetrics* **144**:3, 239-247. [[Crossref](#)]
1021. Wei Zhao, Xinyu Li, Xinghai Xia, Zhengnan Gao, Cheng Han. 2019. Iodine Nutrition During Pregnancy: Past, Present, and Future. *Biological Trace Element Research* **188**:1, 196-207. [[Crossref](#)]
1022. Pallavi Panth, Gena Guerin, Nancy M. DiMarco. 2019. A Review of Iodine Status of Women of Reproductive Age in the USA. *Biological Trace Element Research* **188**:1, 208-220. [[Crossref](#)]
1023. M. L. Tanda, C. Cusini, A. Colombo, P. Premoli, S. Rosetti, A. Cromi, E. Piantanida, L. Bartalena. 2019. Iodine supplementation in women of reproductive age: a survey of clinical practice among Italian gynecologists and midwives. *Journal of Endocrinological Investigation* **42**:3, 353-355. [[Crossref](#)]
1024. Maryam Kianpour, Ashraf Aminorroaya, Massoud Amini, Awat Feizi, Mohsen Janghorbani, Saeideh Shokri, Sima Aminorroaya Yamini, Maryam Farghadani, Ali Hekmatnia, Hossien Gharib. 2019. Reference Intervals for Thyroid Hormones During the First Trimester of Gestation: A Report from an Area with a Sufficient Iodine Level. *Hormone and Metabolic Research* **51**:03, 165-171. [[Crossref](#)]
1025. Shiva Borzouei, Mohammad Taghi Goodarzi, Mansoorh Biglari, Fathollah Nazari, Zahra Shivapour. 2019. The Prevalence of Thyroid Disorders in Pregnant Women of Hamadan. *Avicenna Journal of Nursing and Midwifery Care* **27**:1, 11-17. [[Crossref](#)]
1026. Laura J McNally, Catherine I Ofiaeli, Samson O Oyibo. 2019. Treatment-refractory hypothyroidism. *BMJ* **36**, l579. [[Crossref](#)]
1027. Jan Jiskra. 2019. Which of thyroid disorders should be managed by general practitioner?. *Medicina pro praxi* **16**:1, 7-13. [[Crossref](#)]
1028. Stine Linding Andersen. 2019. Frequency and outcomes of maternal thyroid function abnormalities in early pregnancy. *Scandinavian Journal of Clinical and Laboratory Investigation* **79**:1-2, 99-107. [[Crossref](#)]
1029. Lee Sun Y.. 2019. Maternal Iodine Deficiency in Pregnancy Is Associated with Decreased Language Skills in Infancy and Toddlerhood. *Clinical Thyroidology* **31**:2, 55-57. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1030. Emerson Charles H.. 2019. Iodine Content Is Low or Absent in Some U.S. Multivitamin and Prenatal Vitamin Brands. *Clinical Thyroidology* **31**:2, 58-60. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]

1031. Razvi Salman, Korevaar Tim I.M., Taylor Peter. 2019. Trends, Determinants, and Associations of Treated Hypothyroidism in the United Kingdom, 2005–2014. *Thyroid* **29**:2, 174–182. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
1032. Dong Allan Chen, Stagnaro-Green Alex. 2019. Differences in Diagnostic Criteria Mask the True Prevalence of Thyroid Disease in Pregnancy: A Systematic Review and Meta-Analysis. *Thyroid* **29**:2, 278–289. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
1033. Alessandro P. Delitala, Giampiero Capobianco, Pier Luigi Cherchi, Salvatore Dessole, Giuseppe Delitala. 2019. Thyroid function and thyroid disorders during pregnancy: a review and care pathway. *Archives of Gynecology and Obstetrics* **299**:2, 327–338. [[Crossref](#)]
1034. Jürgen M. Weiss. 2019. Schilddrüseneinstellung bei Kinderwunsch und in der Schwangerschaft. *Gynäkologische Endokrinologie* **17**:1, 20–24. [[Crossref](#)]
1035. Sergio Donnay Candil, Amelia Oleaga Alday, Elías Álvarez-García. 2019. Reference TSH levels in Spanish pregnant women. Is it possible to unify criteria?. *Endocrinología, Diabetes y Nutrición (English ed.)* **66**:2, 124–131. [[Crossref](#)]
1036. Sergio Donnay Candil, Amelia Oleaga Alday, Elías Álvarez-García. 2019. Valores de referencia de TSH en población gestante española. ¿Podemos unificar criterios?. *Endocrinología, Diabetes y Nutrición* **66**:2, 124–131. [[Crossref](#)]
1037. Whitney W. Woodmansee. 2019. Pituitary Disorders in Pregnancy. *Neurologic Clinics* **37**:1, 63–83. [[Crossref](#)]
1038. Hayden Anthony Homer. 2019. Modern management of recurrent miscarriage. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **59**:1, 36–44. [[Crossref](#)]
1039. Kate Wiles. 2019. Management for women with subclinical hypothyroidism in pregnancy. *Drug and Therapeutics Bulletin* **57**:2, 22–26. [[Crossref](#)]
1040. Mara Simopoulou, Konstantinos Sfakianoudis, Evangelos Maziotis, Sokratis Grigoriadis, Polina Giannelou, Anna Rapani, Petroula Tsioulou, Agni Pantou, Theodoros Kalampokas, Nikolaos Vlahos, Konstantinos Pantos, Michael Koutsilieris. 2019. The Impact of Autoantibodies on IVF Treatment and Outcome: A Systematic Review. *International Journal of Molecular Sciences* **20**:4, 892. [[Crossref](#)]
1041. Yoshihara Ai, Iwaku Kenji, Noh Jaeduk Yoshimura, Watanabe Natsuko, Kunii Yo, Ohye Hidemi, Suzuki Miho, Matsumoto Masako, Suzuki Nami, Tadokoro Rie, Sekiyama Chihiro, Hiruma Marino, Sugino Kiminori, Ito Koichi. 2019. Incidence of Neonatal Hyperthyroidism Among Newborns of Graves' Disease Patients Treated with Radioiodine Therapy. *Thyroid* **29**:1, 128–134. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1042. Patel Aneesh, Lee Sun Y., Stagnaro-Green Alex, MacKay Douglas, Wong Andrea W., Pearce Elizabeth N.. 2019. Iodine Content of the Best-Selling United States Adult and Prenatal Multivitamin Preparations. *Thyroid* **29**:1, 124–127. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1043. Andersen Stine Linding, Andersen Stig. 2019. Timing of Shift in Antithyroid Drug Therapy and Birth Defects. *Thyroid* **29**:1, 155–156. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1044. Antonina A. Smetnik, Alexander N. Sencha, Stanislav V. Pavlovich. Thyroid Disorders and Female Reproductive System Diseases: The Thyroid Gland and Pregnancy 203–213. [[Crossref](#)]
1045. Malini Soundarajan, Peter A. Kopp. Thyroid Hormone Biosynthesis and Physiology 1–17. [[Crossref](#)]
1046. Kelly S. Acharya, Jennifer L. Eaton. Thyroid Autoimmunity and Miscarriage 169–182. [[Crossref](#)]
1047. Benjamin S. Harris, Jennifer L. Eaton. Postpartum Thyroiditis 183–188. [[Crossref](#)]
1048. Nathan King, Lia A. Bernardi. Thyroid Function and Pregnancy 69–78. [[Crossref](#)]
1049. Elizabeth N. Pearce. Hypothyroidism in Pregnancy 101–115. [[Crossref](#)]
1050. Wilburn D. Bolton, Jennifer M. Perkins. Thyrotoxicosis in Pregnancy 117–136. [[Crossref](#)]
1051. Sarah E. Mayson, Linda A. Barbour. Thyroid Nodules and Cancer in Pregnancy 137–156. [[Crossref](#)]
1052. Shweta J. Bhatt, Emily C. Holden, Aimee Seungdamrong. Thyroid Dysfunction and Infertility 157–168. [[Crossref](#)]
1053. Raluca-Alexandra Trifanescu, Catalina Poiana. Diagnosis and Management of Hypothyroidism in Adults 581–592. [[Crossref](#)]
1054. Dev A. Kevat, Lucy Mackillop. Thyroid Disease in Pre- and Post-Pregnancy 629–641. [[Crossref](#)]
1055. Sofie Bliddal, Henriette S. Nielsen, Ulla Feldt-Rasmussen. Thyroid and Infertility 752–761. [[Crossref](#)]
1056. 2019. Prepregnancy counseling. *Fertility and Sterility* **111**:1, 32–42. [[Crossref](#)]
1057. D. Álvarez Ballano, M.O. Bandrés Nivelá, M.L. Gracia Ruiz, A. Ilundain González, P. de Diego García, Y. Blasco Lamarca, A.B. Mañas Martínez, E. García-González, L. Rello Varas, M.A. Sancho Serrano, J.J. Puente Lanzarote. 2019. Intervalos de referencia de hormonas tiroideas en mujeres gestantes mediante 2 inmunoanálisis diferentes: la importancia del método por encima de valores únicos universales, en consonancia con las recomendaciones internacionales 2017. *Clínica e Investigación en Ginecología y Obstetricia* **46**:1, 21–27. [[Crossref](#)]

1058. C C Repelaer van Driel- Delprat, E W C M van Dam, P M van de Ven, S Homsma, L van der Kooij, E Vis, R P Peeters, R Schats, C B Lambalk. 2019. Live birth rate after intrauterine insemination is not different between women with lower quartile versus higher quartile normal range thyroid stimulating hormone levels. *Human Reproduction Open* **2019**:1. . [\[Crossref\]](#)
1059. 2019. ACOG Committee Opinion No. 762: Prepregnancy Counseling. *Obstetrics & Gynecology* **133**:1, e78-e89. [\[Crossref\]](#)
1060. Song-Qing Deng, Hai-Tian Chen, Dong-Yu Wang, Bin Liu, Han-Qing Chen, Zi-Lian Wang. 2019. Maternal Thyroid-Stimulating Hormone Level and Thyroid Peroxidase Antibody Status in the First and Second Trimester of Pregnancy and Their Relationship with the Risk of Gestational Diabetes Mellitus. *Maternal-Fetal Medicine* **1**:2, 81-85. [\[Crossref\]](#)
1061. Daowen Zhang, Keying Cai, Guixia Wang, Shuhang Xu, Xiaodong Mao, Ang Zheng, Chao Liu, Kuanlu Fan. 2019. Trimester-specific reference ranges for thyroid hormones in pregnant women. *Medicine* **98**:4, e14245. [\[Crossref\]](#)
1062. Josef Köhrle. 2019. The Colorful Diversity of Thyroid Hormone Metabolites. *European Thyroid Journal* **8**:3, 115-129. [\[Crossref\]](#)
1063. Diogo Mendes, Carlos Alves, Nuno Silverio, Francisco Batel Marques. 2019. Prevalence of Undiagnosed Hypothyroidism in Europe: A Systematic Review and Meta-Analysis. *European Thyroid Journal* **8**:3, 130-143. [\[Crossref\]](#)
1064. Ilaria Muller, Carla Moran, Beatriz Lecumberri, Brigitte Decallonne, Neil Robertson, Joanne Jones, Colin M. Dayan. 2019. 2019 European Thyroid Association Guidelines on the Management of Thyroid Dysfunction following Immune Reconstitution Therapy. *European Thyroid Journal* **8**:4, 173-185. [\[Crossref\]](#)
1065. Ako Oiwa, Kesami Minemura, Shin-ichi Nishio, Masanori Yamazaki, Mitsuhsa Komatsu. 2019. Implications of thyroid autoimmunity in infertile women with subclinical hypothyroidism in the absence of both goiter and anti-thyroid antibodies: lessons from three cases. *Endocrine Journal* **66**:2, 193-198. [\[Crossref\]](#)
1066. Florina Păuleț, Alexandru Baroș, Crenguța Șerboiu, Monica Mihaela Cîrstoiu. 2019. Ultrasound findings of hypothyroidism during pregnancy. *Ginecologia.ro* **4**:26, 15. [\[Crossref\]](#)
1067. Shui-Boon Soh, Tar-Choon Aw. 2019. Laboratory Testing in Thyroid Conditions - Pitfalls and Clinical Utility. *Annals of Laboratory Medicine* **39**:1, 3. [\[Crossref\]](#)
1068. Hyemi Kwon. 2019. Letter: Thyroid-Stimulating Hormone Reference Ranges in the First Trimester of Pregnancy in an Iodine-Sufficient Country (Endocrinol Metab 2018;33:466-72, Carmen Castillo et al.). *Endocrinology and Metabolism* **34**:1, 93. [\[Crossref\]](#)
1069. Lorena Mosso. 2019. Response: Thyroid-Stimulating Hormone Reference Ranges in the First Trimester of Pregnancy in an Iodine-Sufficient Country (Endocrinol Metab 2018;33:466-72, Carmen Castillo et al.). *Endocrinology and Metabolism* **34**:2, 213. [\[Crossref\]](#)
1070. Elizabeth N. Pearce. 2019. Management of Thyrotoxicosis: Preconception, Pregnancy, and the Postpartum Period. *Endocrine Practice* **25**:1, 62-68. [\[Crossref\]](#)
1071. Valentin V. Fadeyev. 2018. Review of American Thyroid Association guidelines for the diagnosis and management of thyroid disease during pregnancy and the postpartum. *Clinical and experimental thyroidology* **14**:3, 128-139. [\[Crossref\]](#)
1072. Qi Zhou, Yanli Zhang, Jianhua Zhou, Xiangrong Yang, Yuqiang Huang, Huafeng Li, Chunling Ma. 2018. Analysis of detection results of thyroid function-related indexes in pregnant women and establishment of the reference interval. *Experimental and Therapeutic Medicine* . [\[Crossref\]](#)
1073. Stephanie Allelein, Matthias Schott. 2018. Schilddrüse und Schwangerschaft. *CME* **15**:12, 9-18. [\[Crossref\]](#)
1074. Frédéric Illouz, Dominique Luton, Michel Polak, Alix Besançon, Claire Bornaud. 2018. Graves' disease and pregnancy. *Annales d'Endocrinologie* **79**:6, 636-646. [\[Crossref\]](#)
1075. Harry Gibson, Catherine Nelson-Piercy. 2018. Endocrine disorders in pregnancy. *Medicine* **46**:12, 738-742. [\[Crossref\]](#)
1076. Ping Li, Shuo Lin, Ling Li, Jinhui Cui, Shuisheng Zhou, Jianhui Fan. 2018. Effect of mildly elevated thyroid-stimulating hormone during the first trimester on adverse pregnancy outcomes. *BMC Endocrine Disorders* **18**:1. . [\[Crossref\]](#)
1077. Gorkem Tuncay, Abdullah Karaer, Ebru İnci Coşkun, Demet Baloglu, Ayşe Nihan Tecellioglu. 2018. The impact of thyroid-stimulating hormone levels in euthyroid women on intrauterine insemination outcome. *BMC Women's Health* **18**:1. . [\[Crossref\]](#)
1078. Julie Bienertová-Vášků, Markéta Grulichová, Ondřej Mikeš, Filip Zlámál, Tomáš Pruša, Aneta Pohořalá, Lenka Andrášková, Hyněk Pikhart. 2018. Estimated dietary iodine intake as a predictor of placental size: evidence from the ELSPAC study. *Nutrition & Metabolism* **15**:1. . [\[Crossref\]](#)
1079. Tim I. M. Korevaar. 2018. The upper limit for TSH during pregnancy: why we should stop using fixed limits of 2.5 or 3.0 mU/l. *Thyroid Research* **11**:1. . [\[Crossref\]](#)
1080. Kris Poppe, Flora Veltri, Candice Autin. 2018. 'Does levothyroxine improve pregnancy outcomes in euthyroid women with thyroid autoimmunity undergoing assisted reproductive technology?'. *Thyroid Research* **11**:1. . [\[Crossref\]](#)

1081. Caroline T. Nguyen, Elizabeth B. Sasso, Lorayne Barton, Jorge H. Mestman. 2018. Graves' hyperthyroidism in pregnancy: a clinical review. *Clinical Diabetes and Endocrinology* 4:1. . [\[Crossref\]](#)
1082. Violeta Mladenovic. 2018. Thyroid and Pregnancy. *Serbian Journal of Experimental and Clinical Research*, ahead of print. [\[Crossref\]](#)
1083. H. Lewis, R. Egerman, A. Kazory, M. Sattari, C. O. Шурпак. 2018. Diabetes and Pregnancy: Risks and Opportunities. *Family Medicine* 0:5, 53-59. [\[Crossref\]](#)
1084. Ivana Chandra, Lizhou Sun. 2018. Preterm and term preeclampsia: differences in biochemical parameter and pregnancy outcomes. *Postgraduate Medicine* 130:8, 703-707. [\[Crossref\]](#)
1085. Yonghong Sheng, Dongping Huang, Shun Liu, Xuefeng Guo, Jiehua Chen, Yantao Shao, Guoqiang Zhang, Liangjia Wei, Xiaoyun Zeng, Xiaoqiang Qiu. 2018. Reference Intervals of Thyroid Hormones and Correlation of BMI with Thyroid Function in Healthy Zhuang Ethnic Pregnant Women. *BioMed Research International* 2018, 1-8. [\[Crossref\]](#)
1086. Pearce Elizabeth N.. 2018. Levothyroxine Treatment Improves Miscarriage Risk, But Not Live Birth Rates, in Women with Subclinical Hypothyroidism or TPO Antibody Positivity Undergoing ART. *Clinical Thyroidology* 30:11, 526-528. [\[Citation\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
1087. Nazeri Pantea, Tahmasebinejad Zhale, Mehrabi Yadollah, Hedayati Mehdi, Mirmiran Parvin, Azizi Fereidoun. 2018. Lactating Mothers and Infants Residing in an Area with an Effective Salt Iodization Program Have No Need for Iodine Supplements: Results from a Double-Blind, Placebo-Controlled, Randomized Controlled Trial. *Thyroid* 28:11, 1547-1558. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
1088. Magdy Mohamed Allam, Hanaa Tarek Hussein Elzawawy. 2018. Induction of remission in autoimmune polyglandular syndrome type three (APS III): An old drug with new perspectives. *Clinical Case Reports* 6:11, 2178-2184. [\[Crossref\]](#)
1089. Patricia C. Weissenfels, Joachim Woelfle, Eckhard Korsch, Matthias Joergens, Bettina Gohlke. 2018. Inconsistencies in the management of neonates born to mothers with "thyroid diseases". *European Journal of Pediatrics* 177:11, 1711-1718. [\[Crossref\]](#)
1090. Tian Chen, Piye Niu, Fanling Kong, Yuwei Wang, Yi Bai, Dong Yu, Jiaxin Jia, Luping Yang, Zhongjian Fu, Renbo Li, Jingguang Li, Lin Tian, Zhiwei Sun, Dejun Wang, Zhixiong Shi. 2018. Disruption of thyroid hormone levels by decabrominated diphenyl ethers (BDE-209) in occupational workers from a deca-BDE manufacturing plant. *Environment International* 120, 505-515. [\[Crossref\]](#)
1091. Aly Bernard Khalil, Bashir Taha Salih, Onismos Chinengo, Ma Remy D. Bardies, Andrew Turner, Laila O. Abdel Wareth. 2018. Trimester specific reference ranges for serum TSH and Free T4 among United Arab Emirates pregnant women. *Practical Laboratory Medicine* 12, e00098. [\[Crossref\]](#)
1092. Helle Katrine Knutsen, Jan Alexander, Lars Barregård, Margherita Bignami, Beat Brüsweiler, Sandra Ceccatelli, Bruce Cottrill, Michael Dinovi, Lutz Edler, Bettina Grasl-Kraupp, Christer Hogstrand, Carlo Stefano Nebbia, Isabelle P Oswald, Annette Petersen, Martin Rose, Alain-Claude Roudot, Tanja Schwerdtle, Christiane Vleminckx, Günter Vollmer, Heather Wallace, Peter Fürst, Helen Håkansson, Thorhallur Halldorsson, Anne-Katrine Lundebye, Raimo Pohjanvirta, Lars Rylander, Andrew Smith, Henk van Loveren, Ine Waalkens-Berendsen, Marco Zeilmaker, Marco Binaglia, José Ángel Gómez Ruiz, Zsuzsanna Horváth, Eugen Christoph, Laura Ciccolallo, Luisa Ramos Bordajandi, Hans Steinkellner, Laurentius (Ron) Hoogenboom. 2018. Risk for animal and human health related to the presence of dioxins and dioxin-like PCBs in feed and food. *EFSA Journal* 16:11. . [\[Crossref\]](#)
1093. Peter N. Taylor, Stamatios Zouras, Thinzar Min, Kalyani Nagarajah, John H. Lazarus, Onyebuchi Okosieme. 2018. Thyroid Screening in Early Pregnancy: Pros and Cons. *Frontiers in Endocrinology* 9. . [\[Crossref\]](#)
1094. Yang Liu, Lin Li, Jie Yu, Yu-Xia Fan, Xiu-Bo Lu. 2018. Carbon nanoparticle lymph node tracer improves the outcomes of surgical treatment in papillary thyroid cancer. *Cancer Biomarkers* 23:2, 227-233. [\[Crossref\]](#)
1095. Lijie Zhao, Guojing Jiang, Xingfang Tian, Xiaoqin Zhang, Ting Zhu, Beibei Chen, Yongxiang Wang, Qingliang Ma. 2018. Initiation timing effect of levothyroxine treatment on subclinical hypothyroidism in pregnancy. *Gynecological Endocrinology* 34:10, 845-848. [\[Crossref\]](#)
1096. Hattori Naoki, Aisaka Kohzo, Chihara Kazuhisa, Shimatsu Akira. 2018. Current Thyrotropin Immunoassays Recognize Macro-Thyrotropin Leading to Hyperthyrotropinemia in Females of Reproductive Age. *Thyroid* 28:10, 1252-1260. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#)
1097. Abel Marianne Hope, Korevaar Tim I.M., Erlund Iris, Villanger Gro Dehli, Caspersen Ida Henriette, Arohonka Petra, Alexander Jan, Meltzer Helle Margrete, Brantsæter Anne Lise. 2018. Iodine Intake is Associated with Thyroid Function in Mild to Moderately Iodine Deficient Pregnant Women. *Thyroid* 28:10, 1359-1371. [\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#) [\[PDF Plus\]](#) [\[Supplementary Material\]](#)
1098. Sanjay Kalra, A. K. Das, Sarita Bajaj, Banshi Saboo, Deepak Khandelwal, Mangesh Tiwaskar, Navneet Agarwal, Pritam Gupta, Rakesh Sahay, Sameer Aggarwal, Sujoy Ghosh, Vijay Negalur, A. G. Unnikrishnan, Ganapathi Bantwal, Rashmi Aggarwal,

- Harshal Chaudhari, Nitin Mulgaonkar. 2018. Diagnosis and Management of Hypothyroidism: Addressing the Knowledge–Action Gaps. *Advances in Therapy* **35**:10, 1519–1534. [[Crossref](#)]
1099. Martin P. Hansen, George J. Kahaly. 2018. Hypothyreose beim Hausarzt: wie abklären, wie behandeln?. *MMW - Fortschritte der Medizin* **160**:17, 42–46. [[Crossref](#)]
1100. Susana Ares Segura, Carmen Temboury Molina, María Jesús Chueca Guindulain, Gema Grau Bolado, María Jesus Alija Merillas, María Caimari Jaume, Paula Casano Sancho, José Carlos Moreno Navarro, José Manuel Rial Rodríguez, Amparo Rodríguez Sánchez. 2018. Recommendations for the diagnosis and followup of the foetus and newborn child born to mothers with autoimmune thyroid disease. *Anales de Pediatría (English Edition)* **89**:4, 254.e1–254.e7. [[Crossref](#)]
1101. Susana Ares Segura, Carmen Temboury Molina, María Jesús Chueca Guindulain, Gema Grau Bolado, María Jesus Alija Merillas, María Caimari Jaume, Paula Casano Sancho, José Carlos Moreno Navarro, José Manuel Rial Rodríguez, Amparo Rodríguez Sánchez. 2018. Recomendaciones para el diagnóstico y seguimiento del feto y del recién nacido hijo de madre con patología tiroidea autoinmune. *Anales de Pediatría* **89**:4, 254.e1–254.e7. [[Crossref](#)]
1102. Paula Fernández Martínez, Rocío Aguado García, David Emilio Barajas Galindo, Ana Hernández Moreno, Mirian Alejo Ramos, Sara García Arias, María D. Ballesteros Pomar, Isidoro Manuel Cano Rodríguez. 2018. Influence of thyroid peroxidase antibodies on TSH levels of pregnant women and maternal–fetal complications. *Endocrinología, Diabetes y Nutrición (English ed.)* **65**:8, 444–450. [[Crossref](#)]
1103. Paula Fernández Martínez, Rocío Aguado García, David Emilio Barajas Galindo, Ana Hernández Moreno, Mirian Alejo Ramos, Sara García Arias, María D. Ballesteros Pomar, Isidoro Manuel Cano Rodríguez. 2018. Influencia de los anticuerpos antiperoxidasa tiroidea en los valores de TSH de gestantes y en las complicaciones materno-fetales. *Endocrinología, Diabetes y Nutrición* **65**:8, 444–450. [[Crossref](#)]
1104. Luis Javier Morales-García, María Pacheco-Delgado, Santiago Prieto Menchero, Daniel Pineda Tenor. 2018. Influencia del informe de laboratorio en el diagnóstico de la disfunción tiroidea en la gestante: más allá del asterisco. *Revista del Laboratorio Clínico* **11**:4, 186–192. [[Crossref](#)]
1105. Luba Rakhlin, Stephanie Fish. 2018. Pregnancy as a risk factor for thyroid cancer progression. *Current Opinion in Endocrinology & Diabetes and Obesity* **25**:5, 326–329. [[Crossref](#)]
1106. Arash Derakhshan, Tim I M Korevaar, Peter N Taylor, Deborah Levie, Monica Guxens, Vincent W V Jaddoe, Scott M Nelson, Henning Tiemeier, Robin P Peeters. 2018. The Association of Maternal Thyroid Autoimmunity During Pregnancy With Child IQ. *The Journal of Clinical Endocrinology & Metabolism* **103**:10, 3729–3736. [[Crossref](#)]
1107. Halenur Bozdog, Esra Akdeniz. 2018. Does severe vitamin D deficiency impact obstetric outcomes in pregnant women with thyroid autoimmunity?. *The Journal of Maternal-Fetal & Neonatal Medicine* **122**, 1–11. [[Crossref](#)]
1108. Farebrother Jessica, Zimmermann Michael B., Abdallah Fatma, Assey Vincent, Fingerhut Ralph, Gichohi-Wainaina Wanjiku N., Hussein Izzeldin, Makokha Anselimo, Sagno Kalil, Untoro Juliawati, Watts Michael, Andersson Maria. 2018. Effect of Excess Iodine Intake from Iodized Salt and/or Groundwater Iodine on Thyroid Function in Nonpregnant and Pregnant Women, Infants, and Children: A Multicenter Study in East Africa. *Thyroid* **28**:9, 1198–1210. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
1109. Georg Zettinig, Wolfgang Buchinger, Corinna Mann. 2018. Die interdisziplinäre Schilddrüse – Fertilität und Schwangerschaft. *Journal für Gynäkologische Endokrinologie/Schweiz* **21**:3, 108–111. [[Crossref](#)]
1110. Yung-Hsiang Lin, Ko-Jung Chen, Yun-Shing Peng, Pau-Chung Chen, Yao-Hsu Yang. 2018. Type 1 diabetes impairs female fertility even before it is diagnosed. *Diabetes Research and Clinical Practice* **143**, 151–158. [[Crossref](#)]
1111. Matthew T. Drake. 2018. Hypothyroidism in Clinical Practice. *Mayo Clinic Proceedings* **93**:9, 1169–1172. [[Crossref](#)]
1112. Gesthimani Mintziori, Dimitrios G. Goulis. 2018. In vitro fertilization/intracytoplasmic insemination and thyroid function: reviewing the evidence. *Metabolism* **86**, 44–48. [[Crossref](#)]
1113. Débora Ayres Saraiva, Nathalie Anne de Oliveira e Silva de Moraes, Carolina Martins Corcino, Tatiana Martins Benvenuto Louro Berbara, Annie Schtscherbyna, Mischelle Santos, Heron Botelho, Mario Vaisman, Patrícia de Fátima dos Santos Teixeira. 2018. Iodine status of pregnant women from a coastal Brazilian state after the reduction in recommended iodine concentration in table salt according to governmental requirements. *Nutrition* **53**, 109–114. [[Crossref](#)]
1114. Sara Watutantrige-Fernando, Susi Barollo, Loris Bertazza, Elisabetta Cavedon, Simona Censi, Jacopo Manso, Federica Vianello, Caterina Mian, Sabrina Andrighettoni, Andrea Maria Chiamenti, Saverio Chilese, Monica Cibir, Oscar Cora, Lucia Cortese, Fiorella Costa, Rita De Noni, Marianna Elia, Giuseppina Girlando, Sibilla Levorato, Renato Ranieri, Valeria Rebonato, Giovanni Sandri, Riccardo Sciacco, Mauro Soppelsa, Antonio Stano, Stefania Tessari, Angiola Vanzo, Laura Zadra, Lorena Zambelli. 2018. Efficacy of educational intervention to improve awareness of the importance of iodine, use of iodized salt, and dietary iodine intake in northeastern Italian schoolchildren. *Nutrition* **53**, 134–139. [[Crossref](#)]

1115. A. Ortega Carpio, I. Vázquez Rico, M.A. Castaño López, L. Duarte González, M. Montilla Álvaro, A. Ruiz Reina. 2018. Intervalos de referencia de la tirotropina durante la gestación en la provincia de Huelva. *Medicina de Familia. SEMERGEN* **44**:6, 372-379. [[Crossref](#)]
1116. Onyebuchi E. Okosieme, Ishrat Khan, Peter N. Taylor. 2018. Preconception management of thyroid dysfunction. *Clinical Endocrinology* **89**:3, 269-279. [[Crossref](#)]
1117. Jennifer M Yamamoto, Jamie L Benham, Kara A Nerenberg, Lois E Donovan. 2018. Impact of levothyroxine therapy on obstetric, neonatal and childhood outcomes in women with subclinical hypothyroidism diagnosed in pregnancy: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* **8**:9, e022837. [[Crossref](#)]
1118. Arash Derakhshan, Huan Shu, Maarten A C Broeren, Ralph A de Poortere, Sverre Wikström, Robin P Peeters, Barbara Demeneix, Carl-Gustaf Bornehag, Tim I M Korevaar. 2018. Reference Ranges and Determinants of Thyroid Function During Early Pregnancy: The SELMA Study. *The Journal of Clinical Endocrinology & Metabolism* **103**:9, 3548-3556. [[Crossref](#)]
1119. Songlin Yu, Yicong Yin, Qian Cheng, Jianhua Han, Xinqi Cheng, Ye Guo, Dandan Sun, Shaowei Xie, Ling Qiu. 2018. Validation of a simple inductively coupled plasma mass spectrometry method for detecting urine and serum iodine and evaluation of iodine status of pregnant women in Beijing. *Scandinavian Journal of Clinical and Laboratory Investigation* **78**:6, 501-507. [[Crossref](#)]
1120. Xiaotong Gao, Yongze Li, Jiashu Li, Aihua Liu, Wei Sun, Weiping Teng, Zhongyan Shan. 2018. Gestational TSH and FT4 Reference Intervals in Chinese Women: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology* **9**. [[Crossref](#)]
1121. Teng Xiaochun, Shan Zhongyan, Li Chenyan, Yu Xiaohui, Mao Jinyuan, Wang Weiwei, Xie Xiaochen, Du Jianling, Zhang Shaowei, Gao Zhengnan, Zhang Xiaomei, Li Ling, Fan Chenling, Teng Weiping. 2018. Iron Deficiency May Predict Greater Risk for Hypothyroxinemia: A Retrospective Cohort Study of Pregnant Women in China. *Thyroid* **28**:8, 968-975. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
1122. Shirley Cueva, Channing Burks, Dana McQueen, Marla S. Barkoff, Mary D. Stephenson. 2018. Maternal antithyroid antibodies and euploid miscarriage in women with recurrent early pregnancy loss. *Fertility and Sterility* **110**:3, 452-458. [[Crossref](#)]
1123. Giovanna Mantovani, Murat Bastepe, David Monk, Luisa de Sanctis, Susanne Thiele, Alessia Usardi, S. Faisal Ahmed, Roberto Bufo, Timothée Choplin, Gianpaolo De Filippo, Guillemette Devernois, Thomas Eggermann, Francesca M. Elli, Kathleen Freson, Aurora García Ramirez, Emily L. Germain-Lee, Lionel Groussin, Neveen Hamdy, Patrick Hanna, Olaf Hiort, Harald Jüppner, Peter Kamenický, Nina Knight, Marie-Laure Kottler, Elvire Le Norcy, Beatriz Lecumberri, Michael A. Levine, Outi Mäkitie, Regina Martin, Gabriel Ángel Martos-Moreno, Masanori Minagawa, Philip Murray, Arrate Pereda, Robert Pignolo, Lars Rejnmark, Rebecca Rodado, Anya Rothenbuhler, Vrinda Saraff, Ashley H. Shoemaker, Eileen M. Shore, Caroline Silve, Serap Turan, Philip Woods, M. Carola Zillikens, Guiomar Perez de Nanclares, Agnès Linglart. 2018. Diagnosis and management of pseudohypoparathyroidism and related disorders: first international Consensus Statement. *Nature Reviews Endocrinology* **14**:8, 476-500. [[Crossref](#)]
1124. Victor Pop, Maarten Broeren, Hennie Wijnen, Joyce Endendijk, Anneloes van Baar, Wilmar Wiersinga, Graham R Williams. 2018. Longitudinal Trajectories of Gestational Thyroid Function: A New Approach to Better Understand Changes in Thyroid Function. *The Journal of Clinical Endocrinology & Metabolism* **103**:8, 2889-2900. [[Crossref](#)]
1125. Deborah Levie, Tim I M Korevaar, Sarah C Bath, Albert Dalmau-Bueno, Mario Murcia, Mercedes Espada, Mariana Dineva, Jesús M Ibarluzea, Jordi Sunyer, Henning Tiemeier, Marisa Rebagliato, Margaret P Rayman, Robin P Peeters, Mònica Guxens. 2018. Thyroid Function in Early Pregnancy, Child IQ, and Autistic Traits: A Meta-Analysis of Individual Participant Data. *The Journal of Clinical Endocrinology & Metabolism* **103**:8, 2967-2979. [[Crossref](#)]
1126. Nathalie Anne de Oliveira e Silva de Moraes, Annie Schtscherbyna Almeida de Assis, Carolina Martins Corcino, Débora Ayres Saraiva, Tatiana Martins Benvenuto Louro Berbara, Carolina Donner de Drummond Ventura, Mario Vaisman, Patrícia de Fátima dos Santos Teixeira. 2018. Recent recommendations from ATA guidelines to define the upper reference range for serum TSH in the first trimester match reference ranges for pregnant women in Rio de Janeiro. *Archives of Endocrinology and Metabolism* **62**:4, 386-391. [[Crossref](#)]
1127. José Augusto Sgarbi. 2018. Region-specific reference intervals for TSH in pregnancy: time for changes in Brazil. *Archives of Endocrinology and Metabolism* **62**:4, 383-385. [[Crossref](#)]
1128. Chien-Yu Lin, Cheng-Kai Li, Jui-Ming Liu, Ren-Jun Hsu, Heng-Chang Chuang, Fung-Wei Chang. 2018. Postpartum Depression and Subsequent Autoimmune Diseases in Taiwan. *International Journal of Environmental Research and Public Health* **15**:8, 1783. [[Crossref](#)]
1129. Sima Nazarpour, Fahimeh Ramezani Tehrani, Maryam Rahmati, Sonia Minooee, Masoumeh Simbar, Mahsa Noroozadeh, Fereidoun Azizi. 2018. Validation of Billewicz Scoring System for Detection of Overt Hypothyroidism During Pregnancy. *International Journal of Endocrinology and Metabolism* **16**:3. [[Crossref](#)]
1130. Steve L Hyer. 2018. Caring for women with thyroid disorders in pregnancy. *British Journal of Midwifery* **26**:7, 434-439. [[Crossref](#)]

1131. Pearce Elizabeth N.. 2018. Higher Preconception Maternal Iodine Intakes Are Associated with Higher Child IQ. *Clinical Thyroidology* **30**:7, 302-304. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1132. Tomáš Šálek, Ishraq Dhaifalah, Dagmar Langova, Jana Havalová. 2018. Maternal thyroid-stimulating hormone reference ranges for first trimester screening from 11 to 14 weeks of gestation. *Journal of Clinical Laboratory Analysis* **32**:6, e22405. [[Crossref](#)]
1133. K. A. Iwen, H. Lehnert. 2018. Schilddrüse und Schwangerschaft. *Der Internist* **59**:7, 654-660. [[Crossref](#)]
1134. Simone De Leo, Elizabeth N Pearce. 2018. Autoimmune thyroid disease during pregnancy. *The Lancet Diabetes & Endocrinology* **6**:7, 575-586. [[Crossref](#)]
1135. Anupam Kotwal, Marius Stan. 2018. Thyrotropin Receptor Antibodies—An Overview. *Ophthalmic Plastic & Reconstructive Surgery* **34**:4S, S20-S27. [[Crossref](#)]
1136. Shruti Rawal, Michael Y Tsai, Stefanie N Hinkle, Yeyi Zhu, Wei Bao, Yuan Lin, Pranati Panuganti, Paul S Albert, Ronald C W Ma, Cuilin Zhang. 2018. A Longitudinal Study of Thyroid Markers Across Pregnancy and the Risk of Gestational Diabetes. *The Journal of Clinical Endocrinology & Metabolism* **103**:7, 2447-2456. [[Crossref](#)]
1137. Howard B A Baum. 2018. Clinical Excellence in Endocrinology. *The Journal of Clinical Endocrinology & Metabolism* **103**:7, 2430-2435. [[Crossref](#)]
1138. Wieland Kiess, Melanie Penke, Julia Gesing, Annette Stoltze, Antje Körner, Roland Pfäffle, Jürgen Kratzsch. 2018. Congenital hypothyroidism. *Journal of Pediatric Endocrinology and Metabolism* **31**:6, 595-596. [[Crossref](#)]
1139. Sawka Anna M., Carty Sally E., Haugen Bryan R., Hennessey James V., Kopp Peter A., Pearce Elizabeth N., Sosa Julie A., Tufano Ralph P., Jonklaas Jacqueline. 2018. American Thyroid Association Guidelines and Statements: Past, Present, and Future. *Thyroid* **28**:6, 692-706. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1140. Jan Boucek, Jorine Haan, Michael J. Halaska, Jan Plzak, Kristel Calsteren, Christianne J. M. Groot, Karina Dahl Steffensen, Robert Fruscio, Elske T. Massolt, Philipp Klaritsch, Paolo Zola, Frédéric Amant. 2018. Maternal and obstetrical outcome in 35 cases of well-differentiated thyroid carcinoma during pregnancy. *The Laryngoscope* **128**:6, 1493-1500. [[Crossref](#)]
1141. Brigitte Decallonne, Emmanuel Bartholomé, Valérie Delvaux, Miguel D'haeseleer, Souraya El Sankari, Pierrette Seeldrayers, Bart Van Wijmeersch, Chantal Daumerie. 2018. Thyroid disorders in alemtuzumab-treated multiple sclerosis patients: a Belgian consensus on diagnosis and management. *Acta Neurologica Belgica* **118**:2, 153-159. [[Crossref](#)]
1142. Georg Zettinig, Wolfgang Buchinger, Corinna Mann. 2018. Die interdisziplinäre Schilddrüse – Fertilität und Schwangerschaft. *Journal für Gynäkologische Endokrinologie/Österreich* **28**:2, 62-65. [[Crossref](#)]
1143. Faiza Lamine, Sara De Giorgi, Laura Marino, Marina Michalaki, Gerasimos P. Sykiotis. 2018. Subclinical hypothyroidism: new trials, old caveats. *Hormones* **17**:2, 231-236. [[Crossref](#)]
1144. Maïa Banigé, Michel Polak, Dominique Luton, Alexandra Benachi, Valérie Biran, Mostafa Mokhtari, Claire Boithias, Valérie Champion, Luc Desfrere, Yves Ville, Pierre-Henri Jarreau, Marc Dommergues, Frederic Brioude, Laurent Mandelbrot, Delphine Mitanchez. 2018. Prediction of Neonatal Hyperthyroidism. *The Journal of Pediatrics* **197**, 249-254.e1. [[Crossref](#)]
1145. V. Bergink, V. J. M. Pop, P. R. Nielsen, E. Agerbo, T. Munk-Olsen, X. Liu. 2018. Comorbidity of autoimmune thyroid disorders and psychiatric disorders during the postpartum period: a Danish nationwide register-based cohort study. *Psychological Medicine* **48**:8, 1291-1298. [[Crossref](#)]
1146. Tim I. M. Korevaar, Robin P. Peeters. 2018. Antithyroid drugs and congenital malformations. *Nature Reviews Endocrinology* **14**:6, 328-329. [[Crossref](#)]
1147. Spyridoula Maraka, Naykky M Singh Ospina, George Mastorakos, Derek T O'Keeffe. 2018. Subclinical Hypothyroidism in Women Planning Conception and During Pregnancy: Who Should Be Treated and How?. *Journal of the Endocrine Society* **2**:6, 533-546. [[Crossref](#)]
1148. Mario Rotondi, Valentina Capelli, Francesca Coperchini, Sara Pinto, Laura Croce, Massimo Tonacchera, Luca Chiovato. 2018. Post-partum and non-post-partum relapsing Graves' hyperthyroidism display different response to anti-thyroid drugs. *European Journal of Endocrinology* **178**:6, 589-594. [[Crossref](#)]
1149. Juneo F Silva, Natália M Ocarino, Rogéria Serakides. 2018. Thyroid hormones and female reproduction†. *Biology of Reproduction* **113**. . [[Crossref](#)]
1150. Pedro Wesley Rosario, Luis Fernando Faria Oliveira, Maria Regina Calsolari. 2018. Maternal hypothyroxinemia in the first trimester of gestation and association with obstetric and neonatal outcomes and iron deficiency: a prospective Brazilian study. *Archives of Endocrinology and Metabolism* . [[Crossref](#)]
1151. Pearce Elizabeth N.. 2018. Neonatal TSH <0.90 mIU/L Increases the Risk for Hyperthyroidism in Infants of Mothers with Graves' Disease. *Clinical Thyroidology* **30**:5, 217-219. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]

1152. Bedaiwy Mohamed A., Abdel-Rahman Mohamed Y., Tan Justin, AbdelHafez Faten F., Abdelkareem Amr O., Henry Drisana, Lisonkova Sarka, Hurd William W., Liu James H.. 2018. Clinical, Hormonal, and Metabolic Parameters in Women with Subclinical Hypothyroidism and Polycystic Ovary Syndrome: A Cross-Sectional Study. *Journal of Women's Health* **27**:5, 659-664. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1153. Rotondi Mario, Chiovato Luca, Pacini Furio, Bartalena Luigi, Vitti Paolo. 2018. Management of Subclinical Hypothyroidism in Pregnancy: A Comment from the Italian Society of Endocrinology and the Italian Thyroid Association to the 2017 American Thyroid Association Guidelines—"The Italian Way". *Thyroid* **28**:5, 551-555. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1154. Lican Han, Wei Zheng, Yanhong Zhai, Xin Xie, Jingnan Zhang, Shaozu Zhang, Zhen Zhao, Zheng Cao. 2018. Reference intervals of trimester-specific thyroid stimulating hormone and free thyroxine in Chinese women established by experimental and statistical methods. *Journal of Clinical Laboratory Analysis* **32**:4, e22344. [[Crossref](#)]
1155. Sidra Hanif, Asim Ilyas, Munir H. Shah. 2018. Statistical Evaluation of Trace Metals, TSH and T4 in Blood Serum of Thyroid Disease Patients in Comparison with Controls. *Biological Trace Element Research* **183**:1, 58-70. [[Crossref](#)]
1156. Joseph M. Braun, Aimin Chen, Andrew Hoofnagle, George D. Papandonatos, Medina Jackson-Browne, Russ Hauser, Megan E. Romano, Margaret R. Karagas, Kimberly Yoltan, R. Thomas Zoeller, Bruce P. Lanphear. 2018. Associations of early life urinary triclosan concentrations with maternal, neonatal, and child thyroid hormone levels. *Hormones and Behavior* **101**, 77-84. [[Crossref](#)]
1157. Peter N. Taylor, Diana Albrecht, Anna Scholz, Gala Gutierrez-Buey, John H. Lazarus, Colin M. Dayan, Onyebuchi E. Okosieme. 2018. Global epidemiology of hyperthyroidism and hypothyroidism. *Nature Reviews Endocrinology* **14**:5, 301-316. [[Crossref](#)]
1158. Marta Hernández, Carolina López, Berta Soldevila, Laura Cecenarro, María Martínez-Barahona, Elisabet Palomera, Ferran Rius, Albert Lecube, Maria José Pelegay, Jordi García, Dídac Mauricio, Manel Puig Domingo. 2018. Impact of TSH during the first trimester of pregnancy on obstetric and foetal complications: Usefulness of 2.5 mIU/L cut-off value. *Clinical Endocrinology* **88**:5, 728-734. [[Crossref](#)]
1159. Kris Poppe, Candice Autin, Flora Veltri, Pierre Kleynen, Lidia Grabczan, Serge Rozenberg, Lieveke Ameye. 2018. Thyroid Autoimmunity and Intracytoplasmic Sperm Injection Outcome: A Systematic Review and Meta-Analysis. *The Journal of Clinical Endocrinology & Metabolism* **103**:5, 1755-1766. [[Crossref](#)]
1160. Verônica Carneiro Borges Mito, Ana Carolina de Castro Nassif Gomes Monteiro, Rosalinda Yossie Asato de Camargo, Andréia Rodrigues Borel, Regina Maria Catarino, Sergio Kobayashi, Maria Cristina Chammas, Suemi Marui. 2018. High prevalence of iodine deficiency in pregnant women living in adequate iodine area. *Endocrine Connections* **7**:5, 762-767. [[Crossref](#)]
1161. Juliane Léger, Anne Forhan, Sophie Dos Santos, Béatrice Larroque, Emmanuel Ecosse, Marie-Aline Charles, Barbara Heude. 2018. Developmental milestones at one year for the offspring of mothers with congenital hypothyroidism: a population-based study. *European Journal of Endocrinology* **178**:5, 471-480. [[Crossref](#)]
1162. Kwanhoon Jo, Dong-Jun Lim. 2018. Wider reference ranges of serum thyroid stimulating hormone in pregnant Korean women: is it time to stop overtreatment?. *The Korean Journal of Internal Medicine* **33**:3, 493-496. [[Crossref](#)]
1163. Sakiko Kobayashi. 2018. V. Significance of Antihypertensive Therapy in Patients with Acromegaly or Thyroid Dysfunctions. *Nihon Naika Gakkai Zasshi* **107**:4, 688-695. [[Crossref](#)]
1164. Yingying Zhou, Xinyi Wang, Yuanyuan Zhang, Tong Zhao, Zhongyan Shan, Weiping Teng. 2018. Circulating MicroRNA Profile as a Potential Predictive Biomarker for Early Diagnosis of Spontaneous Abortion in Patients With Subclinical Hypothyroidism. *Frontiers in Endocrinology* **9**. . [[Crossref](#)]
1165. Fereidoun Azizi, Ladan Mehran, Farhad Hosseinpahan, Hossein Delshad, Atieh Amouzegar. 2018. Secondary and tertiary preventions of thyroid disease. *Endocrine Research* **43**:2, 124-140. [[Crossref](#)]
1166. Gideon Koren, Asher Ornoy. 2018. The role of the placenta in drug transport and fetal drug exposure. *Expert Review of Clinical Pharmacology* **11**:4, 373-385. [[Crossref](#)]
1167. Pearce Elizabeth N.. 2018. First-Trimester Maternal Thyroid Function Is Not Associated with Child Scores on Standardized Educational Tests. *Clinical Thyroidology* **30**:4, 179-181. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1168. Yuan Fang, Liang Yao, Jing Sun, Jian Zhang, Yanxia Li, Ruifei Yang, Kehu Yang, Limin Tian. 2018. Appraisal of clinical practice guidelines on the management of hypothyroidism in pregnancy using the Appraisal of Guidelines for Research and Evaluation II instrument. *Endocrine* **60**:1, 4-14. [[Crossref](#)]
1169. Susanne Dold, Michael B Zimmermann, Tomislav Jukic, Zvonko Kusic, Qingzhen Jia, Zhongna Sang, Antonio Quirino, Teofilo OL San Luis, Ralph Fingerhut, Roland Kupka, Arnold Timmer, Greg S Garrett, Maria Andersson. 2018. Universal Salt Iodization Provides Sufficient Dietary Iodine to Achieve Adequate Iodine Nutrition during the First 1000 Days: A Cross-Sectional Multicenter Study. *The Journal of Nutrition* **148**:4, 587-598. [[Crossref](#)]

1170. William Thompson, Ginny Russell, Genevieve Baragwanath, Justin Matthews, Bijay Vaidya, Jo Thompson-Coon. 2018. Maternal thyroid hormone insufficiency during pregnancy and risk of neurodevelopmental disorders in offspring: A systematic review and meta-analysis. *Clinical Endocrinology* **88**:4, 575-584. [[Crossref](#)]
1171. Lauren E Johns, Kelly K Ferguson, David E Cantonwine, Bhramar Mukherjee, John D Meeker, Thomas F McElrath. 2018. Subclinical Changes in Maternal Thyroid Function Parameters in Pregnancy and Fetal Growth. *The Journal of Clinical Endocrinology & Metabolism* **103**:4, 1349-1358. [[Crossref](#)]
1172. Priya Gupta, Jaime Gahche, Kirsten Herrick, Abby Ershow, Nancy Potischman, Cria Perrine. 2018. Use of Iodine-Containing Dietary Supplements Remains Low among Women of Reproductive Age in the United States: NHANES 2011–2014. *Nutrients* **10**:4, 422. [[Crossref](#)]
1173. Chang-Eun Park. 2018. Evaluation of Pregnancy and Thyroid Function. *The Korean Journal of Clinical Laboratory Science* **50**:1, 1-10. [[Crossref](#)]
1174. Anita Banerjee, Catherine Williamson. Endocrine and Metabolic Emergencies in Pregnancy 85-97. [[Crossref](#)]
1175. . Thyroid Disorders 243-251. [[Crossref](#)]
1176. Fanni Pääkkilä, Tuija Männistö, Anna-Liisa Hartikainen, Eila Suvanto. 2018. Maternal Thyroid Function During Pregnancy and the Child's Linguistic and Sensory Development in the Northern Finland Birth Cohort 1986. *Frontiers in Endocrinology* **9**. . [[Crossref](#)]
1177. Polina V. Popova, Ekaterina S. Shilova, Alexandra S. Tkachuk, Alexandra V. Dronova, Anna D. Anopova, Alla E. Nikolaeva, Elena N. Grineva. 2018. First trimester thyroid function in pregnant women residing in Saint Petersburg (Russia): reference values and risk of gestational diabetes. *Diabetes mellitus* **21**:1, 34-41. [[Crossref](#)]
1178. Spyridoula Maraka, Derek T. O'Keeffe, Naykky Singh Ospina. 2018. Commentary: Effect of Levothyroxine on Miscarriage among Women with Normal Thyroid Function and Thyroid Autoimmunity Undergoing In Vitro Fertilization and Embryo Transfer: A Randomized Clinical Trial. *Frontiers in Endocrinology* **9**. . [[Crossref](#)]
1179. Leung Angela M.. 2018. No Benefit of Levothyroxine Among Pregnant Hypothyroid and/or Hypothyroxinemic Women on Offspring IQ at Age 9 years. *Clinical Thyroidology* **30**:3, 100-103. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1180. Korevaar Tim I.M.. 2018. Methimazole Has a Dose-Dependent Association With Congenital Malformations, but Switching to PTU in the First Trimester Seems Too Late. *Clinical Thyroidology* **30**:3, 104-107. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1181. Pearce Elizabeth N.. 2018. Low Urinary Iodine Concentrations Predict Decreased Fertility. *Clinical Thyroidology* **30**:3, 135-137. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1182. K. Laubner, H. Kleinwechter. 2018. Diabetes mellitus Typ 1 und Schwangerschaft. *Der Diabetologe* **14**:2, 113-125. [[Crossref](#)]
1183. Valdelena Alessandra da Silva, Robson José de Almeida, Marcelle Pimenta Cavalcante, Lucio Antonio Pereira Junior, Francine Machado Reis, Mariana Fabbris Pereira, Teresa S. Kasamatsu, Cleber P. Camacho. 2018. Two Thyroid Stimulating Hormone assays correlated in clinical practice show disagreement in subclinical hypothyroidism patients. *Clinical Biochemistry* **53**, 13-18. [[Crossref](#)]
1184. Sima Nazarpour, Fahimeh Ramezani Tehrani, Masoumeh Simbar, Sonia Minooee, Maryam Rahmati, Mohammad Ali Mansournia, Fereidoun Azizi. 2018. Establishment of trimester-specific reference range for thyroid hormones during pregnancy. *Clinical Biochemistry* **53**, 49-54. [[Crossref](#)]
1185. Stephanie L. Samuels, Sisi M. Namoc, Andrew J. Bauer. 2018. Neonatal Thyrotoxicosis. *Clinics in Perinatology* **45**:1, 31-40. [[Crossref](#)]
1186. Monika Chaudhari, Jonathan L. Slaughter. 2018. Thyroid Function in the Neonatal Intensive Care Unit. *Clinics in Perinatology* **45**:1, 19-30. [[Crossref](#)]
1187. Alyse S. Goldberg, Shoba Sujana Kumar, Ellen Greenblatt, Iliana C. Lega, Heather Shapiro. 2018. Delaying Thyroxine Until Positive Beta-Human Chorionic Gonadotropin is Safe for Patients Receiving Fertility Therapy: Applying New ATA Guidelines to Subclinical Hypothyroidism. *Journal of Obstetrics and Gynaecology Canada* **40**:3, 299-303. [[Crossref](#)]
1188. Sima Nazarpour, Fahimeh Ramezani Tehrani, Masoumeh Simbar, Maryam Tohidi, Sonia Minooee, Maryam Rahmati, Fereidoun Azizi. 2018. Effects of Levothyroxine on Pregnant Women With Subclinical Hypothyroidism, Negative for Thyroid Peroxidase Antibodies. *The Journal of Clinical Endocrinology & Metabolism* **103**:3, 926-935. [[Crossref](#)]
1189. Ann M Gronowski. 2018. Evaluation of Thyroid Function during Pregnancy: Have We Taken a Wrong Turn?. *Clinical Chemistry* **64**:3, 439-441. [[Crossref](#)]
1190. Barbara Torlinska, Sarah Bath, Aisha Janjua, Kristien Boelaert, Shiao-Yng Chan. 2018. Iodine Status during Pregnancy in a Region of Mild-to-Moderate Iodine Deficiency is not Associated with Adverse Obstetric Outcomes; Results from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Nutrients* **10**:3, 291. [[Crossref](#)]

1191. van Dijk Myrthe M., Smits Iris H., Fliers Eric, Bisschop Peter H.. 2018. Maternal Thyrotropin Receptor Antibody Concentration and the Risk of Fetal and Neonatal Thyrotoxicosis: A Systematic Review. *Thyroid* **28**:2, 257-264. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
1192. Alexandra Miranda, Nuno Sousa. 2018. Maternal hormonal milieu influence on fetal brain development. *Brain and Behavior* **8**:2, e00920. [[Crossref](#)]
1193. Mariacarla Moleti, Thomas Brix, Francesco Trimarchi, Laszlo Hegedüs. 2018. Screening delle tireopatie gestazionali, trattamento dell'ipotiroidismo subclinico o dell'ipotiroidemia in gravidanza. Siamo ancora in cerca di risposte?. *L'Endocrinologo* **19**:1, 40-42. [[Crossref](#)]
1194. María Martínez, Berta Soldevila, Anna Lucas, Inés Velasco, Lluís Vila, Manel Puig-Domingo. 2018. Hypothyroidism during pregnancy and its association to perinatal and obstetric morbidity: a review. *Endocrinología, Diabetes y Nutrición (English ed.)* **65**:2, 107-113. [[Crossref](#)]
1195. María Martínez, Berta Soldevila, Anna Lucas, Inés Velasco, Lluís Vila, Manel Puig-Domingo. 2018. Hypothyroidism during pregnancy and its association to perinatal and obstetric morbidity: a review. *Endocrinología, Diabetes y Nutrición* **65**:2, 107-113. [[Crossref](#)]
1196. Tim I M Korevaar, Victor J Pop, Layal Chaker, Mariette Goddijn, Yolanda B de Rijke, Peter H Bisschop, Maarten A Broeren, Vincent W V Jaddoe, Marco Medici, Theo J Visser, Eric A P Steegers, Tanja G Vrijkotte, Robin P Peeters. 2018. Dose Dependency and a Functional Cutoff for TPO-Antibody Positivity During Pregnancy. *The Journal of Clinical Endocrinology & Metabolism* **103**:2, 778-789. [[Crossref](#)]
1197. Tahereh Orouji Jokar, Lindsay T Fourman, Hang Lee, Katherine Mentzinger, Pouneh K Fazeli. 2018. Higher TSH Levels Within the Normal Range Are Associated With Unexplained Infertility. *The Journal of Clinical Endocrinology & Metabolism* **103**:2, 632-639. [[Crossref](#)]
1198. Stine Linding Andersen, Stig Andersen, Zeyan Liew, Peter Vestergaard, Jørn Olsen. 2018. Maternal Thyroid Function in Early Pregnancy and Neuropsychological Performance of the Child at 5 Years of Age. *The Journal of Clinical Endocrinology & Metabolism* **103**:2, 660-670. [[Crossref](#)]
1199. Flora Veltri, Pierre Kleynen, Lidia Grabczan, Alexandra Salajan, Serge Rozenberg, Thierry Pepersack, Kris Poppe. 2018. Pregnancy outcomes are not altered by variation in thyroid function within the normal range in women free of thyroid disease. *European Journal of Endocrinology* **178**:2, 189-197. [[Crossref](#)]
1200. Inés Velasco, Peter Taylor. 2018. The role of levothyroxine in obstetric practice. *Annals of Medicine* **50**:1, 57-67. [[Crossref](#)]
1201. Pearce Elizabeth N.. 2018. Levothyroxine Treatment Does Not Reduce Risk for Premature Delivery Among TPO-Negative Pregnant Women with a TSH of 2.5 to 10.0 mIU/L. *Clinical Thyroidology* **30**:1, 7-9. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1202. Chen Angela C., Livhits Masha J., Du Lin, Wu James X., Kuo Eric J., Yeh Michael W., Leung Angela M.. 2018. Recent Pregnancy Is Not Associated with High-Risk Pathological Features of Well-Differentiated Thyroid Cancer. *Thyroid* **28**:1, 68-71. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1203. Giovanni Ceccarini, Ferruccio Santini, Paolo Vitti. Tests of Thyroid Function 33-55. [[Crossref](#)]
1204. Bijay Vaidya, Shiao-Yng Chan. Thyroid Physiology and Thyroid Diseases in Pregnancy 673-708. [[Crossref](#)]
1205. Gesthimani Mintziori, Dimitrios G. Goulis, Basil C. Tarlatzis. Thyroid Function and Pregnancy Outcome After ART: What Is the Evidence? 113-117. [[Crossref](#)]
1206. Veronica Gillispie, Brittany McKinley. Obstetrics: The Hospitalist's Approach to the Pregnant Patient 159-178. [[Crossref](#)]
1207. Mohamad Irani, Samantha M. Pfeifer. Treatment of Subclinical Hypothyroidism in the Infertile Female 115-131. [[Crossref](#)]
1208. Catharina Bullmann. Schilddrüsenfunktionsstörungen in der Reproduktionsmedizin 1-9. [[Crossref](#)]
1209. K. Lorenz, M. Elwerr. Maligne Schilddrüsentumoren 1-34. [[Crossref](#)]
1210. Rajeev Parameswaran, Amit Agarwal. Evidence-Based Surgery 3-10. [[Crossref](#)]
1211. Jacques Orgiazzi. Antithyroid Drugs 683-687. [[Crossref](#)]
1212. Dina M. Abdelsamad, Raymond W. Ke, William H. Kutteh. Thyroid Function and Reproduction 77-81. [[Crossref](#)]
1213. Mark P.J. Vanderpump. Epidemiology of Thyroid Disease 486-495. [[Crossref](#)]
1214. Lise Bernard, Marie Batisse, Igor Tauveron, Brigitte Sallerin, Philippe Caron, Florent Puisset. Traitement des dysthyroïdies 361-377.e1. [[Crossref](#)]
1215. Meera Mallya, Amanda L. Ogilvy-Stuart. 2018. Thyrotropic hormones. *Best Practice & Research Clinical Endocrinology & Metabolism* **32**:1, 17-25. [[Crossref](#)]

1216. Marta Lucas Javato, Inmaculada Domínguez Pascual, Ana Isabel Álvarez Ríos, Manuel Conde Sánchez, Juan Miguel Guerrero Montavez. 2018. Gestante eutiroides con anticuerpos estimulantes del receptor de tirotrópina positivos: manifestación tardía de hipertiroidismo neonatal. A propósito de un caso. *Revista del Laboratorio Clínico* 11:1, 47-50. [[Crossref](#)]
1217. 2018. ACOG Practice Bulletin No. 189: Nausea And Vomiting Of Pregnancy. *Obstetrics & Gynecology* 131:1, e15-e30. [[Crossref](#)]
1218. Won Sang Yoo. 2018. Recurrent Painless Thyroiditis in Patients with History of Postpartum Thyroiditis. *International Journal of Thyroidology* 11:1, 49. [[Crossref](#)]
1219. Flora Veltri, Kris Poppe. 2018. Variables Contributing to Thyroid (Dys)Function in Pregnant Women: More than Thyroid Antibodies?. *European Thyroid Journal* 7:3, 120-128. [[Crossref](#)]
1220. Macy T. Hardley, Andrew H. Chon, Jorge Mestman, Caroline T. Nguyen, Mitchell E. Geffner, Ramen H. Chmait. 2018. Iodine-Induced Fetal Hypothyroidism: Diagnosis and Treatment with Intra-Amniotic Levothyroxine. *Hormone Research in Paediatrics* 90:6, 419-423. [[Crossref](#)]
1221. Leqi He, Chunmei Shen, Yanan Zhang, Zaoping Chen, Heyuan Ding, Jun Liu, Bingbing Zha. 2018. Evaluation of serum ferritin and thyroid function in the second trimester of pregnancy. *Endocrine Journal* 65:1, 75-82. [[Crossref](#)]
1222. Inés Velasco, Peter Taylor. 2018. Identifying and treating subclinical thyroid dysfunction in pregnancy: emerging controversies. *European Journal of Endocrinology* 178:1, D1-D12. [[Crossref](#)]
1223. Pedro Wesley Rosario. 2018. Selective screening for thyroid dysfunction in pregnant women: How often do low-risk women cease to be treated following the new guidelines of the American Thyroid Association?. *Archives of Endocrinology and Metabolism* 62:6, 641-643. [[Crossref](#)]
1224. Abby Ershow, Sheila Skeaff, Joyce Merkel, Pamela Pehrsson. 2018. Development of Databases on Iodine in Foods and Dietary Supplements. *Nutrients* 10:1, 100. [[Crossref](#)]
1225. Tae Yong Kim. 2018. Thyroid-Stimulating Hormone Reference Ranges in Early Pregnancy: Possible Influence of Iodine Status. *Endocrinology and Metabolism* 33:4, 445. [[Crossref](#)]
1226. Carmen Castillo, Nicole Lustig, Paula Margozzini, Andrea Gomez, María Paulina Rojas, Santiago Muzzo, Lorena Mosso. 2018. Thyroid-Stimulating Hormone Reference Ranges in the First Trimester of Pregnancy in an Iodine-Sufficient Country. *Endocrinology and Metabolism* 33:4, 466. [[Crossref](#)]
1227. Sanjay Kalra, Sameer Agarwal, Rashmi Aggarwal, Salam Ranabir. 2018. Trimester-specific thyroid-stimulating hormone: An Indian perspective. *Indian Journal of Endocrinology and Metabolism* 22:1, 1. [[Crossref](#)]
1228. Subramanian Kannan, Shriraam Mahadevan, Alben Sigamani. 2018. A systematic review on normative values of trimester-specific thyroid function tests in Indian women. *Indian Journal of Endocrinology and Metabolism* 22:1, 7. [[Crossref](#)]
1229. Subramanian Kannan, Sanjay Kalra. 2018. Normative Trimester-Specific Thyroid Function Data from India: The State of the Nation. *Indian Journal of Endocrinology and Metabolism* 22:1, 5. [[Crossref](#)]
1230. Jee Hee Yoon, Ho-Cheol Kang. 2018. Interpretation of puzzling thyroid function tests. *Journal of the Korean Medical Association* 61:4, 241. [[Crossref](#)]
1231. Chang Hoon Yim. 2018. Thyroid dysfunction during pregnancy. *Journal of the Korean Medical Association* 61:4, 253. [[Crossref](#)]
1232. Darya E. Soboleva, Svetlana V. Dora, Tatyana L. Karonova, Anna R. Volkova, Elena N. Grineva. 2017. Iodine status of Saint Petersburg adult residents. *Clinical and experimental thyroidology* 13:4, 23-29. [[Crossref](#)]
1233. Rudolf Hoermann, John E. M. Midgley, Rolf Larisch, Johannes W. Dietrich. 2017. Recent Advances in Thyroid Hormone Regulation: Toward a New Paradigm for Optimal Diagnosis and Treatment. *Frontiers in Endocrinology* 8. . [[Crossref](#)]
1234. Kapoor Ekta, Faubion Stephanie, Hines Stephanie, Stuenkel Cynthia A.. 2017. Women's Health Endocrine Update. *Journal of Women's Health* 26:12, 1365-1368. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1235. Lee Sun Y., McCarthy Alicia M., Stohl Hindi, Ibrahim Sherrine, Jeong Christina, Braverman Lewis E., Ma Wendy, He Xuemei, Mestman Jorge H., Schuller Kristin E., Jahreis Katherine A., Pearce Elizabeth N., Leung Angela M.. 2017. Urinary Iodine, Perchlorate, and Thiocyanate Concentrations in U.S. Lactating Women. *Thyroid* 27:12, 1574-1581. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1236. Sofie Bliddal, Ulla Feldt-Rasmussen. 2017. Is Thyroid Replacement Therapy Necessary in Fertility Patients with "Normal" TSH?. *Journal of Fetal Medicine* 4:4, 171-176. [[Crossref](#)]
1237. Urszula Ambroziak, Sandra Hybsier, Urszula Shahnazaryan, Małgorzata Krasnodębska-Kiljańska, Eddy Rijntjes, Zbigniew Bartoszewicz, Tomasz Bednarczuk, Lutz Schomburg. 2017. Severe selenium deficits in pregnant women irrespective of autoimmune thyroid disease in an area with marginal selenium intake. *Journal of Trace Elements in Medicine and Biology* 44, 186-191. [[Crossref](#)]

1238. Catherine Ley, Lauren Pischel, Julie Parsonnet. 2017. Triclosan and triclocarban exposure and thyroid function during pregnancy —A randomized intervention. *Reproductive Toxicology* **74**, 143-149. [[Crossref](#)]
1239. VJ Pop, MA Broeren, WM Wiersinga, A Stagnaro-Green. 2017. Thyroid disease symptoms during early pregnancy do not identify women with thyroid hypofunction that should be treated. *Clinical Endocrinology* **87**:6, 838-843. [[Crossref](#)]
1240. María Castillo Lara, Ángel Vilar Sánchez, Consuelo Cañavate Solano, Estefanía Soto Pazos, María Iglesias Álvarez, Carmen González Macías, Carmen Ayala Ortega, Luis Javier Moreno Corral, Juan Jesús Fernández Alba. 2017. "Hypothyroidism screening during first trimester of pregnancy". *BMC Pregnancy and Childbirth* **17**:1. . [[Crossref](#)]
1241. YunYing Cai, LanPing Zhong, Jie Guan, RuiJin Guo, Ben Niu, YanPing Ma, Heng Su. 2017. Outcome of in vitro fertilization in women with subclinical hypothyroidism. *Reproductive Biology and Endocrinology* **15**:1. . [[Crossref](#)]
1242. Naoko Arata. 2017. Management of Endocrine Disorders during Pregnancy. *Nihon Naika Gakkai Zasshi* **106**:11, 2432-2438. [[Crossref](#)]
1243. Spaulding Stephen W.. 2017. Does Antithyroid Drug Treatment Improve Outcomes in Wild-Type Offspring of Mothers with Thyroid Hormone Resistance?. *Clinical Thyroidology* **29**:11, 426-429. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1244. Pearce Elizabeth N.. 2017. TPO Antibody Positivity in Pregnancy Is Associated with Metabolic Syndrome, Increased Waist Circumference, and Higher BMI in Offspring at Age 16. *Clinical Thyroidology* **29**:11, 430-432. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1245. Meredith M. Howley, Sarah C. Fisher, Alissa R. Van Zutphen, Dorothy K. Waller, Suzan L. Carmichael, Marilyn L. Browne. 2017. Thyroid Medication Use and Birth Defects in the National Birth Defects Prevention Study. *Birth Defects Research* **109**:18, 1471-1481. [[Crossref](#)]
1246. Sarah C Bath. 2017. Iodine supplementation in pregnancy in mildly deficient regions. *The Lancet Diabetes & Endocrinology* **5**:11, 840-841. [[Crossref](#)]
1247. Sueppong Gowachirapant, Nidhi Jaiswal, Alida Melse-Boonstra, Valeria Galetti, Sara Stinca, Ian Mackenzie, Susan Thomas, Tinku Thomas, Pattanee Winichagoon, Krishnamachari Srinivasan, Michael B Zimmermann. 2017. Effect of iodine supplementation in pregnant women on child neurodevelopment: a randomised, double-blind, placebo-controlled trial. *The Lancet Diabetes & Endocrinology* **5**:11, 853-863. [[Crossref](#)]
1248. Aimee Seungdamrong, Anne Z. Steiner, Clarisa R. Gracia, Richard S. Legro, Michael P. Diamond, Christos Coutifaris, William D. Schlaff, Peter Casson, Gregory M. Christman, Randal D. Robinson, Hao Huang, Ruben Alvero, Karl R. Hansen, Susan Jin, Esther Eisenberg, Heping Zhang, Nanette Santoro. 2017. Preconceptional antithyroid peroxidase antibodies, but not thyroid-stimulating hormone, are associated with decreased live birth rates in infertile women. *Fertility and Sterility* **108**:5, 843-850. [[Crossref](#)]
1249. Anna-Leena Heikkinen, Fanni Pääkilä, Anna-Liisa Hartikainen, Marja Vääräsmäki, Tuija Männistö, Eila Suvanto. 2017. Maternal Thyroid Antibodies Associates With Cardiometabolic Risk Factors in Children at the Age of 16. *The Journal of Clinical Endocrinology & Metabolism* **102**:11, 4184-4190. [[Crossref](#)]
1250. Azizi Fereidoun, Malboosbaf Ramin. 2017. Long-Term Antithyroid Drug Treatment: A Systematic Review and Meta-Analysis. *Thyroid* **27**:10, 1223-1231. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1251. Thomas Stulnig. 2017. Schilddrüse und weibliche Fertilität. *Journal für Gynäkologische Endokrinologie/Österreich* **27**:3, 102-107. [[Crossref](#)]
1252. Thomas Stulnig. 2017. Schilddrüse und weibliche Fertilität. *Journal für Gynäkologische Endokrinologie/Schweiz* **20**:3, 109-114. [[Crossref](#)]
1253. Rainbow Rubin, Michelle Pearl, Martin Kharrazi, Benjamin C. Blount, Mark D. Miller, Elizabeth N. Pearce, Liza Valentin-Blasini, Gerald DeLorenze, Jane Liaw, Andrew N. Hoofnagle, Craig Steinmaus. 2017. Maternal perchlorate exposure in pregnancy and altered birth outcomes. *Environmental Research* **158**, 72-81. [[Crossref](#)]
1254. Mahboubeh Ghanbari, Asghar Ghasemi. 2017. Maternal hypothyroidism: An overview of current experimental models. *Life Sciences* **187**, 1-8. [[Crossref](#)]
1255. Tim I. M. Korevaar, Marco Medici, Theo J. Visser, Robin P. Peeters. 2017. Thyroid disease in pregnancy: new insights in diagnosis and clinical management. *Nature Reviews Endocrinology* **13**:10, 610-622. [[Crossref](#)]
1256. Stine L. Andersen. 2017. Risk of embryopathies with use of antithyroidal medications. *Current Opinion in Endocrinology & Diabetes and Obesity* **24**:5, 364-371. [[Crossref](#)]
1257. Katsuhiko Hamada, Tetsuya Mizokami, Tetsushi Maruta, Kiichiro Higashi, Kaoru Konishi, Naoko Momotani, Junichi Tajiri. 2017. Effects of Inorganic Iodine Therapy Administered to Lactating Mothers With Graves Disease on Infant Thyroid Function. *Journal of the Endocrine Society* **1**:10, 1293-1300. [[Crossref](#)]

1258. Stine Linding Andersen, Stefan Lönn, Peter Vestergaard, Ove Törring. 2017. Birth defects after use of antithyroid drugs in early pregnancy: a Swedish nationwide study. *European Journal of Endocrinology* **177**:4, 369-378. [[Crossref](#)]
1259. Pearce Elizabeth N.. 2017. Hypothyroid Symptoms in Pregnant Women Fail to Predict Hypothyroid Status. *Clinical Thyroidology* **29**:9, 332-334. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1260. Chen Chien-Wen, Huang Ya-Li, Tzeng Chii-Ruey, Huang Rui-Lan, Chen Chi-Huang. 2017. Idiopathic Low Ovarian Reserve Is Associated with More Frequent Positive Thyroid Peroxidase Antibodies. *Thyroid* **27**:9, 1194-1200. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1261. Anagnostis Panagiotis, Lefkou Eleftheria, Goulis Dimitrios G.. 2017. Re: "Guidelines of the American Thyroid Association for the Diagnosis and Management of Thyroid Disease During Pregnancy and the Postpartum" by Alexander et al. (Thyroid 2017;27:315-389). *Thyroid* **27**:9, 1209-1210. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1262. Atanaska Elenkova, Iliana Atanasova, Georgi Kirilov, Emil Natchev, Ralitz Ivanova, Roussanka Kovatcheva, Silvia Vandeva, Dimitar Tcharaktchiev, Sabina Zacharieva. 2017. Autoimmune hypothyroidism is three times more frequent in female prolactinoma patients compared to healthy women: data from a cross-sectional case-control study. *Endocrine* **57**:3, 486-493. [[Crossref](#)]
1263. Abha Maheshwari, Priya Bhide, Jyotsna Pundir, Siladitya Bhattacharya. 2017. Routine serum thyroid-stimulating hormone testing —optimizing pre-conception health or generating toxic knowledge?. *Human Reproduction* **32**:9, 1779-1785. [[Crossref](#)]
1264. Tim I M Korevaar, Eric A P Steegers, Laya Chaker, Marco Medici, Vincent W V Jaddoe, Theo J Visser, Yolanda B de Rijke, Robin P Peeters. 2017. Thyroid Function and Premature Delivery in TPO Antibody-Negative Women: The Added Value of hCG. *The Journal of Clinical Endocrinology & Metabolism* **102**:9, 3360-3367. [[Crossref](#)]
1265. Shannon D Sullivan, Erin Downs, Geanina Popoveniuc, Alexander Zeymo, Jacqueline Jonklaas, Kenneth D Burman. 2017. Randomized Trial Comparing Two Algorithms for Levothyroxine Dose Adjustment in Pregnant Women With Primary Hypothyroidism. *The Journal of Clinical Endocrinology & Metabolism* **102**:9, 3499-3507. [[Crossref](#)]
1266. Lee Sun Y., Stagnaro-Green Alex, MacKay Douglas, Wong Andrea W., Pearce Elizabeth N.. 2017. Iodine Contents in Prenatal Vitamins in the United States. *Thyroid* **27**:8, 1101-1102. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1267. Jackie Gilbert. 2017. Hypothyroidism. *Medicine* **45**:8, 506-509. [[Crossref](#)]
1268. Alex Stagnaro-Green. 2017. Second trimester levothyroxine treatment for subclinical hypothyroidism or hypothyroxinaemia of pregnancy does not improve cognitive outcomes of children. *Evidence Based Medicine* **22**:4, 149-149. [[Crossref](#)]
1269. Apostolos Chatzitomaris, Rudolf Hoermann, John E. Midgley, Steffen Hering, Aline Urban, Barbara Dietrich, Assjana Abood, Harald H. Klein, Johannes W. Dietrich. 2017. Thyroid Allostasis-Adaptive Responses of Thyrotropic Feedback Control to Conditions of Strain, Stress, and Developmental Programming. *Frontiers in Endocrinology* **8**. . [[Crossref](#)]
1270. Flavia Di Bari, Roberta Granese, Maria Le Donne, Roberto Vita, Salvatore Benvenega. 2017. Autoimmune Abnormalities of Postpartum Thyroid Diseases. *Frontiers in Endocrinology* **8**. . [[Crossref](#)]
1271. Ishrat Khan, Onyebuchi Okosieme, John Lazarus. 2017. Antithyroid drug therapy in pregnancy: a review of guideline recommendations. *Expert Review of Endocrinology & Metabolism* **12**:4, 269-278. [[Crossref](#)]
1272. Anderson Philip O.. 2017. Choosing Medication Alternatives During Breastfeeding, Avoiding Alternative Facts. *Breastfeeding Medicine* **12**:6, 328-330. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1273. D. Führer, W. A. Scherbaum. 2017. Neues zur Endokrinologie. *Der Diabetologe* **13**:5, 342-354. [[Crossref](#)]
1274. Amanda L. Ogilvy-Stuart. 2017. Neonatal Thyrotoxicosis. *NeoReviews* **18**:7, e422-e430. [[Crossref](#)]
1275. Ines Bucci, Cesidio Giuliani, Giorgio Napolitano. 2017. Thyroid-Stimulating Hormone Receptor Antibodies in Pregnancy: Clinical Relevance. *Frontiers in Endocrinology* **8**. . [[Crossref](#)]
1276. María C. Opazo, Pablo A. González, Betsi D. Flores, Luis F. Venegas, Eduardo A. Albornoz, Pablo Cisternas, Karen Bohmwald, Pamela A. Nieto, Susan M. Bueno, Alexis M. Kalergis, Claudia A. Riedel. 2017. Gestational Hypothyroxinemia Imprints a Switch in the Capacity of Astrocytes and Microglial Cells of the Offspring to React in Inflammation. *Molecular Neurobiology* **22**. . [[Crossref](#)]
1277. Cristina Modesti, Paola Aceto, Lucia Masini, Celestino Pio Lombardi, Rocco Bellantone, Liliana Sollazzi. 2017. Approach to thyroid carcinoma in pregnancy. *Updates in Surgery* **69**:2, 261-265. [[Crossref](#)]
1278. Mariacarla Moleti, Francesco Trimarchi, Francesco Vermiglio. 2017. Quando e che cosa deve comunicare al neonatologo/pediatra, l'endocrinologo che segue una gestante con problemi tiroidei?. *L'Endocrinologo* **18**:S1, 59-60. [[Crossref](#)]
1279. Roy Moncayo, Helga Moncayo. 2017. A post-publication analysis of the idealized upper reference value of 2.5 mIU/L for TSH: Time to support the thyroid axis with magnesium and iron especially in the setting of reproduction medicine. *BBA Clinical* **7**, 115-119. [[Crossref](#)]

1280. Shi Chen, Xiang Zhou, Huijuan Zhu, Hongbo Yang, Fengying Gong, Linjie Wang, Man Zhang, Yu Jiang, Chengsheng Yan, Jianqiang Li, Qing Wang, Shikun Zhang, Hui Pan. 2017. Preconception TSH and pregnancy outcomes: a population-based cohort study in 184 611 women. *Clinical Endocrinology* **86**:6, 816-824. [[Crossref](#)]
1281. Flora Veltri, Julie Belhomme, Pierre Kleynen, Lidia Grabczan, Serge Rozenberg, Thierry Pepersack, Kris Poppe. 2017. Maternal thyroid parameters in pregnant women with different ethnic backgrounds: Do ethnicity-specific reference ranges improve the diagnosis of subclinical hypothyroidism?. *Clinical Endocrinology* **86**:6, 830-836. [[Crossref](#)]
1282. Maïa Banigé, Candice Estellat, Valerie Biran, Luc Desfrere, Valerie Champion, Alexandra Benachi, Yves Ville, Marc Dommergues, Pierre-Henri Jarreau, Mostafa Mokhtari, Claire Boithias, Frederic Brioude, Laurent Mandelbrot, Pierre-François Ceccaldi, Delphine Mitanchez, Michel Polak, Dominique Luton. 2017. Study of the Factors Leading to Fetal and Neonatal Dysthyroidism in Children of Patients With Graves Disease. *Journal of the Endocrine Society* **1**:6, 751-761. [[Crossref](#)]
1283. Maria Le Donne, Carmela Mento, Salvatore Settineri, Alessandro Antonelli, Salvatore Benvenga. 2017. Postpartum Mood Disorders and Thyroid Autoimmunity. *Frontiers in Endocrinology* **8**. . [[Crossref](#)]
1284. Pearce Elizabeth N.. 2017. Higher Preconception TSH Values Are Associated With Adverse Obstetric Outcomes. *Clinical Thyroidology* **29**:5, 189-191. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1285. Asha Hesarghatta Shyamasunder, Prakash Abraham. 2017. Measuring TSH receptor antibody to influence treatment choices in Graves' disease. *Clinical Endocrinology* **86**:5, 652-657. [[Crossref](#)]
1286. Sofie Bliddal, Malene Boas, Linda Hilsted, Lennart Friis-Hansen, Anders Juul, Torben Larsen, Ann Tabor, Jens Faber, Dorthe Hansen Precht, Ulla Feldt-Rasmussen. 2017. Increase in thyroglobulin antibody and thyroid peroxidase antibody levels, but not preterm birth-rate, in pregnant Danish women upon iodine fortification. *European Journal of Endocrinology* **176**:5, 603-612. [[Crossref](#)]
1287. Mary Barna Bridgeman. Thyroid Disorders: Hyper- and Hypothyroidism . [[Crossref](#)]
1288. Hershman Jerome M.. 2017. Levothyroxine Therapy of Subclinical Hypothyroidism or Hypothyroxinemia in Pregnancy Does Not Improve Cognitive Function in the Offspring. *Clinical Thyroidology* **29**:4, 132-135. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1289. Leung Angela M.. 2017. Thyroid Nodules Do Not Increase in Size During Pregnancy. *Clinical Thyroidology* **29**:4, 150-152. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1290. Alex Stagnaro-Green. 2017. Thyroid and pregnancy — time for universal screening?. *Nature Reviews Endocrinology* **13**:4, 192-194. [[Crossref](#)]
1291. Dianjun Sun, Karen Codling, Suying Chang, Shubin Zhang, Hongmei Shen, Xiaohui Su, Zupei Chen, Robert W. Scherpbier, Jun Yan. 2017. Eliminating Iodine Deficiency in China: Achievements, Challenges and Global Implications. *Nutrients* **9**:4, 361. [[Crossref](#)]
1292. Korevaar Tim I.M.. 2017. Evidence-Based Tightrope Walking: The 2017 Guidelines of the American Thyroid Association for the Diagnosis and Management of Thyroid Disease During Pregnancy and the Postpartum. *Thyroid* **27**:3, 309-311. [[Citation](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
1293. Giovanni Ceccarini, Ferruccio Santini, Paolo Vitti. Tests of Thyroid Function 1-23. [[Crossref](#)]
1294. Bijay Vaidya, Shiao-Yng Chan. Thyroid Physiology and Thyroid Diseases in Pregnancy 1-36. [[Crossref](#)]
1295. Doina Piciu. The Radiation Protection 21-29. [[Crossref](#)]
1296. Doina Piciu, Patriciu Achimaş-Cadariu, Alexandru Irimie, Andra Piciu. The Thyroid Gland 61-177. [[Crossref](#)]
1297. Hyung Wook Choi, You Jung Han, Dong Wook Kwak, So Young Park, Sung Hoon Kim, Hyun Koo Yoon, Chang Hoon Yim. 2017. Maternal Thyroid Function during the First Trimester of Pregnancy in Korean Women. *International Journal of Thyroidology* **10**:1, 36. [[Crossref](#)]
1298. N. V. Shidlovskaya, V. A. Petrukhin, F. F. Burumkulova, N. F. Bashakin, I. V. Barinova. 2017. Thyroid cancer and pregnancy. Features of course and perinatal outcomes. *Rossiiskii vestnik akushera-ginekologa* **17**:6, 83. [[Crossref](#)]
1299. Massimo Giusti, Lucia Conte, Anna Maria Repetto, Stefano Gay, Paola Marroni, Miranda Mittica, Michele Mussap. 2017. Detection of Polyethylene Glycol Thyrotropin (TSH) Precipitable Percentage (Macro-TSH) in Patients with a History of Thyroid Cancer. *Endocrinology and Metabolism* **32**:4, 460. [[Crossref](#)]
1300. Gin Chiu, Xiaotian Zhang, Enfa Zhao, Baomin Liu. 2017. Maternal Thyroid Disease and Neonatal Low Birth Weight: A Systematic Review and Meta-Analysis. *Open Journal of Obstetrics and Gynecology* **07**:07, 778-789. [[Crossref](#)]